



HAL
open science

Bleomycin exerts ambivalent antitumor immune effect by triggering both immunogenic cell death and proliferation of regulatory T cells.

Hélène Bugaut, Mélanie Bruchard, Hélène Berger, Valentin Derangère, Ludivine Odoul, Romain Euvrard, Sylvain Ladoire, Fanny Chalmin, Frédérique Végran, Cédric Rébé, et al.

► To cite this version:

Hélène Bugaut, Mélanie Bruchard, Hélène Berger, Valentin Derangère, Ludivine Odoul, et al.. Bleomycin exerts ambivalent antitumor immune effect by triggering both immunogenic cell death and proliferation of regulatory T cells.. PLoS ONE, 2013, 8 (6), pp.e65181. 10.1371/journal.pone.0065181 . inserm-00857882

HAL Id: inserm-00857882

<https://inserm.hal.science/inserm-00857882>

Submitted on 4 Sep 2013

HAL is a multi-disciplinary open access archive for the deposit and dissemination of scientific research documents, whether they are published or not. The documents may come from teaching and research institutions in France or abroad, or from public or private research centers.

L'archive ouverte pluridisciplinaire **HAL**, est destinée au dépôt et à la diffusion de documents scientifiques de niveau recherche, publiés ou non, émanant des établissements d'enseignement et de recherche français ou étrangers, des laboratoires publics ou privés.

Bleomycin Exerts Ambivalent Antitumor Immune Effect by Triggering Both Immunogenic Cell Death and Proliferation of Regulatory T Cells

Hélène Bugaut^{1,2,3}, Mélanie Bruchard^{1,2,3}, Hélène Berger^{1,2,3}, Valentin Derangère^{1,2,3}, Ludivine Odoul⁴, Romain Euvrard^{1,2,3}, Sylvain Ladoire^{1,2,3,5}, Fanny Chalmin^{1,2,3}, Frédérique Végran^{1,2,3}, Cédric Rébé^{1,2,3,5}, Lionel Apetoh^{1,2,3}, François Ghiringhelli^{1,2,3,5*}, Grégoire Mignot^{1,2,3*}

1 INSERM, U866, Dijon, France, 2 INSERM AVENIR Team, Dijon, France, 3 Faculté de Médecine, Université de Bourgogne, Dijon, France, 4 University of Bern, Institute of Pathology, Bern, Switzerland, 5 Centre Georges François Leclerc, Dijon, France

Abstract

Bleomycin (BLM) is an anticancer drug currently used for the treatment of testis cancer and Hodgkin lymphoma. This drug triggers cancer cell death via its capacity to generate radical oxygen species (ROS). However, the putative contribution of anticancer immune responses to the efficacy of BLM has not been evaluated. We make here the observation that BLM induces immunogenic cell death. In particular, BLM is able to induce ROS-mediated reticulum stress and autophagy, which result in the surface exposure of chaperones, including calreticulin and ERp57, and liberation of HMBG1 and ATP. BLM induces anti-tumor immunity which relies on calreticulin, CD8⁺ T cells and interferon- γ . We also find that, in addition to its capacity to trigger immunogenic cell death, BLM induces expansion of Foxp3⁺ regulatory T (Treg) cells via its capacity to induce transforming growth factor beta (TGF β) secretion by tumor cells. Accordingly, Treg cells or TGF β depletion dramatically potentiates the antitumor effect of BLM. We conclude that BLM induces both anti-tumor CD8⁺ T cell response and a counteracting Treg proliferation. In the future, TGF β or Treg inhibition during BLM treatment could greatly enhance BLM anti-tumor efficacy.

Citation: Bugaut H, Bruchard M, Berger H, Derangère V, Odoul L, et al. (2013) Bleomycin Exerts Ambivalent Antitumor Immune Effect by Triggering Both Immunogenic Cell Death and Proliferation of Regulatory T Cells. PLoS ONE 8(6): e65181. doi:10.1371/journal.pone.0065181

Editor: John W. Glod, Robert Wood Johnson Medical School, United States of America

Received: January 10, 2013; **Accepted:** April 23, 2013; **Published:** June 7, 2013

Copyright: © 2013 Bugaut et al. This is an open-access article distributed under the terms of the Creative Commons Attribution License, which permits unrestricted use, distribution, and reproduction in any medium, provided the original author and source are credited.

Funding: This work was supported by special grants from the Fondation de France, the Association pour la Recherche sur le Cancer, the Institut National de la Santé et de la Recherche Médicale, the Ligue Contre le Cancer (Région Bourgogne), the Fondation pour la Recherche Médicale, and the Institut National du Cancer. GM was funded by the Association pour la Recherche sur le Cancer and FV by the Ligue Nationale contre le cancer. LA was supported by Agence Nationale pour la Recherche [ANR-10-PDOC-014-01]. HB is funded by École de l'Inserm Liliane Bettencourt. The funders had no role in study design, data collection and analysis, decision to publish, or preparation of the manuscript.

Competing Interests: The authors have declared that no competing interests exist.

* E-mail: fghiringhelli@cgfl.fr (FG); gregoire.mignot@free.fr (GM)

☯ These authors contributed equally to this work.

Introduction

The aim of anti-cancer therapy is the eradication of tumor cells in the patient's body, including the smallest metastasis or single-cell localization that could not be removed by surgery. Chemotherapy and radiotherapy are currently used to complement surgery to kill residual disease or for the treatment of metastatic disease. However, tumor cells can develop some resistance and escape cytotoxic treatment, causing tumor relapse. In this context, immunotherapy is seen as one of the ultimate anti-cancer strategies, as the immune system could reshape its actions against an evolving malignancy and eliminate tumor cells resistant to chemotherapy. Accumulating evidence shows that conventional chemotherapies, in addition to their direct cytotoxic effect, could trigger an antitumor immune response. In particular, we have shown that some chemotherapies activate natural killer (NK) cells [1,2], some drugs target suppressive cells such as regulatory T cells (Treg) or myeloid-derived suppressor cells (MDSC) [3,4] and some other drugs also induce a particular mode of cancer cell death called "immunogenic cell death" (ICD) [5]. ICD relies on the

capacity of drugs to induce three main checkpoints. The first one is the exposure of 'eat-me' signals on cell surface, such as calreticulin (CRT), caused by endoplasmic reticulum (RE) stress [6], which favors cancer cell phagocytosis by dendritic cells [7]. The second is the secretion of an endogenous Toll-like receptor 4 (TLR4) ligand High-mobility group box 1 (HMGB1) [8], which is required for efficient processing of tumor antigen by dendritic cells. The third signal is the release of ATP which induces the production of interleukin (IL)-1 β by dendritic cells and favors CD8⁺ T cell immune response [9].

Bleomycin (BLM) is an anti-tumor antibiotic glycopeptide produced by the bacterium *Streptomyces* [10]. BLM causes breaks in DNA, similar to those obtained with radiotherapy [11]. This DNA damage has been demonstrated to be mediated through induction of oxidative stress [12,13]. BLM is indicated, in association with other cytotoxic agents, for the treatment of cancer testis and Hodgkin disease [14,15]. The particularity of these two diseases is the high rate of cure obtained by chemotherapy. So, we proposed to test the antitumor immune response induced by BLM to determine if this mechanism could participate in the antitumor

effect of bleomycin and may explain the high capacity of BLM-containing regimens to cure cancer.

In this study, we analyzed the implication of ICD, effector cells (such as CD8⁺ T and NK cells), as well as modulator cells (such as dendritic cells, MDSC and Treg) in BLM antitumor efficacy. We describe how BLM treatment induces an immunogenic apoptosis of tumor cells, anti-tumor CD8⁺ T cell response and the production of the tolerogenic cytokine transforming growth factor beta (TGF β) by tumor cells which mediates regulatory T cell accumulation *in vivo*.

Materials and Methods

Cell Culture

MCA205 (fibrosarcoma), B16F10 (melanoma) cell lines, syngeneic of C57BL/6 mice, and CT26 (colon carcinoma) cell line, syngeneic of Balb/c mice, were cultured at 37°C in 5% CO₂ in RPMI 1640 (Lonza) enriched with 10% fetal bovine serum, 0.4 mmol/L of sodium pyruvate, 4 mmol/L of HEPES, and antibiotics (penicillin, streptomycin, and amphotericin B). All cell lines were obtained from the American Type Culture Collection, excepted for B16-OVA, which was a kind gift from Pr Laurence Zitvogel.

Mouse Models

Mice. All experiments were approved by the Université de Bourgogne's Animal Experimental Ethics Comitee. Female C57BL/6 or Balb/c mice were purchased from the Centre d'Élevage Janvier (Le Genest St. Isle, France) and used between 6 and 10 weeks of age. OT-I transgenic mice (C57BL/6-Tg(TCR α /TCR β)1100mjb) were provided by C.Borg (INSERM U645, Besançon, France). Foxp3-EGFP [16] and CD45.1 syngeneic mice were from CDTA (UMR 7355 CNRS, Orléans, France).

Tumor models. CT26 tumor cells were injected into the right flank of Balb/c mice (1.10⁶ cells). Tumor size was monitored three times a week with a caliper. In some experiments, 1.10⁶ cells were injected intraperitoneally to induce peritoneal carcinosis.

Treatments. Mice treated by BLM received 5 intraperitoneal injections (20 mg/kg) every other day, starting when tumor size reached approximately 25 mm². Mice receiving LPS (InVivoGen, Toulouse, France) were treated intraperitoneally (40 μ g) the day before sacrifice. Mice treated with IL-2 received 2 intraperitoneal injections per day (1.10⁵ UI, Chiron), starting two days before sacrifice. Anti-CD8 (2.43), anti-CD4 (GK1.5), anti-CD25 (PC-61.5.3) and anti-TGF β (1D11) antibodies were bought from BioXcell (West Lebanon, NH, USA) and used to eliminate CD8⁺, CD4⁺, CD25⁺ Treg lymphocytes or TGF β *in vivo*, respectively. Injections were performed intraperitoneally, 150 μ g per mouse 12 hours before and 12 hours after first BLM injection, and then 100 μ g per mouse twice a week.

Vaccination. B16F10 cells expressing chicken ovalbumin (B16-OVA) were treated for 24 hours using the indicated drugs, and were then injected (5.10⁵ cells) into the footpad of C57BL/6 mice. Positive control was 1 mg of ovalbumin (Calbiochem, Merck France) protein plus 10 μ g CpG ODN 1668 (InVivoGen). Five days later, draining lymph nodes were harvested, mechanically disrupted and restimulated for 72 hours with PBS, OVA (1 g/L) or anti-CD3 (2 μ g/mL, 145.2C11, BioXcell) plus anti-CD28 (2 μ g/mL, PV1, BioXcell). Then supernatants were collected and IFN γ concentration was determined using ELISA (BD biosciences, le Pont de Claix, France).

In vitro Treatments

Tumor cells were treated for 24 hours with 30 μ M Mitomycin C, 0.25 μ M doxorubicin, 30 mM N-acetyl-cystein, or 15 μ g/mL BLM for CT26 and MCA205 cells, 150 μ g/mL BLM for B16F10.

Suppression assays. We used OT-I suppression assay [4,17,18], adapted as follow. Five days after PBS or BLM treatment, spleens from tumor bearing mice were harvested and mechanically disrupted. MDSC were labeled as PE-Cy7 Gr1⁺ cells and isolated with anti-PE-Cy7 magnetic beads (Miltenyi). CD4⁺ lymphocytes were purified using MACS CD4⁺ T cells isolation kit II (Miltenyi, Paris, France), and then Treg were purified among CD4⁺ lymphocytes, using CD25 microbeads (Miltenyi): CD4⁺ CD25⁺ were labelled Treg and CD4⁺ CD25⁻ cells were used as Tconv. Separately, OT-I transgenic mice were sacrificed and lymph nodes and spleens harvested. CD8⁺ lymphocytes were then purified using MACS CD8⁺ T cells isolation kit II. CD8⁺ OT-I lymphocytes were cocultured for 72 hours with SIINFEKL peptide (10 μ g/mL, Tebu) and purified Treg or Tconv (1:1 cells) from tumor bearing mice. IFN γ concentration in the supernatants was determined by ELISA (BD bioscience).

Cell transfer. CD45.2 Foxp3-EGFP mice were sacrificed then lymph nodes and spleens were harvested. These organs were mechanically disrupted and the single cell suspension was enriched in CD4⁺ lymphocytes using MACS CD4⁺ T cells isolation kit II (Miltenyi). Then CD4⁺ GFP⁻ CD62L⁺ cells (naïve CD4⁺ T lymphocytes) and CD4⁺ GFP⁺ cells (Treg) were sorted using FACS Aria III (BD). *In vivo*, sorted naïve T cells and Treg cells were injected intravenously into CD45.1 C57BL/6 CT26 tumor bearing mice. CD45.1 mice which had received Treg were then treated with PBS, BLM or IL-2. CD45.1 mice which had received naïve CD4⁺ T cells were treated with PBS or BLM. The day before sacrifice, CD45.1 mice which had received Treg were treated by EdU (70 mg/kg, intraperitoneal). Five days after BLM treatment, all CD45.1 mice were sacrificed and spleens and tumors harvested. Organs were mechanically disrupted, and a CD4 CD45.2 Foxp3 EdU (Click-iT EdU kit, Invitrogen, Saint Aubin, France) staining was performed. Cells were then analysed by flow cytometry. *In vitro*, Treg were cultured with supernatant from PBS-or BLM-treated CT26 cells in presence of anti-CD3 and anti-CD28 antibodies. Three days later, cells were incubated 3 h with BrdU, then handled with BD BrdU Flow Kit, according to the manufacturer's protocol and FACS-analysed.

Flow Cytometry

Five days after last BLM treatment, tumor bearing mice were sacrificed and organs were harvested, and mechanically disrupted. The cell suspensions from spleen or tumor were separated with lymphocyte separation medium (Eurobio) to isolate live mononuclear cells. Cells were stained with the antibodies described below, according to the manufacturer's protocol. Foxp3 staining was performed using mouse regulatory T cell staining kit (eBiosciences). Dead cells were excluded by DAPI labeling (1 μ g/mL). Following antibodies were from BD Biosciences : APC-H7 anti-CD19, PerCP anti-CD3, V500 anti-CD4, V450 anti-CD8, PE anti-DX5, FITC anti-CD11c, PE anti-Ly6G, PerCP anti-CD4, APC anti-CD3, FITC anti-CD40, PE anti-CD86, Alexa700 anti-CD62L, V450 anti-CD45.2, APC anti-CD11c.; the following antibodies were from BioLegend: Alexa700 anti-F4/80, APC anti-LAP; the following antibodies were from eBiosciences: APC anti-CD11b, PerCP-Cy5.5 anti-Ly6C, eFluor450 anti-Foxp3, Alexa700 anti-CHM-II. Other antibodies were also used: Alexa647 azide (Invitrogen), anti-CRT rabbit whole serum (Abcam), anti-ERp57 rabbit whole serum (Abcam), Alexa 680 anti-rabbit (Invitrogen). Data were acquired with LSR II

(BD), and analysed with FlowJo (Tristar). Cell sorting was done with a BD ARIA III. Cell populations were defined as below: B lymphocytes: CD19⁺; CD4⁺ and CD8⁺ T lymphocytes: CD3⁺ CD4⁺ or CD3⁺ CD8⁺, respectively; NK lymphocytes: CD19⁻ CD3⁻ DX5⁺; dendritic cells: CD19⁻ CD3⁻ CD11c high; monocytes/macrophages: CD11b⁺ F4/80⁺; monocytic or granulocytic MDSCs: CD11b⁺ Ly6C⁺ or CD11b⁺ Ly6G⁺, respectively; Treg: CD3⁺ CD4⁺ Foxp3⁺; dendritic cells maturation: fluorescence intensity of CD40 and CD86 labelling in live CD11c high MHC-II⁺ cells.

Autophagy Detection

Cyto-ID kit (Enzo) was used according to the manufacturer's protocol for flow cytometry and immunofluorescence. Anti-LC3 (clone 5F10, Nanotools) was used on cells fixed and permeabilized with BD fix/perm kit and revealed with alexa 488 anti-mouse (InvitroGen).

ROS Detection

Cells were incubated 2 hours with the indicated drugs, washed, then incubated with 100 μ M DCFH₂/DA (Sigma) for 50 min at 37°C, and analysed by flow cytometry.

Immunoblot Analysis

Samples were lysed in lysis buffer containing 1% SDS (Promega), 10 mM Tris HCl pH 7.5, 1 mM NaF (Sigma), 1 mM EDTA (Promega), 2 mM sodium orthovanadate (Sigma) and antiprotease cocktail (Roche, Mannheim, Germany). The protein concentration for each sample was determined by the DC protein assay according to the manufacturer's protocol (Bio-rad, Marnes-la-Coquette, France). After immunoblotting, immunochemical detection was revealed with an appropriate horseradish peroxidase-labeled secondary antibody (Jackson ImmunoResearch). Specific bands were visualized using a chemiluminescence detection system (Santa Cruz, Heidelberg, Germany), according to the manufacturer's protocol. Controls of equal loading of proteins were performed with anti-actin antibody. The following antibodies were used: monoclonal rabbit anti-p-eIF2 α (Cell Signaling), monoclonal rabbit anti-eIF2 α (Cell Signaling), monoclonal mouse anti- β -actin (Sigma).

Extracellular ATP Concentration

Supernatants were centrifugated twice (425 g, 3 min, room temperature) to eliminate cellular debris. Then ATP concentration was determined with the Cell Titer Glo Viability Assay Kit (Promega), according to the manufacturer's protocol.

ELISA

The following kits were used according to the manufacturer's protocol: HMGB1 detection kit (Chondrex), IFN γ ELISA kit (BD), TGF β 1 Ready-SET-Go ELISA kit (eBiosciences).

Primers and PCR

Nucleic acid purification was performed using Trizol (Invitrogen), according to the manufacturer's protocol. The M-MLV Reverse Transcriptase kit (Invitrogen) was used according to the manufacturer's protocol to perform reverse transcription. RT-qPCR was performed using the *Power SYBR*[®] Green PCR Master Mix kit (Applied Biosystems), according to the manufacturer's protocol with a 7500 Fast Real Time PCR system (Applied Biosystems). The following primers were used: *b-actin* (internal control): forward 5'-ATGGAGGGGAATACAGCCC-3', reverse 5'-TTCTTTGCAGCTCCTTCGTT-3'; *Tgfb1*: forward 5'-

CAACCCAGGTGCTTCCTAAA-3', reverse 5'-GGA-GAGCCCTGGATACCAAC-3'.

shRNA Transduction

CT26 cells were transduced with MISSION Lentiviral Transduction Particles (Sigma-Aldrich) to silence CRT or with negative control particles according to the manufacturer's instructions.

Statistical Analysis

Data were analysed with GraphPad Prism 5 Software. Comparisons between control and test conditions were performed with student t test or Mann&whitney test; group comparison were performed with Kruskal-Wallis test followed by Dunn's post-test. Tumor growth comparisons were performed by two-way ANOVA method.

Results

BLM Induces Stigmata of Immunogenic Cell Death

BLM antitumor effect is due to its capacity to induce radical oxygen species (ROS) production [19,20]. Because ROS generation could lead to reticulum stress [21], we hypothesized that BLM could be able to cause ROS-dependent ER stress and the induction of ICD. We found that BLM, like doxorubicin (DOX), an anthracycline and immunogenic drug [7], triggers eIF2 α phosphorylation, a marker of reticulum stress in CT26 cells (Figure 1a). Chemotherapy-induced reticulum stress could trigger calreticulin (CRT) and the endoplasmic reticulum chaperone ERp57 translocation at the cell surface [22,23]. *In vitro*, BLM induced translocation of CRT and ERp57 onto tumor cell plasma membrane in three mouse cancer models (Figure 1b). As expected, BLM effects are comparable to DOX effects, whereas mitomycin C (Mito C), a non-immunogenic drug [7], did not induce CRT or ERp57 exposure, (Figure 1b). Importantly, we showed that these translocations are aborted by using N-acetylcystein (NAC) (Figure 1c), a ROS inhibitor (Figure S1). The second determinant of ICD is the capacity of chemotherapy to induce the release of HMGB1 which triggers TLR4 on antigen presenting cells [8]. Using ELISA, we observed that BLM induces similar release of HMGB1 as doxorubicin (Figure 1d). The third element of ICD is the capacity of the chemotherapeutic agent to trigger autophagy and subsequent ATP secretion which activates dendritic cells via P2RX7 [9,24]. We observed that BLM induced, like DOX, LC3 puncta in immunofluorescence and Cyto-ID labeling in flow cytometry, two markers of autophagy vacuoles (Figure 1e, 1f and Figure S2). Of note, mitomycin C was unable to induce these signs of autophagy (Figure S2). Similarly to DOX, BLM also induced ATP release from CT26 cells (Figure 1g). The ROS inhibitor NAC also blunted both autophagy and ATP release (Figure 1f and 1g).

Together, these data demonstrate that BLM triggers all characteristic events of immunogenic cell death [7–9] in a ROS-dependent manner.

BLM-induced Immunogenic Cell Death Drives CD8⁺ Dependent Antitumor Immunity

To determine whether BLM-treated tumor cells induced antitumor immune response *in vivo*, we vaccinated naïve mice by injecting B16-OVA cells treated with PBS, BLM or DOX in the footpad. We observed that BLM- and DOX- treated cells are able to induce ovalbumin-dependent IFN γ production by T cells from the draining lymph node (Figure 2a). We also performed *in vivo* vaccination with control or CRT-knocked down BLM-treated dying tumor cells, followed by a challenge with live tumor cells.




Figure 1. BLM treatment triggers characteristic events of immunogenic cell death. A: Western blotting analysis of phosphorylated, total eIF2 α and β -actin in CT26 treated 24 h with the indicated drugs. B: indicated tumor cells were treated *in vitro* with different chemotherapeutic drugs for 24 h, then CRT (left panel) or Erp57 (right panel) exposure was detected by flow cytometry. C: tumor cells were treated with BLM, with or without N-acetylcystein (NAC) for 24 h, then CRT (left panel) or Erp57 (right panel) exposure was detected by flow cytometry. C: HMGB1 was titrated with ELISA method in supernatant of CT26 cells treated 24 h with indicated drugs. E: CT26 cells were cultured on glass slides and treated with indicated drugs for 24 h. The cells were then submitted to anti-LC3 (green) and DAPI (blue) labeling and observed under an epifluorescence microscope. F: CT26 cells were treated with indicated drugs for 24 h, then stained using Cyto-ID autophagy detection kit and analysed by flow cytometry. G: ATP

was titrated in supernatant of CT26 treated with indicated drugs for 24 h. All data presented are representative of one out of three experiments. B-D, E and F show mean \pm SEM.

doi:10.1371/journal.pone.0065181.g001

We could observe that the BLM-treated control CT26 cells were able to induce protective immunity (100% protection, $n = 10$), in contrast to mitomycin-treated CT26 (20% protection, $n = 10$) and CRT-invalidated BLM-treated cells (30% protection, $n = 10$). Furthermore, in the setting of growing s/c CT26 tumors, we observed that 5 injections of BLM delayed the growth of control shRNA transfected CT26 cells but had no effect on CRT shRNA transfected CT26 cells (Figure 2b). Finally, IFN γ or CD8 depletion completely prevents BLM antitumor effect *in vivo* (Figure 2c and 2d). Together, these data strongly suggest that BLM exerts a calreticulin-, IFN γ - and CD8-dependent antitumor effect.

BLM Shows No Effect on Innate Immune Cells

In addition to its capacity to elicit ICD of tumor cells, chemotherapy could harness the host immune system by inducing dendritic cell maturation [25], NK cell activation [1] or by eliminating immunosuppressive cells such as Treg and MDSC [3,4,26].

To address these possibilities in the BLM treatment, we have first assessed if BLM injection did significantly modify the number of B cells, CD4 $^+$ T cells, CD8 $^+$ T cells, NK cells, dendritic cells, monocytes/macrophages or MDSC in spleen of tumor-bearing mice (Figure 3a). We observed that BLM treatment did not change the proportion of all the cell types analyzed. We further tested whether BLM promoted the maturation of antigen presenting cells by testing the effect of a single systemic injection of BLM or lipopolysaccharide (LPS) on the maturation pattern of splenic dendritic cells (DC). While LPS induced as expected a massive up-regulation of CD86 and CD40, BLM administration did not affect the expression of these markers, ruling out any direct DC-activating properties of BLM (Figure 3b).

We also checked NK cell activation in tumor draining lymph nodes. We did not observe any NK activation, assessed by CD69 expression, after either PBS, one or five BLM injections (Figure 3c).

Finally, we evaluated the frequency of MDSC in the spleen of BLM-treated tumor-bearing mice. We found that BLM did not affect the number of myeloid or granulocytic MDSC subsets (Figure 3d) and did not affect MDSC immunosuppressive function, as assessed by MDSC capacity to blunt OT-I capacity to produce IFN γ (Figure 3e).

Altogether, these data suggest that BLM does not affect innate immune cells such as dendritic cells, NK cells or MDSC.

BLM Mediates Regulatory T Cell Proliferation via Tumor Cell-derived TGF β

While we detected no change in most immune cell populations (Figure 3a), we observed an accumulation of Foxp3 $^+$ Treg in lymphoid organs of mice treated with BLM and bearing CT26 (Figure 4a) or EL4 (not shown) tumors. We then determined the biological function of these Treg. We sorted CD4 $^+$ CD25 $^+$ (Treg) and CD4 $^+$ CD25 $^-$ (conventional T) T cell subsets from the spleen from PBS- or BLM-treated tumor-bearing mice, and tested their capacity to blunt IFN γ production by OT-I cells in an *in vitro* assay. We observed as expected that conventional T cells were unable to blunt OT-I capacity to produce IFN γ . In contrast, Treg from untreated or BLM-treated mice had strong and similar capacities to blunt OT-I IFN γ production (Figure 4b).

Higher Treg presence could be due to differentiation of naive T cells into Treg or expansion of pre-existing Treg. To test these

hypotheses, we transferred CD45.2 CD4 $^+$ GFP $^+$ (Treg) or CD45.2 CD4 $^+$ GFP $^-$ CD62L $^+$ (naive T cells) from Foxp3-EGFP mice into CD45.1 tumor bearing mice (Figure 4c). We observed that BLM induces proliferation of Treg at similar level to IL-2 injection, a known Treg inducer (Figure 4d) [27,28]. In contrast, BLM was unable to induce differentiation of naive T cells into Treg (Figure 4e).

Treg cell proliferation is known to be driven by IL-2 production [28,29] or TGF β [30,31]. We tested the presence of IL-2 and TGF β in ascites of a CT26 peritoneal carcinosis model. While we could not detect any IL-2 in ascites from PBS or BLM-treated mice (not shown), we observed an induction of TGF β production in the ascites of BLM-treated animals (Figure 4f). *In vitro*, we also observed that BLM treatment induced TGF β production by CT26 both at the mRNA (Figure S3) and protein level (Figure 4g). Similarly, we observed that BLM could induce *tgfb* mRNA expression in B16F10, EL4 and MCA205 cells (not shown).

To validate that TGF β was also produced *in vivo* by tumor cells, we determined the expression of latency-associated protein (LAP), a protein associated with TGF β , in cancer and immune cells. Only tumor cells isolated from subcutaneous tumor showed increased LAP expression after BLM treatment (Figure 4h and Figure S4).

To test if this TGF β production by tumor cells may be responsible for Treg accumulation in BLM-treated mice, we sorted CD4 $^+$ GFP $^+$ from Foxp3-EGFP mice and stimulated them with T cell receptor (TCR) triggering. In some conditions, supernatant from PBS- or BLM-treated CT26 cells was added, with or without anti-TGF β blocking antibody. After 72 h of stimulation, the proliferation of Treg was assessed by a BrdU incorporation assay. We could observe that supernatant from BLM-treated CT26 induces proliferation of Treg, and that this effect is dependent on TGF β (Figure 4i). *In vivo*, we also observed that inhibition of TGF β with a blocking anti-TGF β antibody prevents the BLM-induced accumulation of Treg in tumor-bearing mice (Figure 4j).

Taken together, these data strongly suggest that BLM induces Treg accumulation through enhancement of tumor-produced TGF β .

TGF β or Regulatory T cell Depletion Potentiate the Antitumor Effects of BLM

To assess the therapeutic relevance of these observations, we tested the effect of Treg depletion on the antitumor effect of BLM. In the colon carcinoma model CT26. We first observed that the antitumor effect of BLM was drastically improved by CD4 $^+$ T cell depletion (Figure 5a). To validate that the antitumor effect of CD4 depletion is mediated essentially by Treg, we tested the effect of an anti-CD25, which is a membranous marker for mouse Treg, depleting antibody. We observed that CD25 depletion also enhanced the antitumor efficacy of BLM (Figure 5b).

We demonstrated above (Figure 4) that BLM induces TGF β production by tumor cells and Treg expansion in tumor bearing mice. To determine if this phenomenon impacts on BLM antitumor effect, we treated tumor-bearing mice with either BLM, anti-TGF β or both. While the anti-TGF β antibody did not significantly impact on tumor growth, TGF β depletion enhanced the antitumor efficacy of BLM (Figure 5c).

Taken together, these data highlight that BLM treatment causes TGF β -dependent proliferation of Treg, which decreases its antitumor efficacy.




Figure 2. BLM in vivo antitumor effect through immune-based mechanisms. A: Footpad injection of different vaccines was performed: PBS-, DOX- or BLM-treated B16-OVA, or ovalbumin protein. 5 days later, the popliteal lymph nodes were harvested and cells were rechallenged with OVA peptide SIINFEKL. After 3 days of culture, IFN γ was titrated in the supernatant using ELISA method. B: shControl CT26 (left panel) or shCRT CT26 (right panel) were injected in the flank of mice. When the tumor reached about 25 mm², mice were treated with PBS or BLM and tumor growth was monitored with a caliper over time. C and D: CT26 cells were injected in the flank of mice. When the tumor reached about 25 mm², mice were treated with PBS or BLM. Some of the mice received injection of isotype control or depleting anti-CD8 (C) or anti-IFN γ (D). We monitored tumor growth with a caliper over time. All data presented are representative of one out of two experiments. Graphs show mean \pm SEM. doi:10.1371/journal.pone.0065181.g002

Discussion

Our study reveals that BLM can be counted among the immunogenic anticancer drugs which trigger ICD along with oxaliplatin and anthracyclines [7,8,32]. Here, by studying the

immune-based effects of BLM anti-cancer treatment, we show that BLM is capable to cause ER and oxidative stress, CRT exposure, HMGB1 release, autophagy and ATP release, providing all the elements required to induce an ICD, leading to IFN γ and CD8⁺ T




Figure 3. BLM has no obvious effect on innate immune cells. A: Spleens from CT26 tumor-bearing mice, treated with PBS or BLM, were harvested and the cells were analyzed through flow cytometry. The graph shows proportion of different splenocyte populations identified by immunostaining. B: Mice received a single injection of PBS, LPS or BLM. One day later, spleens were collected and the maturation status of dendritic cells defined as CD11c^{hi} and MHC-II⁺ was assessed by flow cytometry, with CD40 (left panel) and CD86 (right panel) staining. C: We injected CT26 cells in the thigh of mice. When the tumor reached about 25 mm², we treated them with PBS, single or five BLM injections. One day after last injection, inguinal lymph nodes were collected and NK cell activation was analysed through CD69 immunostaining. D: Same as A with further identification of MDSC sub-populations with Ly6G and Ly6C immunostaining. The cells are gated as CD11b^{hi} among the live cells. E: MDSC from PBS- or BLM- treated CT26-bearing mice were isolated from spleen and co-cultured with OT-I in presence of SIINFEKL. After 3 days of culture, IFN γ was titrated in the supernatant using ELISA method. All data presented are representative of one out of three experiments. Graphs show mean \pm SEM. doi:10.1371/journal.pone.0065181.g003

cell response against tumor cells. Importantly, BLM has a two-edged effect, because BLM also induces the production of TGF β by tumor cells and consequent Treg accumulation which limits its anti-tumor efficacy. These data demonstrate that BLM may have ambivalent immune effects which limit the antitumor efficacy of this anticancer agent. This is similar to our recent conclusions on the immune effects of 5-fluorouracil (5 FU) on tumor immunity. We indeed described that 5 FU induces MDSC death, which restores CD8⁺ T cell antitumor immunity and causes initial

efficacy of chemotherapy [4]. But while MDSC die, they release IL-1 β , which induces accumulation of deleterious Th17 cells, promoting later tumor relapse [33]. Together, these observations about 5 FU and BLM underscore the importance to perform a systematic analysis of the immune effect of anticancer agents on the different aspects of the immune response.

Besides, cardiac glycosides, such as digoxin, were recently identified as strong immunogenic cell death inducers, using a high throughput strategy [34]. In this study, it was shown in rodent




Figure 4. BLM induces in vivo Treg accumulation through TGF β production. A: Splens from CT26 tumor-bearing mice, treated with PBS or BLM, were harvested and the cells were analyzed through flow cytometry for Foxp3⁺CD4⁺ Treg detection. B: Treg from PBS- or BLM- treated CT26-bearing mice were isolated from spleen and co-cultured with OT-I in presence of SIINFEKL (SII). After 3 days of culture, IFN γ was titrated in the supernatant using ELISA method. C: Schematic representation for D and E panel experiments. D: GFP⁺ CD4⁺ cells were sorted from CD45.2 FOXP3-EGFP mice, then injected i.v. in CD45.1 mice bearing CT26 tumor. The mice then received PBS, IL-2 or BLM treatment. All mice received EdU injection.

One day after treatment, spleens and tumors were collected and the proliferation status of transferred cells was assessed by revealing EdU by flow cytometry. E: CD4⁺ CD62L⁺ GFP⁻ naive T cells were sorted from CD45.2 FOXP3-EGFP mice, and injected i.v. in CD45.1 mice bearing CT26 tumor. The mice then received PBS or BLM treatment. Spleens were harvested and the cells were analyzed through flow cytometry for Treg detection. F: Mice were injected with 1.10⁵ CT26 cells i.p. Ten days later mice received PBS or BLM injection. The following day ascites were collected, and TGFβ was assessed using ELISA method. G: CT26 cells were treated with PBS or BLM *in vitro* for 24 h, then the supernatant was collected and assessed for TGFβ using ELISA method. H: Mice bearing CT26 tumor cells were treated with PBS or BLM. The day after, CT26 tumors were collected and tumor cells were separated from the tumor infiltrating lymphocytes. Tumor cells were stained for LAP and analyzed by flow cytometry. I: GFP⁺ CD4⁺ cells were sorted from CD45.2 FOXP3-EGFP mice, then cultured *in vitro* under TCR-stimulating conditions. We added culture supernatant of CT26 treated for 24 h with PBS or BLM. In some wells, blocking anti-TGFβ antibody was added. After three days of culture, the cells were incubated with BrdU for 3 h, and then proceeded to BrdU detection by flow cytometry. J: Same as A, and mice received injection of blocking anti-TGFβ antibody. All data presented are representative of one out of two (panels B, D, E, G and I) or three experiments (panels A, F, H and J). Graphs show mean ± SEM.

models that treatment of tumor cells with digoxin can enhance the effects of non-immunogenic cytotoxic drugs, such as cisplatin [32], due to their capacity to convert non immunogenic to immunogenic cell death. However, the effect of BLM on the immune response was not detected using this strategy. So we believe that systematic analysis of immune properties of chemotherapeutic




Figure 5. BLM anti-tumor effect is enhanced by Treg or TGFβ depletion. Five hundred thousand CT26 cells were injected in the flank of mice. When the tumor reached about 25 mm², mice were treated with PBS or BLM. Some of the mice received injections of isotype control antibody or depleting anti-CD4 (A), anti-CD25 (B) or blocking anti-TGFβ (C) antibodies. Tumor growth was monitored with a caliper over time. All data presented are representative of one out of two experiments. Graphs show mean ± SEM.

agents on each immune cell subset is complementary to high throughput strategy.

Foxp3⁺CD4⁺CD25⁺ Treg were initially defined as a subpopulation of suppressor T cells that mediate immune tolerance by suppressing autoreactive T cells [35,36]. In human neoplasia, Treg cells accumulate in tumors, draining lymph nodes and blood, and their induction is often correlated with poor outcome both in human and rodent cancers [37]. Activation and expansion of Treg subset was determined to be dependent on IL-2 [28,29] and TGFβ [30,31]. BLM treatment of lung fibroblasts results in an increased transcription of *tgfβ1* and increased TGFβ protein [38,39]. This production of TGFβ by fibroblasts in the lung after BLM injection is responsible for the induction of lung fibrosis, one of the major side-effects of BLM [40,41]. Here we made the original observation that BLM could also trigger TGFβ production by tumor cells. We may suspect that this increase in TGFβ expression could result in immunosubversion of anticancer immune response. In particular, we showed that BLM-induced TGFβ production by tumor cells drives Treg expansion and may thus inhibit the CD8⁺ antitumor immune response induced by BLM-driven immunogenic cell death. In the B16F10 model, while either high or low BLM dosage could induce the production of TGFβ by B16F10, only high concentration of BLM could induce ICD (not shown). We may therefore raise the hypothesis that in some cases the ICD effect which only occurs with high dosage of BLM is less likely to happen *in vivo* and that the TGFβ-Treg driven induction would be predominant. We also show that Treg depletion or TGFβ inhibition significantly enhance BLM anti-tumor efficacy in our *in vivo* rodent model. Taken together with the current literature, our data suggest that TGFβ and Treg are responsible for both limited antitumor effect and major side effect of BLM [40,41]. This may give the rationale to disrupt this phenomenon by combining BLM and Treg inhibitors, such as cyclophosphamide [3]. In the future, the association between BLM and novel drugs like anti-human CD25 (such as daclizumab) [42] or inhibitors of TGFβ signaling (such as Smad inhibitors) [43] may be beneficial for both anti-cancer efficacy and side-effect management.

In conclusion, this study supports the ambivalent immunological property of BLM by its capacity to trigger immunogenic cell death and Treg accumulation.

Supporting Information

Figure S1 BLM treatment induces ROS production. CT26 cells were treated 24 h *in vitro* with PBS, Mitomycin C (Mito C), Doxorubicine (DOX) or bleomycin (BLM), with addition of PBS or N-acetylcystein (NAC). After 24 h of culture, production of radical species was assessed by DCFH₂/DA detection by flow cytometry. Data presented are representative of one out of three experiments. Graphs show mean ± SEM.

Figure S2 BLM treatment induces autophagy. CT26 cells were cultured on glass slide, and treated with PBS, Mito C, DOX or BLM for 24 h (scale bar: 10 μ m). Upper panel: cells were fixed and permeabilized, then labeled with anti-LC3 antibody (green) and DAPI (blue). Lower panel: cells were stained using the Cyto-ID autophagy detection kit, showing autophagosomes (green) and Hoescht 33342 (blue). Data presented are representative of one out of two experiments. (TIF)

Figure S3 BLM treatment induces *tgfb* transcription in tumor cells. CT26 cells were treated *in vitro* with PBS, BLM or DOX. After 24 h, cells were collected and their mRNA isolated, then submitted to RT-qPCR for *tgfb* expression. Results are shown standardized to *actin* expression. Data presented are representative of one out of three experiments. (TIF)

Figure S4 BLM *in vivo* treatment induces no noticeable modification of Lap in host immune cells. Mice bearing CT26 tumor cells were treated with PBS or BLM. The day after, CT26 tumors and spleen cells were collected. The tumor cells were separated from the tumor infiltrating lymphocytes. Spleen cells and tumor-infiltrating lymphocytes were stained for LAP and analysed by flow cytometry. Data presented are representative of one out of three experiments. (TIF)

Author Contributions

Conceived and designed the experiments: FG GM. Performed the experiments: H. Bugaut MB H. Berger VD LO RE SL FC FV CR. Analyzed the data: CR LA FG GM. Contributed reagents/materials/analysis tools: CR. Wrote the paper: FG GM.

References

- Hervieu A, Rebe C, Vegran F, Chalmin F, Bruchard M, et al. (2012) Dacarbazine-Mediated Upregulation of NKG2D Ligands on Tumor Cells Activates NK and CD8 T Cells and Restrains Melanoma Growth. *J Invest Dermatol*.
- Mignot G, Ullrich E, Bonmort M, Menard C, Apetoh L, et al. (2008) The critical role of IL-15 in the antitumor effects mediated by the combination therapy imatinib and IL-2. *J Immunol* 180: 6477–6483.
- Ghiringhelli F, Menard C, Puig PE, Ladoire S, Roux S, et al. (2007) Metronomic cyclophosphamide regimen selectively depletes CD4+CD25+ regulatory T cells and restores T and NK effector functions in end stage cancer patients. *Cancer Immunol Immunother* 56: 641–648.
- Vincent J, Mignot G, Chalmin F, Ladoire S, Bruchard M, et al. (2010) 5-Fluorouracil selectively kills tumor-associated myeloid-derived suppressor cells resulting in enhanced T cell-dependent antitumor immunity. *Cancer Res* 70: 3052–3061.
- Apetoh L, Locher C, Ghiringhelli F, Kroemer G, Zitvogel L (2011) Harnessing dendritic cells in cancer. *Semin Immunol* 23: 42–49.
- Tesniere A, Panaretakis T, Kepp O, Apetoh L, Ghiringhelli F, et al. (2008) Molecular characteristics of immunogenic cancer cell death. *Cell Death Differ* 15: 3–12.
- Obeid M, Tesniere A, Ghiringhelli F, Fimia GM, Apetoh L, et al. (2007) Calreticulin exposure dictates the immunogenicity of cancer cell death. *Nat Med* 13: 54–61.
- Apetoh L, Ghiringhelli F, Tesniere A, Obeid M, Ortiz C, et al. (2007) Toll-like receptor 4-dependent contribution of the immune system to anticancer chemotherapy and radiotherapy. *Nat Med* 13: 1050–1059.
- Ghiringhelli F, Apetoh L, Tesniere A, Aymeric L, Ma Y, et al. (2009) Activation of the NLRP3 inflammasome in dendritic cells induces IL-1 β -dependent adaptive immunity against tumors. *Nat Med* 15: 1170–1178.
- Ryzhkov VK, Zeidlits VN, Tikhonov KB (1977) [Use of protamine sulfate as a vasodilator in liver and spleen angiography]. *Vestn Rentgenol Radiol*: 41–45.
- Dresp J, Schmid E, Bauchinger M (1978) The cytogenetic effect of bleomycin on human peripheral lymphocytes *in vitro* and *in vivo*. *Mutat Res* 56: 341–353.
- Caspary WJ, Lanzo DA, Niziak C (1982) Effect of deoxyribonucleic acid on the production of reduced oxygen by bleomycin and iron. *Biochemistry* 21: 334–338.
- Hay J, Shahzeidi S, Laurent G (1991) Mechanisms of bleomycin-induced lung damage. *Arch Toxicol* 65: 81–94.
- Meyer RM, Gospodarowicz MK, Connors JM, Pearcey RG, Wells WA, et al. (2012) ABVD alone versus radiation-based therapy in limited-stage Hodgkin's lymphoma. *N Engl J Med* 366: 399–408.
- Williams SD, Birch R, Einhorn LH, Irwin L, Greco FA, et al. (1987) Treatment of disseminated germ-cell tumors with cisplatin, bleomycin, and either vinblastine or etoposide. *N Engl J Med* 316: 1435–1440.
- Bettelli E, Carrier Y, Gao W, Korn T, Strom TB, et al. (2006) Reciprocal developmental pathways for the generation of pathogenic effector TH17 and regulatory T cells. *Nature* 441: 235–238.
- Chalmin F, Ladoire S, Mignot G, Vincent J, Bruchard M, et al. (2010) Membrane-associated Hsp72 from tumor-derived exosomes mediates STAT3-dependent immunosuppressive function of mouse and human myeloid-derived suppressor cells. *J Clin Invest* 120: 457–471.
- Chalmin F, Mignot G, Bruchard M, Chevriaux A, Vegran F, et al. (2012) Stat3 and Gfi-1 transcription factors control Th17 cell immunosuppressive activity via the regulation of ectonucleotidase expression. *Immunity* 36: 362–373.
- Burger RM, Peisach J, Horwitz SB (1981) Activated bleomycin. A transient complex of drug, iron, and oxygen that degrades DNA. *J Biol Chem* 256: 11636–11644.
- Wallach-Dayana SB, Izbicki G, Cohen PY, Gerstl-Golan R, Fine A, et al. (2006) Bleomycin initiates apoptosis of lung epithelial cells by ROS but not by Fas/FasL pathway. *Am J Physiol Lung Cell Mol Physiol* 290: L790–L796.
- Garg AD, Krysko DV, Verfaillie T, Kaczmarek A, Ferreira GB, et al. (2012) A novel pathway combining calreticulin exposure and ATP secretion in immunogenic cancer cell death. *EMBO J* 31: 1062–1079.
- Panaretakis T, Kepp O, Brockmeier U, Tesniere A, Bjorklund AC, et al. (2009) Mechanisms of pre-apoptotic calreticulin exposure in immunogenic cell death. *EMBO J* 28: 578–590.
- Panaretakis T, Joza N, Modjtahedi N, Tesniere A, Vitale I, et al. (2008) The co-translocation of ERp57 and calreticulin determines the immunogenicity of cell death. *Cell Death Differ* 15: 1499–1509.
- Michaud M, Martins I, Sukkurwala AQ, Adjemian S, Ma Y, et al. (2011) Autophagy-dependent anticancer immune responses induced by chemotherapeutic agents in mice. *Science* 334: 1573–1577.
- Tanaka H, Matsushima H, Nishibu A, Clausen BE, Takashima A (2009) Dual therapeutic efficacy of vinblastine as a unique chemotherapeutic agent capable of inducing dendritic cell maturation. *Cancer Res* 69: 6987–6994.
- Suzuki E, Kapoor V, Jassar AS, Kaiser LR, Albelda SM (2005) Gemcitabine selectively eliminates splenic Gr-1+CD11b+ myeloid suppressor cells in tumor-bearing animals and enhances antitumor immune activity. *Clin Cancer Res* 11: 6713–6721.
- Thornton AM, Donovan EE, Piccirillo CA, Shevach EM (2004) Cutting edge: IL-2 is critically required for the *in vitro* activation of CD4+CD25+ T cell suppressor function. *J Immunol* 172: 6519–6523.
- Melencio L, McKallip RJ, Guan H, Ramakrishnan R, Jain R, et al. (2006) Role of CD4(+)CD25(+) T regulatory cells in IL-2-induced vascular leak. *Int Immunol* 18: 1461–1471.
- Thornton CA, Upham JW, Wikstrom ME, Holt BJ, White GP, et al. (2004) Functional maturation of CD4+CD25+CTLA4+CD45RA+ T regulatory cells in human neonatal T cell responses to environmental antigens/allergens. *J Immunol* 173: 3084–3092.
- Huber S, Schramm C, Lehr HA, Mann A, Schmitt S, et al. (2004) Cutting edge: TGF- β signaling is required for the *in vivo* expansion and immunosuppressive capacity of regulatory CD4+CD25+ T cells. *J Immunol* 173: 6526–6531.
- Ghiringhelli F, Puig PE, Roux S, Parcellier A, Schmitt E, et al. (2005) Tumor cells convert immature myeloid dendritic cells into TGF- β -secreting cells inducing CD4+CD25+ regulatory T cell proliferation. *J Exp Med* 202: 919–929.
- Tesniere A, Schlemmer F, Boige V, Kepp O, Martins I, et al. (2010) Immunogenic death of colon cancer cells treated with oxaliplatin. *Oncogene* 29: 482–491.
- Bruchard M, Mignot G, Derangere V, Chalmin F, Chevriaux A, et al. (2012) Chemotherapy-triggered cathepsin B release in myeloid-derived suppressor cells activates the Nlrp3 inflammasome and promotes tumor growth. *Nat Med*.
- Menger L, Vacchelli E, Adjemian S, Martins I, Ma Y, et al. (2012) Cardiac glycosides exert anticancer effects by inducing immunogenic cell death. *Sci Transl Med* 4: 143ra199.
- Sakaguchi S (2000) Regulatory T cells: key controllers of immunologic self-tolerance. *Cell* 101: 455–458.
- Shevach EM (2002) CD4+ CD25+ suppressor T cells: more questions than answers. *Nat Rev Immunol* 2: 389–400.
- Nishikawa H, Sakaguchi S (2010) Regulatory T cells in tumor immunity. *Int J Cancer* 127: 759–767.
- Breen E, Shull S, Burne S, Absher M, Kelley J, et al. (1992) Bleomycin regulation of transforming growth factor- β mRNA in rat lung fibroblasts. *Am J Respir Cell Mol Biol* 6: 146–152.
- Li H, He B, Que C, Weng B (1996) Expression of TGF- β 1, PDGF and IGF-1 mRNA in lung of bleomycin-A5-induced pulmonary fibrosis in rats. *Chin Med J (Engl)* 109: 533–536.

40. Gaudie J, Kolb M, Ask K, Martin G, Bonniaud P, et al. (2006) Smad3 signaling involved in pulmonary fibrosis and emphysema. *Proc Am Thorac Soc* 3: 696–702.
41. Khalil N, Greenberg AH (1991) The role of TGF-beta in pulmonary fibrosis. *Ciba Found Symp* 157: 194–207; discussion 207–111.
42. Rech AJ, Vonderheide RH (2009) Clinical use of anti-CD25 antibody daclizumab to enhance immune responses to tumor antigen vaccination by targeting regulatory T cells. *Ann N Y Acad Sci* 1174: 99–106.
43. Calone I, Souchelnytskyi S (2012) Inhibition of TGFbeta signaling and its implications in anticancer treatments. *Exp Oncol* 34: 9–16.