

Fruit and vegetable intakes and asthma in the E3N study.

Isabelle Romieu, Raphaëlle Varraso, Valérie Avenel, Bénédicte Leynaert, Francine Kauffmann, Françoise Clavel-Chapelon

► **To cite this version:**

Isabelle Romieu, Raphaëlle Varraso, Valérie Avenel, Bénédicte Leynaert, Francine Kauffmann, et al.. Fruit and vegetable intakes and asthma in the E3N study.. Thorax, BMJ Publishing Group, 2006, 61 (3), pp.209-15. <10.1136/thx.2004.039123>. <inserm-00143617>

HAL Id: inserm-00143617

<http://www.hal.inserm.fr/inserm-00143617>

Submitted on 27 Apr 2007

HAL is a multi-disciplinary open access archive for the deposit and dissemination of scientific research documents, whether they are published or not. The documents may come from teaching and research institutions in France or abroad, or from public or private research centers.

L'archive ouverte pluridisciplinaire **HAL**, est destinée au dépôt et à la diffusion de documents scientifiques de niveau recherche, publiés ou non, émanant des établissements d'enseignement et de recherche français ou étrangers, des laboratoires publics ou privés.

Fruit and vegetable intakes and asthma in the E3N study

I Romieu^{1,*}, R Varraso^{2,3}, V Avenel⁴, B Leynaert⁵, F Kauffmann², F Clavel-Chapelon⁴

¹ Instituto Nacional de Salud Pública, México, ² INSERM, U780, Villejuif, France, ³ Université Paris 11, Kremlin-Bicêtre, France, ⁴ INSERM, Equipe Nutrition, Hormones et Cancer, Institut Gustave Roussy, Villejuif, France, ⁵ INSERM, U700, Paris, and Université Paris Diderot, Faculté de Médecine Xavier Bichat, Paris, France.

Background: A study was undertaken to investigate whether dietary intake predicted the prevalence of adult asthma among French women participating in the E3N study.

Methods: Of 68 535 women who completed a food frequency questionnaire in 1993 which included 238 food items, 2145 (3.1%) reported having asthma. The distribution of food intake was divided into quartiles (Q₁–Q₄) and the prevalence of asthma was compared between the different quartiles (lowest as reference) using logistic regression models on cross sectional data.

Results: After adjusting for age, body mass index, menopausal status, smoking status, total caloric intake, physical activity, and use of dietary supplements, women who had a greater intake of tomatoes (OR_{Q₁–Q₄} 0.85 95% CI 0.75 to 0.96, test for trend p=0.02), carrots (OR_{Q₁–Q₄} 0.81 95% CI 0.72 to 0.92, test for trend p=0.0003), and leafy vegetables (OR_{Q₁–Q₄} 0.82 95% CI 0.73 to 0.93, test for trend p=0.0009) had a lower prevalence of asthma. Apples were marginally related to the prevalence of asthma. No other fruits or vegetables were significantly associated with asthma prevalence.

Conclusions: These results suggest that the intake of some vegetables may decrease the prevalence of adult asthma.

Environmental factors appear to be critical for the increase in the prevalence of asthma recently observed in most developed countries. Changes in diet such as decreased consumption of fresh fruit and vegetables might have a major role.^{1–3} However, the epidemiological evidence for the role of food intake in the prevalence of asthma among adults is still scarce. Previous studies have reported that a greater intake of fresh fruit is related to a lower prevalence of respiratory symptoms⁴ and to a lower incidence of nonspecific lung diseases,⁵ but no specific foods were identified. Although vitamin C is believed to be the principal candidate for the apparent benefit of plant foods,⁶ a prospective study⁷ found no beneficial effect of dietary vitamin C intake and intervention trials provided conflicting results.⁸ Antioxidants other than the vitamin C contained in fruit and vegetables might play an important role in asthma. In a recent casecontrol study, Shaheen *et al*⁹ reported that the consumption of apples and red wine were negatively related to asthma, suggesting the beneficial effect of flavonoids.

There are a number of reasons for exploring the effects of fruit and vegetables rather than individual nutrients. Food contains several substances and an analysis focusing on nutrients might miss important components and interactions between nutrients. Also, people eat food and not nutrients, so foods are easier to convert into dietary recommendation. For these reasons, public health programmes have focused on the promotion of consumption of fruit and vegetables.¹⁰ Although this recommendation was mostly based on results from cancer research, it appears that the consumption of fruit and vegetables might also be beneficial for lung health. It is therefore important to determine which fruit and vegetables are most beneficial for asthma prevention. We used data from the E3N (Etude Epidémiologique auprès des femmes de la MGEN) study, the French branch of the European Prospective Investigation into Cancer and Nutrition (EPIC), to evaluate the association of fruit and vegetable intake with asthma prevalence in a large population of adult women.

* Correspondence to: Dr I Romieu, Instituto Nacional de Salud Publica, Av. Universidad # 655, Col Santa Maria Ahuacatitlan, 62508 Cuernavaca, Morelos, Mexico; iromieu@correo.insp.mx

METHODS

Study population

The E3N study is a prospective investigation of major chronic diseases among a cohort of members from the Mutuelle Générale de l'Éducation Nationale (MGEN), a national health insurance plan covering mostly teachers. The general characteristics of this population have been reported elsewhere.¹¹ This study began in 1990 when a baseline questionnaire (Q₁) was sent to 103 089 women aged 40–65 years. A follow up questionnaire was sent in January 1992 (Q₂) and approximately every 2 years thereafter. In June 1993, in addition to the follow up questionnaire, a dietary assessment was conducted using a dietary history questionnaire (Q₃) to which 75 518 women responded.

Our study population was composed of 68 535 women who answered Q₁, Q₂ and Q₃ and met the inclusion criteria for the analysis (fig 1). Reasons for exclusion were: no answer to the previous questionnaire (Q₂) (n = 4424); unrealistic figures reported to the food questionnaire (Q₃) based on the ratio of energy intake/energy requirement (based on the EPIC methodology¹¹) (n = 1294); asthma diagnosis during childhood or no specified date for asthma diagnosis (n = 1265). In order to study a homogeneous group of asthmatics, analyses were restricted to those who reported adult onset asthma. One important reason for this choice is that longitudinal studies performed in cohorts from childhood to adulthood and during adulthood have shown a low tracking of dietary habits from childhood to adulthood,¹² while tracking of diet in adulthood is acceptable.¹³

The final participation rate was 68 535/98 997 (69.2%). Data were analysed cross sectionally to evaluate the association between fruit and vegetable intake and asthma prevalence.




Figure 1 Design of the study.

Definition of a prevalent case of asthma

Our definition of asthma was based on the presence of an affirmative answer to the questions “Did you have an asthma attack?” and “Age at time of the first attack” on the second follow up questionnaire (Q₃). Among the 68 535 volunteers we identified 2145 women with asthma who reported having the first attack in adulthood and provided the age when they had been diagnosed. These were included in the analysis.

Our classification of asthma was further validated using information gathered in the follow up questionnaire sent in 2003 (Q₇). Asthma was defined by questions based on the ATS questionnaire:¹⁴ “Have you ever had asthma attacks” and, if yes, “Was this diagnosis confirmed by a doctor ?” Of the 68 535 women included in the present analysis, 62 849 provided information on asthma in 2003 (Q₇); 725 women who reported

incident asthma between Q₃ and Q₇ and 374 who reported asthma in childhood in Q₇ were excluded. Based on the 61 750 women with both information at Q₃ and Q₇, the agreement between the asthma definition used in the present analysis (Q₃) and asthma assessed in Q₇ was very strong. Overall, 1661 women (2.7%) were defined as asthmatics in Q₃ and 1909 (3.1%) in Q₇. Most of the women who reported asthma in Q₃ were classified as asthmatics in Q₇. The agreement between the asthma definition used in the present analysis (Q₃) and asthma assessed in Q₇ (ATS definition) was strong (kappa statistic 0.77), and the definition of asthma at Q₃ had a high sensitivity (72.3%) and specificity (99.4%) using asthma assessed in Q₇ as the gold standard.

Food frequency questionnaire

The dietary questionnaire sent to participants in 1993 was composed of two parts with a booklet of photographs to facilitate the estimation of portion sizes. The first part included questions on the consumption (frequency and quantity) of 66 food groups. The second part included qualitative questions. Overall, the questionnaire provided an estimate for daily consumption of 208 food items. Both the questionnaire and the illustrated booklet had been validated previously.^{15, 16} We considered the food classification proposed by EPIC.¹⁷ Total calorie intake was estimated using the food frequency questionnaire and expressed in kcal/day.

Body mass index

Body mass index (BMI) was calculated based on the height and weight reported in the questionnaire administered at baseline. BMI was used as a continuous and categorical variable.

Other information

We used information provided on the 1993 questionnaire (Q₃). Tobacco consumption was categorised as never, past, and current smoker. Current exercise and energy expenditure were calculated by multiplying the duration of daily activities and leisure physical activities. Estimated metabolic energy spent was based on the values proposed by James and Schofield¹⁸ and categorised into quartiles of physical activity. Women were classified as premenopausal, postmenopausal or perimenopausal. Because our population was composed mostly of teachers from the MGEN, years of education were used as a proxy of socioeconomic status. The presence of allergy was classified as either present or absent.

Statistical analysis

The general characteristics of the asthmatic and control women were compared using the χ^2 test statistics for categorical data.¹⁹ Fruit and vegetable intake of the two groups of women were compared after log transformation of the data to normalise the distribution using *t* test statistics.¹⁹ The association between asthma in adulthood and fruit and vegetable intake was studied using logistic regression models,²⁰ adjusting for potential confounding factors including age, total calorie intake, BMI, physical activity, smoking status, menopausal status,²¹⁻²³ and the use of dietary supplements. Because our population came from a fairly homogenous socioeconomic status, adjusting for this variable (years of education) did not modify our estimates. Fruit and vegetable intake was categorised into quartiles based on the distribution among control women. The odds ratios (OR) were determined by comparison with the lowest quartiles. Correction for multiple comparisons was performed by the Bonferroni method.²⁴ The significance of the interaction between smoking status and fruit and vegetable intake was tested. Although the interaction terms were not significant, we further stratified our analysis by smoking status (never smoker, ex-smoker or current smoker) to determine if there was some indication of a modifying effect by this variable. We tested for linear trends in ORs with increasing exposure using the likelihood ratio test.²⁵ All analyses were conducted using SAS Version 8.

RESULTS

Of the 68 535 women included in the analysis, 2145 (3.1%) reported having had an asthma attack in adulthood before application of the food frequency questionnaire in 1993. The characteristics of the women included and excluded from the analysis were similar with regard to age, smoking status, menopausal status, physical activity, socioeconomic status, and use of dietary supplements, but women excluded were more likely to have a lower mean (SD) BMI (22.9 (3.3) kg/m² v 23.3 (3.8) kg/m², p<0.0001), to have a higher mean (SD) total calorie intake (2345.1 (1537.3) kcal/day v 2141.4 (566.0) kcal/day, p<0.0001), and to report allergy (45.5% v 23%, p<0.0001).

The characteristics of the study population are shown in table 1 according to asthma status. Women who suffered from asthma had a larger BMI, smoked less, and reported more frequent use of dietary supplements and allergy than women who did not have asthma.

Table 1 Characteristics of the population according to asthma status

	Asthmatic women (n = 2145)	Control women (n = 66 390)	p value*
Age (years)			
<48	30.6%	31.6%	
48–53	24.5%	24.4%	
53–58	20.9%	20.2%	0.8
58–63	13.6%	13.8%	
>63	10.4%	10.0%	
Physical activity			
1 st quartile	26.1%	25.7%	
2 nd quartile	22.7%	25.3%	
3 rd quartile	24.8%	24.2%	0.04
4 th quartile	26.4%	24.8%	
Caloric intake (kcal/day)			
1 st quartile (<1737)	24.2%	24.5%	
2 nd quartile (1737–2088)	24.2%	25.5%	
3 rd quartile (2088–2490)	25.4%	25.5%	0.2
4 th quartile (>2490)	26.2%	24.5%	
Body mass index (kg/m ²)			
<20	12.8%	15.5%	
20–25	60.9%	64.3%	0.0001
25–30	20.2%	16.6%	
>30	6.1%	3.6%	
Tobacco consumption			
Never smokers	61.4%	64.7%	
Ex-smokers	27.8%	22.6%	0.0001
Current smokers	10.8%	12.7%	
Menopausal status			
Premenopausal	12.0%	12.0%	
Postmenopausal	79.6%	79.8%	0.9
Perimenopausal	8.4%	8.2%	
Education/number of years at school			
<12	0.5%	0.5%	
12	11.0%	11.4%	
14	49.1%	51.2%	0.3
16	20.4%	19.0%	
17	19.0%	17.9%	
Vitamin supplementation use	25.0%	21.1%	0.0001
Allergy	76.4%	21.3%	0.0001

* χ^2 test.

Table 2 presents the dietary intake of specific fruits and vegetables among prevalent cases of asthma and control women. Caloric intake was higher among women reporting asthma. No significant difference in food intake was observed.

Table 2 Mean (SD) fruit and vegetable consumption (g/day) according to asthma status (E3N study, France)

	Asthmatic women (n = 2145)	Control women (n = 66 390)	p value*
Total caloric intake	2167.70 (585.27)	2140.50 (565.39)	0.03
Vegetables†			
Leafy vegetables	67.02 (40.35)	68.05 (39.37)	0.2
Fruity vegetables	69.09 (59.52)	69.81 (56.61)	0.6
Root vegetables	29.42 (26.01)	30.00 (25.29)	0.3
Cabbages	22.75 (23.48)	22.70 (22.86)	0.9
Tomatoes	19.41 (16.49)	19.48 (15.65)	0.8
Carrots	16.21 (15.06)	16.75 (14.85)	0.09
Fruits†			
Fruits	265.05 (175.82)	260.36 (167.30)	0.2
Fruits with citric	48.21 (52.33)	48.61 (50.03)	0.7
Fruits with β -carotene	47.22 (47.73)	47.65 (46.46)	0.7
Nuts and seeds	5.46 (8.09)	5.49 (8.38)	0.9
Apples	20.11 (23.29)	20.81 (22.65)	0.15

*p value from *t* test conducted on log transformed data.

†Food groups: leafy vegetables (chicory, lettuce and spinach), fruity vegetables (artichoke, avocado, tomatoes, cucumber, green beans, eggplants, bell pepper, and zucchini), root vegetables (carrots, radish, beetroot, celeriac, and salsify), fruit (orange, grapefruit, mandarin, apple, pear, banana, kiwi, pineapple, strawberries or raspberries, cherry, peach, melon, apricot, raisins, plums, other fruits), and nuts and seeds.

Table 3 shows the associations between quartiles of fruit and vegetable intake and asthma in adulthood. After adjusting for BMI, physical activity, menopausal status, smoking, and use of dietary supplements, a decreasing trend in asthma prevalence was observed with increasing intake of apples (OR_{Q1-Q4} 0.90, 95% CI 0.80 to 1.02, test for trend *p* = 0.03). Other fruit intake (fruits, citrus fruits, fruits rich in β -carotene) was not related to the prevalence of asthma.

Leafy vegetable intake was inversely related to asthma prevalence with a significant decreasing trend when intake was increased. Women in the highest quartile of leafy vegetable intake (>90.0 g/day) had a 22.0% lower risk of reporting asthma than those in the lowest quartile (\leq 39.3 g/day) (OR 0.82 (95% CI 0.73 to 0.93), test for trend *p* = 0.0009). Intake of fruity vegetables was inversely associated with asthma (test for trend *p* = 0.01); women in the highest quartile of tomato intake (>28.2 g/day) had a 17.6% lower risk of having asthma than those in the lowest quartile (\leq 6.4 g/day) (OR_{Q1-Q4} 0.85 (95% CI 0.75 to 0.96), test for trend *p* = 0.02). Increased intake of root vegetables was also inversely related to asthma prevalence (test for trend *p* = 0.004). Women in the highest quartile of carrot intake (>24.9 g/day) had a 20.0% decrease in the risk of having asthma (OR_{Q1-Q4} 0.81 (95% CI 0.72 to 0.92), test for trend *p* = 0.0003) compared with those with low carrot intake (\leq 3.8 g/day). No significant association was observed with cabbage intake. Similar results were found in multi-food models including leafy vegetables and tomatoes or leafy vegetables and carrots and the potential confounding factors previously mentioned, which suggests that intake of leafy vegetables, tomatoes and carrots may independently play a beneficial role on the prevalence of asthma.

Twenty six foods were studied: leafy vegetables, chicory, lettuce, spinach, fruity vegetables, artichoke, avocado, tomatoes, cucumber, green beans, eggplants, bell pepper, zucchini, root vegetables, carrots, radish, beetroot, celeriac, salsify, fruits, fruits with citrus, fruits with β -carotene, nuts and seeds, apples, blue fish, and other fish. The Bonferroni correction was therefore applied by multiplying by 26 the *p* value presented in the tables. Two of the three main associations remained statistically significant; *p* values of 0.0009 (leafy vegetables), 0.02 (tomatoes) and 0.0003 (carrots) became 0.02, 0.5 and 0.008, respectively.

Table 3 Association of fruit and vegetable intake and asthma prevalence in adulthood

Food groups and foods†	Quartile of intake (g/day)				p test for trend
	Q ₁	Q ₂	Q ₃	Q ₄	
Fruit	≤145.3	>145.3–235.4	>235.4–333.6	>333.6	
Cases (n)	543	522	515	565	
Controls (n)	16339	16700	16789	16562	
Age adjusted OR (95% CI)	1.00	0.94 (0.83 to 1.06)	0.92 (0.81 to 1.04)	1.02 (0.90 to 1.15)	0.8
Multivariate OR (95% CI)*	1.00	0.93 (0.82 to 1.05)	0.91 (0.81 to 1.03)	0.98 (0.87 to 1.11)	0.7
Citrus fruits	≤1.2	>1.2–38.4	>38.4–73.3	>73.3	
Cases (n)	556	538	511	540	
Controls (n)	16375	16606	16745	16664	
Age adjusted OR (95% CI)	1.00	0.96 (0.85 to 1.08)	0.90 (0.80 to 1.02)	0.95 (0.84 to 1.07)	0.3
Multivariate OR (95% CI)*	1.00	0.97 (0.86 to 1.10)	0.90 (0.80 to 1.02)	0.94 (0.83 to 1.06)	0.2
Fruits rich in β-carotene	≤2.6	>2.6–40.1	>40.1–72.1	>72.1	
Cases (n)	556	534	523	532	
Controls (n)	16373	16617	16848	16552	
Age adjusted OR (95% CI)	1.00	0.95 (0.84 to 1.07)	0.91 (0.81 to 1.03)	0.94 (0.84 to 1.07)	0.3
Multivariate OR (95% CI)*	1.00	0.96 (0.85 to 1.08)	0.92 (0.81 to 1.04)	0.94 (0.83 to 1.06)	0.2
Apple	0	>0–16.2	>16.2–31.2	>31.2	
Cases (n)	572	560	498	515	
Controls (n)	16969	16305	16452	16664	
Age adjusted OR (95% CI)	1.00	1.02 (0.91 to 1.15)	0.90 (0.80 to 1.02)	0.91 (0.81 to 1.03)	0.04
Multivariate OR (95% CI)*	1.00	1.04 (0.92 to 1.17)	0.90 (0.80 to 1.02)	0.90 (0.80 to 1.02)	0.03
Nuts and seeds	0	>0–2.9	>2.9–31.2	>31.2	
Cases (n)	631	507	461	546	
Controls (n)	18776	15891	15080	16643	
Age adjusted OR (95% CI)	1.00	0.95 (0.85 to 1.07)	0.91 (0.81 to 1.03)	0.98 (0.87 to 1.11)	0.6
Multivariate OR (95% CI)*	1.00	0.95 (0.85 to 1.07)	0.90 (0.79 to 1.01)	0.93 (0.82 to 1.05)	0.16
Leafy vegetables	≤39.3	>39.3–62.9	>62.9–90.0	>90.0	
Cases (n)	572	542	516	515	
Controls (n)	16335	16694	16797	16564	
Age adjusted OR (95% CI)	1.00	0.93 (0.82 to 1.05)	0.88 (0.78 to 0.99)	0.89 (0.79 to 1.00)	0.03
Multivariate OR (95% CI)*	1.00	0.91 (0.81 to 1.02)	0.85 (0.75 to 0.95)	0.82 (0.73 to 0.93)	0.0009
Fruity vegetables	≤24.9	>24.9–64.8	>64.8–101.6	>101.6	
Cases (n)	578	529	501	537	
Controls (n)	16316	16711	16747	16616	
Age adjusted OR (95% CI)	1.00	0.89 (0.79 to 1.01)	0.85 (0.75 to 0.95)	0.91 (0.81 to 1.03)	0.08
Multivariate OR (95% CI)*	1.00	0.92 (0.81 to 1.04)	0.84 (0.74 to 0.95)	0.87 (0.77 to 0.99)	0.01
Tomatoes	≤6.4	>6.4–17.9	>17.9–28.2	>28.2	
Cases (n)	570	516	546	513	
Controls (n)	16357	16689	16719	16625	
Age adjusted OR (95% CI)	1.00	0.90 (0.79 to 1.00)	0.94 (0.83 to 1.06)	0.89 (0.79 to 1.00)	0.11
Multivariate OR (95% CI)*	1.00	0.91 (0.80 to 1.02)	0.93 (0.83 to 1.05)	0.85 (0.75 to 0.96)	0.02
Root vegetables	≤8.6	>8.6–27.5	>27.5–44.6	>44.6	
Cases (n)	571	552	498	524	
Controls (n)	16331	16673	16757	16629	
Age adjusted OR (95% CI)	1.00	0.95 (0.84 to 1.07)	0.85 (0.75 to 0.96)	0.90 (0.80 to 1.02)	0.03
Multivariate OR (95% CI)*	1.00	0.97 (0.86 to 1.09)	0.85 (0.75 to 0.96)	0.86 (0.77 to 0.98)	0.004
Carrots	≤3.8	>3.8–14.9	>14.9–24.9	>24.9	
Cases (n)	576	557	517	495	
Controls (n)	16313	16710	16698	16669	
Age adjusted OR (95% CI)	1.00	0.95 (0.84 to 1.06)	0.88 (0.78 to 0.99)	0.84 (0.74 to 0.95)	0.002
Multivariate OR (95% CI)*	1.00	0.97 (0.86 to 1.09)	0.86 (0.78 to 0.99)	0.81 (0.72 to 0.92)	0.0003
Cabbage	≤1.3	>1.3–18.1	>18.1–34.8	>34.8	
Cases (n)	568	520	505	552	
Controls (n)	16358	16653	16705	16674	
Age adjusted OR (95% CI)	1.00	0.90 (0.80 to 1.02)	0.87 (0.77 to 0.98)	0.95 (0.85 to 1.07)	0.3
Multivariate OR (95% CI)*	1.00	0.93 (0.82 to 1.05)	0.87 (0.77 to 0.98)	0.91 (0.81 to 1.03)	0.07

*Adjusted for age, total caloric intake, BMI, physical activity, smoking status, menopausal status, and vitamin supplementation use.

†Food groups: leafy vegetables (chicory, lettuce and spinach), fruity vegetables (artichoke, avocado, tomatoes, cucumber, green beans, eggplants, bell pepper, and zucchini), root vegetables (carrots, radish, beetroot, celeriac, and salsify), fruit (orange, grapefruit, mandarin, apple, pear, banana, kiwi, pineapple, strawberries or raspberries, cherry, peach, melon, apricot, raisins, plums, other fruits) and nuts and seeds.

We calculated the association between specific vegetable intake and asthma prevalence for non-smokers, ex-smokers, and current smokers because smokers are known to have a lower serum antioxidant level²⁶ and higher antioxidant requirements²⁷ (table 4). The beneficial effect of leafy vegetable, tomato and carrot intake was slightly stronger in women who reported smoking than in non-smokers, although the small number of asthma cases reporting smoking limited the power of the analysis.

Table 4 Association of leafy vegetables, tomatoes and carrots with asthma prevalence in adulthood according to smoking status

Food groups and foods†	Quartile of intake (g/day)				p test for trend
	Q ₁	Q ₂	Q ₃	Q ₄	
Leafy vegetables	≤39.3	>39.3–62.9	>62.9–90.0	>90.0	
Non-smokers					
Cases (n)	342	324	312	313	
Controls (n)	10369	10693	10781	10249	
Age adjusted OR (95% CI)	1.00	0.92 (0.79 to 1.07)	0.88 (0.75 to 1.03)	0.93 (0.79 to 1.08)	0.2
Multivariate OR (95% CI)*	1.00	0.90 (0.77 to 1.05)	0.85 (0.72 to 0.99)	0.86 (0.74 to 1.01)	0.04
Ex-smokers					
Cases (n)	148	155	137	144	
Controls (n)	3409	3636	3715	3948	
Age adjusted OR (95% CI)	1.00	0.98 (0.78 to 1.24)	0.85 (0.67 to 1.07)	0.83 (0.66 to 1.05)	0.06
Multivariate OR (95% CI)*	1.00	0.97 (0.77 to 1.22)	0.82 (0.65 to 1.05)	0.78 (0.61 to 0.99)	0.01
Current smokers					
Cases (n)	73	52	57	47	
Controls (n)	2246	2042	1977	2018	
Age adjusted OR (95% CI)	1.00	0.78 (0.54 to 1.12)	0.89 (0.62 to 1.26)	0.72 (0.49 to 1.04)	0.13
Multivariate OR (95% CI)*	1.00	0.78 (0.54 to 1.11)	0.88 (0.62 to 1.25)	0.69 (0.47 to 1.00)	0.10
Tomatoes	≤6.4	>6.4–17.9	>17.9–28.2	>28.2	
Non-smokers					
Cases (n)	327	322	336	306	
Controls (n)	10253	10752	10631	10456	
Age adjusted OR (95% CI)	1.00	0.94 (0.80 to 1.10)	0.99 (0.85 to 1.16)	0.92 (0.78 to 1.07)	0.4
Multivariate OR (95% CI)*	1.00	0.96 (0.82 to 1.12)	0.98 (0.85 to 1.16)	0.88 (0.75 to 1.03)	0.17
Ex-smokers					
Cases (n)	154	132	154	144	
Controls (n)	3540	3558	3798	3812	
Age adjusted OR (95% CI)	1.00	0.86 (0.68 to 1.09)	0.94 (0.75 to 1.18)	0.88 (0.69 to 1.10)	0.4
Multivariate OR (95% CI)*	1.00	0.88 (0.69 to 1.11)	0.95 (0.75 to 1.19)	0.85 (0.67 to 1.07)	0.2
Current smokers					
Cases (n)	79	52	44	54	
Controls (n)	2232	2054	1982	2015	
Age adjusted OR (95% CI)	1.00	0.72 (0.50 to 1.02)	0.63 (0.43 to 0.91)	0.75 (0.53 to 1.07)	0.07
Multivariate OR (95% CI)*	1.00	0.74 (0.52 to 1.06)	0.63 (0.43 to 0.92)	0.75 (0.52 to 1.07)	0.06
Carrots	≤3.8	>3.8–14.9	>14.9–24.9	>24.9	
Non-smokers					
Cases (n)	337	334	315	305	
Controls (n)	10002	10467	10812	10811	
Age adjusted OR (95% CI)	1.00	0.95 (0.81 to 1.10)	0.87 (0.74 to 1.01)	0.84 (0.72 to 0.98)	0.01
Multivariate OR (95% CI)*	1.00	0.97 (0.83 to 1.13)	0.87 (0.74 to 1.02)	0.81 (0.69 to 0.95)	0.004
Ex-smokers					
Cases (n)	153	148	145	138	
Controls (n)	3606	3627	3717	3758	
Age adjusted OR (95% CI)	1.00	0.97 (0.77 to 1.22)	0.92 (0.73 to 1.16)	0.86 (0.68 to 1.09)	0.18
Multivariate OR (95% CI)*	1.00	0.99 (0.79 to 1.25)	0.93 (0.73 to 1.17)	0.83 (0.66 to 1.06)	0.10
Current smokers					
Cases (n)	76	63	45	45	
Controls (n)	2360	2285	1863	1775	
Age adjusted OR (95% CI)	1.00	0.85 (0.61 to 1.20)	0.75 (0.52 to 1.09)	0.78 (0.54 to 1.14)	0.13
Multivariate OR (95% CI)*	1.00	0.88 (0.63 to 1.24)	0.76 (0.52 to 1.11)	0.78 (0.53 to 1.14)	0.12

*Adjusted for age, total caloric intake, BMI, physical activity, menopausal status, and vitamin supplementation use.

†Food groups: leafy vegetables (chicory, lettuce and spinach).

DISCUSSION

In this large population of women asthma was less frequent in those with a higher intake of leafy vegetables, tomatoes and carrots, and significant dose responses were observed. The effect persisted after adjusting for potential confounding factors. After correction for multiple comparisons, the main result for leafy vegetables and carrots remained statistically significant. This suggests that intake of specific vegetables may play a beneficial role against the prevalence of asthma in our population.

The strengths of our study include the large number of prevalent cases of asthma, a wide variability in dietary intake,²⁸ and the use of a validated dietary questionnaire with a very detailed assessment. In addition, the questionnaire has proved useful for evaluating the association between dietary intake and cancer.²⁹ However, the cross sectional nature of our analysis limits the interpretation of our results, given that women with asthma might have modified their dietary pattern toward a more “healthy diet” or have avoided some foods because of their symptoms. We do not have information on change of diet over time and therefore cannot assess this. However, it is unlikely that women with asthma symptoms will decrease their intake of tomatoes, carrots, or leafy vegetable because of their symptoms, given that these foods are not known to produce allergic reactions. On the other hand, if women increased their intake of these vegetables, this would tend to underestimate the association; our estimates are therefore likely to be conservative. Correction for multiple comparisons using a conservative approach showed that the main result for leafy vegetables and carrots remained statistically significant.

Women who reported using dietary supplements (n = 14 826) were on average thinner, were more likely to report allergy, and ate significantly more fruit and vegetables than those who did not report using supplements. However, we did not observe a significant interaction between supplement use and the effect of fruit and vegetable intake on asthma. In addition, in a sub-analysis which included only women who did not report using dietary supplements, the results remained similar. We also evaluated the association of food intake and asthma prevalence stratified by time since the first asthma attack because responses to the dietary questionnaire may be influenced by the most recent food intake.¹³ A similar association was observed for each time period, suggesting that the dietary pattern had been almost constant during adulthood, supporting a potential causal association—for example, for leafy vegetable intake with a 20 year delay since first attack (OR_{Q1-Q4} 0.74, 95% CI 0.61 to 0.90, test for trend p = 0.003).

Analyses were conducted in detail for fruit and vegetables in relation to hypotheses regarding protection by antioxidants,¹ and for fish in relation to hypotheses regarding anti-inflammatory aspects of omega 3.¹ We adjusted for total caloric intake and conducted two food models including, in turn, two of the following foods—leafy vegetables, tomatoes, or carrots—but we were not able to adjust for other dietary factors that might be relevant for asthma apart from fish for which we did not find a significant association.⁸ We therefore cannot completely exclude the possibility of confounding by other dietary factors. Although this is a limitation, if confounding is present it should be from dietary factors that exert an even more powerful effect than vegetables. In addition, the decreasing prevalence of asthma with increasing intake of leafy vegetables and carrots provides further evidence for a causal association. We also considered other potential confounding factors, in particular BMI, menopausal status and smoking status. BMI has been related to the frequency of asthma in this³⁰ and other populations,³¹ and a diet poor in fruit and vegetables is usually related to a higher BMI. Lack of control for this variable might therefore have biased the association. Similarly, female reproductive hormones are thought to play a role in asthma²¹⁻²³ (although controversy exists) and should be taken into account. Ex-smokers had a higher frequency of asthma than those who had never smoked, which is in agreement with other studies showing that, as asthma developed, subjects stopped smoking.³² Although our models were adjusted for smoking, there is a possibility for residual confounding by this variable given that smokers are in general more likely to have a diet poor in fruit and vegetables, as observed in our study.

Our definition of asthma was based on one question regarding the occurrence of asthma attacks and the given date of the first attack. Because this information was not confirmed by a physician, we further validated responses with a more detailed questionnaire derived from the ATS questionnaire¹⁴ and observed a high concordance. Although self-reported asthma in older women may have included chronic obstructive pulmonary disease (COPD), 61.4% of the cases reported never smoking and are therefore unlikely to have COPD. However, given that we did not measure pulmonary function, we cannot exclude the possibility of COPD in older women, particularly in ex and current smokers. The attributable risk of atopy to asthma is a matter of debate,³³ asthma may occur without atopy whereas COPD and allergy can occur together. However, the high proportion of women with asthma who reported allergy (76%) represents an argument in favour of the diagnosis. Some

women may have had childhood asthma which, because it remitted and relapsed in adulthood, was classified as adult asthma. These women are likely to report food intake in the same way as women with asthma starting in adulthood. A reporting error, if any, would therefore be random and lead to an underestimation of the association.

One biological mechanism which may be associated with the beneficial effect of vegetable intake on asthma is the effect of antioxidant vitamins¹⁻⁸ such as carotenoids. Carrots and leafy vegetables are rich in carotenoids (α -carotene, β -carotene, lutein and zeaxanthin) which protect cells from oxidative stress.^{34,35} Tomato juice, carrot juice and spinach powder have been shown to increase plasma levels of cystolic glutathione transferase (GST- π), an effect attributed to the upregulation of GST activity by carotenoids.³⁶ Carotenoids are known to modulate immune function by enhancing lymphocyte proliferation and cytotoxic T cell activity, and could modulate the production of PGE₂ by altering the activation of the arachidonic acid cascade from which PGE₂ is derived. In addition, folate—for which leafy vegetables are a major source—has a role in DNA repair and is likely to potentiate the role of antioxidant enzymes.³⁶ Smokers are exposed to an extra burden of oxidative stress and have a significantly lower serum level of antioxidants.²⁶ It is expected that the protective effect of leafy vegetables and carrots should be greater in women who smoke. This is suggested by our data, but the small number of women in this category limits the power of our analysis.

Our findings on the protective effect of vegetables with a high content of carotenoids are consistent with data from a prospective study of asthma onset among women.⁷ Other studies have reported no association between carotenoid intake and asthma, but most of the studies focused only on β -carotene intake and did not assess the impact of other carotenoids.^{1,8} As reported in other studies, we did not observe a protective effect from citrus fruits;^{4,7,9,37} foods rich in vitamin C have been more consistently related to lung function than respiratory symptoms.¹ Caution is needed, however, when interpreting the results since an infrequent consumption of fruit and vegetables may be an indicator of a poor diet and unhealthy habits. We cannot therefore totally exclude residual confounding by life style.

Fruit and vegetables contain thousands of biologically active phytochemicals, but food composition tables often lack this information and the content of specific nutrients may also be subject to error. Focusing on food intake might be more readily applicable for dietary recommendations and might also enhance our understanding of the relation between phytochemicals and asthma. Public health messages need to be simple and recommendations for food intake are easier to follow than recommended dietary intakes of specific nutrients. This has led to the standard recommendation of five portions of fruit and vegetables a day.¹⁰ The World Health Organization has recommended a daily intake of at least 400 g of fruit and vegetables per day, but studies of median fruit and vegetable intake in 10 European countries have shown that a considerable number of consumers failed to meet this target.¹⁷ In our population, 27% of the women had a fruit and vegetable intake under 400 g/day; 65.3% had an intake of <200 g/day leafy, fruity and root vegetables, and 93.4% had an intake of <200 g/day leafy vegetables, tomatoes, and carrots. For the implementation of public health policies, it is important to know which fruit and vegetables might be more likely to prevent specific diseases.

Our results suggest that a higher intake of leafy vegetables, carrots, and tomatoes may have a beneficial effect against adult asthma. Further studies are needed to confirm these results and to consider the effect of fruit and vegetable intake on the severity of the disease.

Competing interests: none declared.

References

- 1 Romieu I, Trenga C. Diet and obstructive lung diseases. *Epidemiol Rev* 2001;23:268–87.
- 2 Heffner JE, Repine JE. Pulmonary strategies of antioxidant defense. *Am Rev Respir Dis* 1989;140:531–54.
- 3 Repine JE, Bast A, Lankhorst I. Oxidative stress in chronic obstructive pulmonary disease. Oxidative Stress Study Group. *Am J Respir Crit Care Med* 1997;156:341–57.
- 4 Kelly Y, Sacker A, Marmot M. Nutrition and respiratory health in adults: findings from the health survey for Scotland. *Eur Respir J* 2003;21:664–71.
- 5 Miedema I, Feskens EJ, Heederik D, et al. Dietary determinants of long-term incidence of chronic nonspecific lung diseases. The Zutphen Study. *Am J Epidemiol* 1993;138:37–45.

- 6 Forastiere F, Pistelli R, Sestini P, et al. Consumption of fresh fruit rich in vitamin C and wheezing symptoms in children. SIDRIA Collaborative Group, Italy (Italian Studies on Respiratory Disorders in Children and the Environment). *Thorax* 2000;55:283–8.
- 7 Troisi RJ, Willett WC, Weiss ST, et al. A prospective study of diet and adult-onset asthma. *Am J Respir Crit Care Med* 1995;151:1401–8.
- 8 McKeever TM, Britton J. Diet and asthma. *Am J Respir Crit Care Med* 2004;170:725–9.
- 9 Shaheen SO, Sterne JA, Thompson RL, et al. Dietary antioxidants and asthma in adults: population-based case-control study. *Am J Respir Crit Care Med* 2001;164:1823–8.
- 10 Potter JD, Finnegan JR, Guinard JX, et al. 5 a day for better health program evaluation report, NIH Publication N0 01-4904. Bethesda, MD: National Cancer Institute, 2000.
- 11 Clavel-Chapelon F, van Liere MJ, Giubout C, et al. E3N, a French cohort study on cancer risk factors. E3N Group. Etude Epidemiologique auprès de femmes de l'Education Nationale. *Eur J Cancer Prev* 1997;6:473–8.
- 12 Bertheke PG, de Vente W, Kemper HC, et al. Longitudinal trends in and tracking of energy and nutrient intake over 20 years in a Dutch cohort of men and women between 13 and 33 years of age: the Amsterdam Growth and Health longitudinal study. *Br J Nutr* 2001;85:375–85.
- 13 Willett WC. Recall of remote diet. In: Willett WC, ed. *Nutritional epidemiology*. New York: Oxford University Press, 1998:148–56.
- 14 Ferris BG. Epidemiology Standardization Project (American Thoracic Society). *Am Rev Respir Dis* 1978;118:1–120.
- 15 Lucas F, Niravong M, Villemainot S, et al. Estimation of food portion size using photographs: Relative validity, strengths, weaknesses and recommendations. *J Human Nutr Diet* 1995;8:65–74.
- 16 Van Liere MJ, Lucas F, Clavel F, et al. Relative validity and reproducibility of a French dietary history questionnaire. *Int J Epidemiol* 1997;26:S128–36.
- 17 Agudo A, Slimani N, Ocke MC, et al. Consumption of vegetables, fruit and other plant foods in the European Prospective Investigation into Cancer and Nutrition (EPIC) cohorts from 10 European countries. *Public Health Nutr* 2002;5:1179–96.
- 18 James WO, Schofield EC. *Human requirements: a manual for planners and nutritionists*. New York: Oxford University Press, 1990.
- 19 Zar J. *Biostatistical analysis*, 3rd. ed. Upper Saddle River, NJ: Prentice-Hall, 1996.
- 20 Kleinbaum DG. *Logistic regression*. New York: Springer, 1994.
- 21 Troisi RJ, Speizer FE, Willett WC, et al. Menopause, postmenopausal estrogen preparations, and the risk of adult-onset asthma. A prospective cohort study. *Am J Respir Crit Care Med* 1995;152:1183–8.
- 22 Barr RG, Camargo Jr CA. Hormone replacement therapy and obstructive airway diseases. *Treat Respir Med* 2004;3:1–7.
- 23 Barr RG, Wentowski CC, Grodstein F, et al. Prospective study of postmenopausal hormone use and newly diagnosed asthma and chronic obstructive pulmonary disease. *Arch Intern Med* 2004;164:379–86.
- 24 Bonferroni CE. *Teoria statistica delle classi e calcolo delle probabilità*, Pubblicazioni del R Istituto Superiore di Scienze Economiche e Commerciali di Firenze, 1936:3–62.
- 25 Woodward M. *Epidemiology study design and data analysis*. Text in statistical science. Florida, USA: Chapman & Hall/CRC, 1999.
- 26 Bendich A. Role of antioxidants in the maintenance of immune functions. In: Frei B, ed. *Natural antioxidant in human health and disease*. New York: Academic Press, 1994:447–467.
- 27 Institute of Medicine. *Dietary reference intakes for vitamin C, vitamin E, selenium, carotenoids*. Washington, DC: National Academy Press, 2000.
- 28 Kesse E, Boutron-Ruault MC, Clavel-Chapelon F. Regional dietary habits of French women born between 1925 and 1950. *Eur J Nutr* 2004;44:285–92.
- 29 Miller AB, Altenburg HP, Bueno-de-Mesquita B, et al. Fruits and vegetables and lung cancer: Findings from the European Prospective Investigation into Cancer and Nutrition. *Int J Cancer* 2004;108:269–76.
- 30 Romieu I, Avenel V, Leynaert B, et al. Body mass index, change in body silhouette, and risk of asthma in the E3N cohort study. *Am J Epidemiol* 2003;158:165–74.
- 31 Chinn S. Asthma and obesity: where are we now? *Thorax* 2003;58:1008–10.
- 32 Siroux VPI, Oryszczyn MP, Le Moual N, et al. Relationships of active smoking to asthma and asthma severity in the EGEA study. *Eur Respir J* 2000;15:470–7.
- 33 Pearce N, Pekkanen J, Beasley R. How much asthma is really attributable to atopy? *Thorax* 1999;54:268–72.
- 34 Sies H. Strategies of antioxidant defense. *Eur J Biochem* 1993;215:213–9.
- 35 Di Mascio P, Kaiser S, Sies H. Lycopene as the most efficient biological carotenoid singlet oxygen quencher. *Arch Biochem Biophys* 1989;274:532–8.
- 36 Lindsay DG, Astley SB. European research on the functional effects of dietary antioxidant: EUROFEDA. *Mol Aspects Med* 2002;23:1–38.
- 37 Soutar A, Seaton A, Brown K. Bronchial reactivity and dietary antioxidants. *Thorax* 1997;52:166–70.