

Phthalates impair germ cell number in the mouse fetal testis by an androgen- and estrogen-independent mechanism.

Abdelali Lehraiki, Chrystèle Racine, Andrée Krust, René Habert, Christine Levacher

► **To cite this version:**

Abdelali Lehraiki, Chrystèle Racine, Andrée Krust, René Habert, Christine Levacher. Phthalates impair germ cell number in the mouse fetal testis by an androgen- and estrogen-independent mechanism.. Toxicological Sciences, Oxford University Press (OUP), 2009, 111 (2), pp.372-82. <10.1093/toxsci/kfp153>. <inserm-00438655>

HAL Id: inserm-00438655

<http://www.hal.inserm.fr/inserm-00438655>

Submitted on 9 Dec 2009

HAL is a multi-disciplinary open access archive for the deposit and dissemination of scientific research documents, whether they are published or not. The documents may come from teaching and research institutions in France or abroad, or from public or private research centers.

L'archive ouverte pluridisciplinaire **HAL**, est destinée au dépôt et à la diffusion de documents scientifiques de niveau recherche, publiés ou non, émanant des établissements d'enseignement et de recherche français ou étrangers, des laboratoires publics ou privés.

Phthalates Impair Germ Cell Number in the Mouse Fetal Testis by an Androgen- and Estrogen-Independent Mechanism

Abdelali Lehraiki,*†‡ Chrystèle Racine,*†‡ Andrée Krust,§ René Habert,*†‡ and Christine Levacher*†‡¹

*Laboratory of Gonad Differentiation and Radiobiology, Stem Cells and Radiation Service, Institute of Cellular and Molecular Radiation Biology, Life Sciences Division, Commissariat à l'Energie Atomique, BP 6, 92265 Fontenay-aux-Roses, France; †Université Paris Diderot-Paris 7, BP 6, 92265 Fontenay-aux-Roses, France; ‡Unité 967, INSERM, BP 6, 92265 Fontenay-aux-Roses, France; and §Institut de Génétique et de Biologie Moléculaire et Cellulaire, (CNRS, INSERM, UDS, Collège de France), BP 10142, 67404 Illkirch-Strasbourg, France

Received June 10, 2009; accepted July 7, 2009

Data from experiments conducted almost exclusively in the rat have established that some phthalates have deleterious effects on the fetal testis probably due to their antiandrogenic and/or estrogenic effects, but their mechanisms of action remain unknown. A recent study reported that phthalates also have deleterious effects on human fetal testis with germ cell number, but not steroidogenesis altered. Therefore, we used organ culture of fetal testes at different stages of development to analyze the direct effects of phthalates on both steroidogenesis and gonocyte development and to determine if the effects of MEHP on these functions reported in the rat can be extended to other mammalian species. We defined specific periods of sensitivity of the fetal mouse testis to MEHP for these two functions and showed that the effects of phthalates on steroidogenesis vary with the developmental stage. Conversely, the strong deleterious effects of phthalates on germ cells were constantly present during the active phases of gonocyte development and thus share no relationship with the steroidogenic status. Moreover, all the effects of phthalates were unchanged in testes from mice deficient for estrogen (ER α KO or ER β KO) or androgen (Tfm) receptors. In conclusion, our results demonstrate that phthalates impair mouse fetal germ cell number similarly to other mammalian species, but are neither estrogenic nor antiandrogenic molecules because their effects do not involve, directly or indirectly, ER or AR.

Key Words: MEHP; phthalates; fetal testis; gonocytes; steroidogenesis; Leydig cell; development; estrogen receptor; androgen receptor.

Phthalate esters are a class of environmental endocrine disrupting chemicals which are mainly used as plasticizers in PVC plastics. They are found in numerous consumer goods and, because they are not covalently bound to the plastic product, they can leach out over time from these products and

can be ingested (Silva *et al.*, 2004). After absorption, phthalates are rapidly hydrolyzed by esterases in the gut and other tissues into a monoester, which is the active molecule (Latini, 2005). For example, di-(2-ethylhexyl) phthalate (DEHP), one of the most widely used phthalates, is metabolized to its monoester metabolite, mono-(2-ethylhexyl) phthalate (MEHP), which is a recognized active testicular toxicant (Fisher, 2004).

Numerous studies in the rat have shown that *in utero* exposure to phthalates results in male reproductive disorders including altered seminiferous cord formation, multinucleated gonocyte (MNG) formation, epididymal agenesis, nipple retention, reduced ano-genital distance, hypospadias and cryptorchidism (Foster, 2006; Gray *et al.*, 2006), and also reduction of fetal testosterone production and Leydig cell Ins13 gene expression (Lehmann *et al.*, 2004; McKinnell *et al.*, 2005; Sharpe and Skakkebaek, 2008), all these effects attesting to a profound alteration of testis development.

During fetal and neonatal development of the testis (Jost and Magre, 1993; Olaso and Habert, 2000), after migration of the primordial germ cells (PGCs) from extra-embryonic mesoderm to the genital ridge at embryonic day 11.5 (E11.5) in the mouse (Yoshimizu *et al.*, 2001), the differentiating Sertoli cells (Wilhelm *et al.*, 2007) surround the germ cells, therefore called gonocytes, to form the seminiferous cords from E12.5. The gonocytes divide actively, many of them undergoing apoptosis by E12.5–14.5 (Wang *et al.*, 1998) and then enter a quiescent period (Nagano *et al.*, 2000; Vergouwen *et al.*, 1991). After birth, they resume mitosis while a second wave of apoptosis occurs (Boulogne *et al.*, 2003; Wang *et al.*, 1998) (Fig. 1) and start to differentiate into spermatogonia. The interstitial region contains mesenchymal cells and steroid-secreting Leydig cells from E12.5 (Livera *et al.*, 2006; O'Shaughnessy *et al.*, 2005). Interactions between the different cell lineages occur early in morphogenesis of the testis and are crucial for its normal development. Androgens mediate a wide range of developmental and physiological responses that are critical for the male reproductive and nonreproductive system (Sharpe, 2006; Welsh *et al.*, 2008). During fetal life, testis development is

¹ To whom correspondence should be addressed at Unité mixte "Cellules Souches et Radiations", Inserm U967/CEA/Université Paris-Diderot, LDRG/SCSR/IRCM/DSV, Centre CEA, BP6 18 Route du Panorama, Bat. 05A RdC, 92265 Fontenay-Aux-Roses, France. Fax: +33-1-46-54-99-06. E-mail: christine.levacher@cea.fr.




FIG. 1. Schematic representation of the development of gonocytes during mouse fetal and neonatal life in relation to the timing and duration of the organ cultures as described in “Material and Methods.”

physiologically modulated by, among other factors (Olaso and Habert, 2000), estrogen and androgen because estrogen receptor beta (ER β) and androgen receptor (AR) inactivation lead to an increase in germ cell number (Delbes *et al.*, 2004; Merlet *et al.*, 2007), whereas testosterone production is enhanced by ER α inactivation (Delbes *et al.*, 2005).

Until now, the mechanisms of action of phthalates have not been elucidated. MEHP activates peroxisome proliferators-activated receptor (PPAR- α) in cell transactivation assays (Corton and Lapinskas, 2005) but phthalate-treated PPAR- α -/- mice develop toxic lesions in the testis (Ward *et al.*, 1998), suggesting that phthalates can act through PPAR- α -independent pathways in mediating testicular toxicity (Bhattacharya *et al.*, 2005). In addition, phthalates show little estrogenic activity (Harris *et al.*, 1997; Takeuchi *et al.*, 2005), but can also reduce estradiol production (Lovekamp-Swan and Davis, 2003), and there is a growing consensus that they are antiandrogenic (Hu *et al.*, 2009). However, phthalates and their mono-phthalate metabolites do not bind to AR (Parks *et al.*, 2000), indicating that they are not direct AR antagonists. Most studies investigating the *in utero* effects of phthalates have been performed in the rat *in vivo* and have focused on the phthalate-induced suppression of testosterone production and Leydig cell aggregation (Fisher *et al.*, 2003; Mylchreest *et al.*, 1999). This has led to a hypothesized causal relationship between reduced testosterone production, altered fetal testicular development and reduced sperm counts in adulthood (Scott *et al.*, 2007; Skakkebaek *et al.*, 2001). However, an *in vitro* study from our laboratory on human fetal testis evidenced a reduction of the number of gonocytes by MEHP in the absence of any alteration of testosterone production (Lambrot *et al.*, 2009). A recent study in the mouse demonstrated also that, as in the rat, *in utero* DBP exposure increases the number of MNGs, but unlike the rat this response occurs in the absence of measurable disruption of testicular testosterone concentrations (Gaido *et al.*, 2007).

Therefore, we used organ culture of fetal mouse testis to establish that there is no relationship between phthalate-induced alterations of fetal steroidogenesis and gonocyte development, unlike what has been concluded from observations in the rat. Furthermore, using mice deficient for AR or ER, we showed that the phthalate-induced alterations are independent of these receptors.

MATERIAL AND METHODS

Chemicals and solutions. The culture medium was phenol red-free Dulbecco’s Modified Eagle Medium/Ham F12 (1:1) (Invitrogen, Carlsbad, CA) supplemented with 80 μ g/ml gentamicin (Invitrogen). Ovine LH (oLH; NIH.LH S19; 1.01 IU/mg) was a gift from Dr A.F. Parlow (National Institute of Diabetes and Digestive and Kidney Diseases, Bethesda, MD). MEHP was from TCI Europe (Antwerp, Belgium). A stock solution (4mM) was made up in dimethyl sulfoxide and then diluted in culture medium. db-cAMP (N6, 2’-O-dibutyryl-cAMP) was from Sigma Aldrich (St Louis, MO).

Animals. C57Bl/6 and transgenic mice were housed under controlled photoperiod (lights on 08:00-20:00) with *ad libitum* access to tap water and a soy and alfalfa-free breeding diet (Global diet 2019, Harlan Teklad, Indianapolis, IN). Mice lacking ER α (ER α -/-) or ER β (ER β -/-) were produced by Dupont *et al.* (2000) and generously provided by Pierre Chambon (Institut de Genetique et de Biologie Moleculaire et Cellulaire, Illkirch, France). Androgen-insensitive *Tfm* mice (C57BL/6J-Aw-J.Cg-EdaTa-6J_-ArTfm, *Tfm*/Y) were from the Jackson Laboratory (Bar Harbor, ME). Transgenic animals were genotyped as described previously (Delbes *et al.*, 2004, 2005; Merlet *et al.*, 2007). Males were caged with females overnight and the day following overnight mating was counted as E0.5. Pregnant mice were killed by cervical dislocation on E13.5, E15.5, or E18.5 and the fetuses were quickly removed from the uterus. Fetuses were dissected under a binocular microscope, their sex was determined on the basis of the gonad morphology and the testes were collected from male fetuses. All animal studies were conducted in accordance with the NIH Guide for Care and Use of Laboratory Animals.

Organ culture and treatment. Organ cultures were performed as previously described (Livera *et al.*, 2006). Briefly, intact E13.5 testes associated with their mesonephros were placed on 10-mm-diameter Millicell CM filters (Millipore, Billerica, MA) (pore size 0.45 μ m). Testes from E15.5 and E18.5 were isolated and cut into six and eight pieces, respectively, and all the pieces from the same testis were placed on a single Millicell filter. The filter bearing the pieces of testes was floated on 0.4 ml of culture medium in tissue culture dishes and incubated at 37°C, in humidified atmosphere containing 95% air/5% CO $_2$. The medium was changed every 24 h and the culture was pursued for 1 or 3 days at E13.5 and for 3 days at E15.5 and E18.5 (Fig. 1). The testes were cultured in the presence or absence of oLH (100 ng/ml) and the effect of 20 or 200 μ M MEHP was estimated by comparing one testis cultured in the presence of MEHP with the other testis from the same fetus cultured in control medium. At the end of culture, testes were fixed for 1 h at room temperature in Bouin’s fluid, embedded in paraffin, and cut into 5- μ m sections. For RNA analysis, testes were immediately dry frozen with liquid nitrogen and stored at -80°C. The entire media were kept at -20°C until testosterone radioimmunoassay. The data were obtained from at least three independently repeated cultures with fetuses from different litters.

Immunohistochemistry. Serial sections were mounted on slides, deparaffinized and rehydrated. They were immunostained by a standard procedure as previously described (Delbes *et al.*, 2007). To unmask the MVH (mouse vasa homolog) protein, sections were microwaved at 750 W for 5 min and at 450W for 3 min in 10mM citrate buffer solution (pH 6.0). For all immunohistological procedures, slides were incubated in 0.3% H $_2$ O $_2$ for 15 min and in 5% bovine serum albumin in phosphate-buffered saline for 30 min to block nonspecific antigens and were then incubated with the primary antibody (rabbit polyclonal anti-AMH (anti-Müllerian hormone) antibody (1/500; Santa Cruz Biotechnology, Santa Cruz, CA), the rabbit polyclonal anti- β HSD antibody (1/5000; provided by Prof A. Payne, Stanford University Medical Center) or the rabbit anti-MVH antibody (1/500; Abcam, Cambridge, UK) overnight at 4°C. The primary antibody was detected by incubation with an appropriate biotinylated secondary antibody (1:200; Vector Laboratories, Burlingame, CA) for a further 30 min at room temperature followed by incubation with avidin-biotin-peroxidase complex (Vector Laboratories) for 30 min. DAB (Vector Laboratories) was used as the chromogen and hematoxylin as the nuclear counterstained. Negative controls were done by omitting the primary antibody.

Testicular cell counting. Germ cells and Leydig cells were identified by immunohistochemical detection of MVH and 3 β HSD, respectively. The counting was done as previously described (Delbes *et al.*, 2004). All counts were performed blind using Histolab analysis software (Microvision Instruments, Evry, France).

Measurement of germ cell apoptosis. Because cleaved caspase-3 is involved in most of the apoptotic pathway, we chose its detection as a marker of apoptosis (Delbes *et al.*, 2004). Gonocytes were distinguished by combining staining for AMH (a Sertoli-specific marker) and cleaved caspase-3 (rabbit polyclonal anti-cleaved caspase-3 1:100; Cell Signaling Technology, Beverly, MA). Cleaved caspase-3 was localized first using DAB (Vector Laboratories) and AMH was then detected using VIP substrate (Vector Laboratories). The percentage of apoptotic germ cells was determined by counting the stained and unstained gonocytes in all sections mounted.

Measurement of BrdU incorporation index. The percentage of germ cells in S-phase was evaluated by measuring the BrdU (5-bromo-2'-deoxyuridine) incorporation by immunohistochemical methods using the Cell Proliferation Kit (GE Healthcare, Buckinghamshire, UK) according to the manufacturer's recommendations. BrdU (1%) was added to the culture medium 3 h before the end of culture. BrdU incorporation into proliferating cells was detected by immunocytochemistry using a mouse anti-BrdU antibody and a peroxidase-linked anti-mouse IgG. Sertoli cells and gonocytes were distinguished by combining staining for AMH (a Sertoli-specific marker) and BrdU. BrdU localization was performed first using DAB (Vector Laboratories) and AMH was then detected using VIP substrate (Vector Laboratories). The BrdU incorporation index was obtained by a blind counting of at least 1000 Sertoli cell or 400 gonocyte stained and unstained nuclei in five nonconsecutive sections from every treatment group experiment. Histolab analysis software was used for counting (Microvision Instruments, Evry, France).

Measurement of testosterone production. Testosterone secretion into the medium was determined in duplicate by direct radioimmunoassay, without extraction as previously described (Habert and Picon, 1984).

RNA extraction and reverse transcription. Total RNA was extracted using the RNeasy Plus mini-Kit (Qiagen, Courtaboeuf, France) and reverse transcribed using the High Capacity cDNA Reverse Transcription Kit (Applied Biosystems, Courtaboeuf, France) according to the manufacturer's instructions.

Real-time quantitative PCR. Q-PCRs were performed on an ABI PRISM 7000 Sequence Detector System using a TaqMan PCR Master Mix (Applied Biosystems). The primers and probes used were assays on demand designed by Applied Biosystems (Table 1). The primers and probes used for 3 β HSD and external standard luciferase were described (Baker and O'Shaughnessy 2001; Merlet *et al.*, 2007). Q-PCR was performed and analyzed as previously described (Merlet *et al.*, 2007). Results were analyzed by using the delta-delta Ct method with the use of luciferase as a normalized control (Merlet *et al.*, 2007). For each treatment, the mRNA levels were determined in samples from three different litters and at least from six to nine testes.

Statistical analysis. All values are expressed as means \pm SEM. The statistical significance of the difference between the mean values for the treated and untreated testes from the same fetus was evaluated using the paired Student's *t*-test. The unpaired Student's *t*-test was used to evaluate the difference between the mean values for two different genotypes.

RESULTS

Effects of MEHP on Leydig Cells

Basal and LH-stimulated testosterone production. In wild-type control testes, as shown in Figure 2, the presence of 200 μ M MEHP during 3 days of culture in basal medium induced a high and significant time-dependent stimulation of

TABLE 1
Primers Used in Real-Time RT-PCR

Gene full name	Gene symbol	Applied Biosystem accession number
3-Hydroxy-3-methylglutaryl coenzyme A reductase	HMG-CoA reductase	Mm01282499_m1
Steroidogenic acute regulatory protein	StAR	Mm00441558_m1
P450 side-chain cleavage enzyme	Cyp11a1	Mm 00490735_m1
P450 17 α hydroxylase	Cyp17a1	Mm00484040_m1
Steroid 5-alpha reductase	Srd5a1	Mm00614213_m1
P450 aromatase	Cyp19a1	Mm00484049_m1
Luteinizing hormone receptor (LH-R)	Lhcgr	Mm00442931_m1
Insulin-like growth factor 3	Insl-3	Mm01340353_m1

testosterone production in E13.5 and E18.5 testes. When the testes were cultured in the presence of 100 ng/ml oLH, testosterone production was increased fourfold in E13.5 testes and more than 10-fold in E18.5 testes compared with basal conditions. In these conditions, the daily testosterone production in E13.5 testes was unaffected by 200 μ M MEHP, but was significantly decreased in a time-dependent manner in E18.5 testes with a strong inhibition on D3.

As expected (Delbes *et al.*, 2005), testosterone production was significantly higher in ER α -/- animals than in respective wild-type controls in basal conditions. The invalidation of ER α did not modify the stimulatory and inhibitory effects of MEHP on testosterone secretion in E18.5 testes in basal and LH-stimulated conditions, respectively (Fig. 2).

Number of Leydig cells. Compared with controls, the number of Leydig cells was not significantly modified after exposure to 200 μ M MEHP in the presence or absence of LH at all the studied stages (Fig. 3A). In addition, no abnormal distribution or clustering of the Leydig cells was observed in any condition (Fig. 3B).

Leydig cell gene expression. Because MEHP had no effect on Leydig cell number, we hypothesized that its effect on steroidogenesis was due to the modulation of the expression of genes coding for proteins involved in cholesterol and testosterone biosynthesis or metabolism. Therefore, mRNA expression of key genes was analyzed by real-time PCR (Fig. 4).

In basal conditions, significant increases in gene expression likely to explain the stimulation of testosterone production by MEHP were observed at the level of HMG-CoA reductase and Cyp17a1 in E13.5 testes and StAR and Cyp17a1 in E18.5 testes (Fig. 4). In LH-stimulated conditions, the decrease in testosterone production in E18.5 testes was related to a 40–50% reduction of the expression levels of all the genes tested, except 5-alpha-reductase, whereas they were not significantly altered in E13.5 testes (Fig. 4) in agreement with the lack of




FIG. 2. Effect of MEHP on testosterone secretion by fetal testes from wild-type and ER α -/- mice in organ culture. Testes at E13.5 (upper panels) and E18.5 (lower panels) were cultured for 3 days (D1–D3) in control medium or in the presence of 200 μ M MEHP. Cultures were performed in the absence (left panels) or presence (right panels) of 100 ng/ml oLH as indicated. Values are mean \pm SEM from 4 to 18 cultures. * p < 0.05, ** p < 0.01, *** p < 0.001 versus corresponding control value in the paired Student's t -test. $^{\circ}p$ < 0.05 in the unpaired Student's t -test for different genotypes.

change in testosterone production. Aromatase (Cyp19a1) and 5-alpha-reductase are required in the conversion of testosterone to estradiol and dihydrotestosterone, respectively, and could modulate the amount of testosterone secreted. The expression of aromatase was decreased in both the absence and presence of LH at all ages studied (Fig. 4), but the expression of 5-alpha-reductase was never modified. The expression of LH-R (LH receptor, Lhcgr) in E13.5 and E18.5 testes was not modified by MEHP in the absence of LH in the culture medium, but was

significantly reduced in its presence (Fig. 4). Interestingly, the expression of Insl-3, the protein which plays a major role in testis descent, was decreased by MEHP in all studied conditions except in E13.5 testes cultured in the absence of LH (Fig. 4).

db-cAMP-stimulated testosterone production. In an attempt to elucidate whether the decrease in LH receptor expression was involved in the inhibition of LH-stimulated




FIG. 3. Effect of MEHP on the number and histological appearance of Leydig cells in mouse testes in organ culture. (A) Number of Leydig cells in E13.5 and E18.5 testes after 3 days of culture in the absence (white bars) or presence of 200 μ M MEHP (black bars) and in the absence or presence of 100 ng/ml oLH. Values are mean \pm SEM from four to five cultures. There was no significant difference between control and MEHP. (B) Leydig cells (arrows), as revealed by immunostaining of 3 β -HSD, in mouse testes at E18.5 after 3 days of culture in control medium or in the presence of 200 μ M MEHP. No morphological alteration or distribution was detectable after MEHP exposure.




FIG. 4. Quantitative real-time PCR analysis of the effect of MEHP (200 μ M) on mRNA levels of a selected set of genes involved in Leydig cell cholesterol metabolism and steroidogenesis, of LH-R (*Lhcgr*) and *Insl3* genes in E13.5 (white bars) and E18.5 (black bars) testes after 3 days of culture in the absence (A) or presence (B) of 100 ng/ml oLH. The results were calculated by the delta-delta Ct method using an external standard (luciferase) added during RNA extraction as an endogenous reference. The levels of mRNA are expressed as mean of relative unit \pm SEM of six to nine cultures with the control animals having a value of 1, with each sample processed in duplicate. * $p < 0.05$, ** $p < 0.01$, *** $p < 0.001$ versus corresponding control value in the paired Student's *t*-test.

testosterone production by MEHP at E18.5, we evaluated the effect of MEHP on db-cAMP-stimulated testosterone production. The inhibition of testosterone production by MEHP (Fig. 5) was similar to that obtained with LH, which means that it exerts its rate-limiting inhibitory effect downstream from LH-R.

Effects of MEHP on Sertoli Cell Proliferation and Function

MEHP did not alter the proliferation of Sertoli cells because the BrdU-labeling index was not modified after 3-day treatment with 200 μ M MEHP at any age studied (results not shown). However, immunostaining of AMH clearly showed that its Sertoli cell content was markedly decreased compared with control in E13.5, E15.5, and E18.5 testes after 3 days of treatment with 200 μ M MEHP (Fig. 6).




FIG. 5. Testosterone secretion by E18.5 testes cultured in the presence of oLH or db-cAMP and in the absence (white bars) or presence (black bars) of 200 μ M MEHP. Values are mean \pm SEM of testosterone secretion on day 3 of culture ($n = 11$ –18 cultures). ** $p < 0.01$, *** $p < 0.001$, versus control value in the paired Student's *t*-test.

Effects of MEHP on Germ Cells

Gonocyte distribution and morphology. After 3 days of culture, the integrity of the seminiferous cord structure was maintained in both control and treated testes at all fetal and neonatal stages studied and at all concentrations of MEHP tested (Fig. 7). However, with 200 μ M MEHP all the gonocytes have disappeared in E13.5dpc testes (see further) and, in E15.5 and E18.5 testes, the gonocytes stayed aggregated in the center of the cord (Figs. 6E and 6F).

MNGs occurred spontaneously in E15.5 and E18.5 control testes after 3 days of culture, that is, during and after the quiescent period, and their number was increased significantly by 200 μ M MEHP with the greatest increase observed in E18.5 testes (Fig. 7A). E18.5 testes from ER β (ER $\beta^{-/-}$) or AR (Tfm) deficient mice, showed similar increase in MNCs (for ER $\beta^{-/-}$: 1.84 \pm 0.33% in control vs. 8.79 \pm 0.94 in MEHP treated, $n = 9$; for Tfm: 2.15 \pm 0.52% in control vs. 12.5 \pm 2.91 in MEHP treated, $n = 4$) than the wild-type (Fig. 7A: 2.02 \pm 0.32% in control vs. 8.79 \pm 0.89 in MEHP treated, $n = 9$). In E13.5 testes, no MNG was detected after 1 or 3 days of culture in control conditions or after 24-h treatment with 20 μ M MEHP (Fig. 7A).

Number of gonocytes. Because MEHP affected testosterone production differently in the presence and absence of LH, we evaluated its effect on gonocytes in these two conditions (Fig. 7B). The presence of LH and, consequently, a high level of testosterone production, did not modify the number of gonocytes in E13.5 and E18.5 control cultures. In contrast, MEHP severely reduced the number of gonocytes in any condition at all the stages studied, except with E15.5 testes cultured for 3 days i.e. during the quiescent period. At E13.5, the gonocytes had completely disappeared after 3 days of culture (D3) in the presence of 200 μ M MEHP and the number of gonocytes was reduced by 40% by 20 μ M MEHP after one day of culture (D1). With E18.5 testes, 200 μ M MEHP reduced the




FIG. 6. AMH immunostaining in mouse testes at E13.5 (A, D), E15.5 (B, E), E18.5 (C, F) cultured for 3 days in control medium (A, B, C) or in the presence of 200µM MEHP (D, E, F). Black arrow: mononucleated gonocyte; orange arrow: MNGs; arrowhead: Sertoli cell.

number of gonocytes by ~40–50% after 3 days of culture (D3), that is, 24 h after resuming mitosis and apoptosis. As expected (Delbes *et al.*, 2004), in testes from E18.5 the number of gonocytes was higher in ERβ^{-/-} mice than in wild-type mice (ERβ^{+/+}) after 3 days of culture. After MEHP treatment, the number of gonocytes was reduced in the same manner in E18.5 testes from wild-type mice and from ERβ^{-/-} or Tfm mice (Fig. 8). Taken together, these results demonstrate that MEHP reduces the number of germ cells throughout fetal/neonatal life except during the quiescent period, independently of the level of testosterone production and of ER and AR pathways.

Germ cell proliferation and apoptosis. The decrease in the number of gonocytes in MEHP-exposed testes could be induced by an increased rate of apoptosis or by a reduced rate of proliferation. The effect of MEHP on proliferation and apoptosis was therefore studied by assessing the percentage of BrdU (Fig. 7D) and cleaved caspase 3-positive (Fig. 7C) gonocytes, respectively.

MEHP greatly increased the rate of apoptotic gonocytes at all the ages studied (Fig. 7C). At the same time, the proliferation of the gonocytes was increased by 40% in E13.5 after 24-h exposure to 20µM MEHP (Fig. 7D), but was reduced by 60% in E18.5 testes after 3 days of exposure to 200µM MEHP. Taken together, these results show that MEHP reduces the number of gonocytes essentially by increasing apoptosis.

DISCUSSION

We have previously shown that ER and AR are involved in fetal testis development (Delbes *et al.*, 2004, 2005; Merlet

et al., 2007). Using an organotypic culture system of fetal testis from wild-type mice or from mice deficient for ER or AR, we present here the first demonstration that phthalate-induced alterations of both steroidogenesis and germ cell number do not involve ER and AR.

Steroidogenesis

Our findings show that testicular alteration of steroidogenesis by phthalates during fetal life is very complex. We report here for the first time a stimulatory effect of MEHP on testosterone production in fetal mouse testis and that MEHP can stimulate, inhibit or have no effect on testosterone production depending on the age of the testis at explantation and on the culture conditions. These results differ from those of numerous experiments conducted in the rat that have led to the conclusion that the main deleterious effect of phthalates concerns their antiandrogenic capacity, because testosterone production by the fetal testis is reduced after *in utero* (Fisher *et al.*, 2003; Foster, 2006) or *in vitro* exposure (Chauvigné *et al.*, 2009; Hallmark *et al.*, 2007) to phthalates. Furthermore, recent experiments on *in utero* exposure to DBP in the mouse (Gaido *et al.*, 2007) and on *in vitro* exposure of fetal human testis to MBP (Hallmark *et al.*, 2007) or MEHP (Lambrot *et al.*, 2009) failed to observe any inhibition of testosterone production or changes in the expression of the steroidogenic enzymes. Moreover, phthalates can stimulate steroidogenesis in prepubertal testis *in vivo* (Lin *et al.*, 2008), in mouse gonadal cell lines or primary Leydig cell cultures (Ge *et al.*, 2007). In all these cases, the mechanisms involved have not been fully elucidated.

In the present study, the MEHP-induced stimulation of steroidogenesis can be ascribed mainly to an increased




FIG. 7. Effect of MEHP on the development of the gonocytes in fetal testes in organ culture. Testes were cultured for 1 day (D1) or 3 days (D3) at E13.5 and 3 days at E15.5 and E18.5 in control medium (white bars) or in the presence of 20µM (gray bars) or 200µM (black bars) MEHP and in the presence or absence of 100 ng/ml oLH as indicated. At the end of the culture, MNGs (A) and total gonocytes (B) were counted as described in Material and Methods. The percentage of cleaved caspase-3-positive gonocytes (C) was used to measure apoptotic activity and the percentage of BrdU-positive gonocytes (D) to evaluate proliferating index. Values are mean \pm SEM from four to nine cultures. * $p < 0.05$, ** $p < 0.01$, *** $p < 0.001$ versus corresponding control value in the paired Student's *t*-test.

transcription of p450C17 and also of HMGCoA-reductase, as reported in gonadal cell lines (Gunnarsson *et al.*, 2008) or of StAR, depending on the developmental stage. In contrast, the inhibitory effect of MEHP on steroidogenesis was obtained by a reduction of the level of the mRNAs of the main genes involved in testosterone biosynthesis while, in the rat it was mainly ascribed to p450C17 and P450sc (Chauvigné *et al.*, 2009; Culty *et al.*, 2008; Hallmark *et al.*, 2007). Additionally, in E18.5 testes, MEHP reduced the expression of LH-R, but this does not seem to be involved in the reduction of steroidogenesis. The reduction of aromatase activity could participate in the stimulation of testosterone production in the absence of LH, that is, in conditions of low testosterone production, mostly by reducing the inhibition of steroidogenesis by estradiol (Delbes *et al.*, 2006, 2007) via ER α (Delbes *et al.*, 2005), the reduction of testosterone transformation being probably negligible because intratesticular estradiol concentration is very low (Delbes *et al.*, 2004). However, we show here that the ER α pathway (Delbes *et al.*, 2006) is involved neither in the stimulatory nor in the inhibitory effects of MEHP on testosterone secretion.

The present results provide new insights into the mechanisms underlying the effects of phthalates on fetal testicular steroidogenesis, because they prove that they are independent of the ER pathway but depend on other factors: 1) the developmental stage, which modulates the maturation and the limiting steps of the steroidogenic pathway, the LH level and the cholesterol availability (present results; Gunnarsson *et al.*, 2008; Hallmark *et al.*, 2007); 2) the species because, *in vitro*, unlike the rat (Chauvigné *et al.*, 2009) and the results we present here in the mouse, human fetal testis is not sensitive to phthalate in terms of steroidogenesis (Hallmark *et al.*, 2007; Lambrot *et al.*, 2009). Aggregation of Leydig cells and even their presence in the seminiferous tubules have also been reported to be a prominent consequence of *in utero* (Lin *et al.*, 2008; Mahood *et al.*, 2005) or *in vitro* (Chauvigné *et al.*, 2009; Hallmark *et al.*, 2007) exposure of rat fetal testis to phthalates, but was not observed in the present experiments on mouse testis *in vitro* or *in vivo* (Gaido *et al.*, 2007) or in human fetal testis *in vitro* (Hallmark *et al.*, 2007; Lambrot *et al.*, 2009).

The importance of an alteration of Leydig cell status during fetal life is related to its possible role in inducing




FIG. 8. Involvement of ER and AR in the effects of MEHP on the number of gonocytes in fetal testes. Testes at E18.5 from wild-type, ERβ^{-/-} and Tfm mice were cultured for 3 days in control medium (white bars) or in the presence of 200μM MEHP (black bars). At the end of the culture, total gonocytes were counted as described in Material and Methods. Values are mean ± SEM from four to nine cultures. **p* < 0.05, ***p* < 0.01 versus corresponding control value in the paired Student's *t*-test for wild-type mice; °*p* < 0.05 in the unpaired Student's *t*-test for two different genotypes.

cryptorchidism and hypospadias (Foster *et al.*, 2001; Henley and Korach, 2006) and concerns not only testosterone but also Insl3 production. Underdeveloped gubernaculum (Barlow and Foster, 2003) and reduced Insl3 expression have been observed following fetal exposure to phthalates in male rats (Lague and Tremblay, 2008; Lehmann *et al.*, 2004; McKinnell *et al.*, 2005). Indeed, Insl3 mRNA was also reduced in the mouse testes after MEHP exposure, except in E13.5 testis in basal conditions when maturation/stimulation of Leydig cells is low.

Gonocyte Development

Our findings clearly show that phthalates strongly impair germ cell development in the fetal mouse testis *in vitro* during critical time-specific windows. In our model, MEHP reduces the number of gonocytes by inducing massive apoptosis during early fetal and late fetal/neonatal life, in relation to the high mitotic and apoptotic activity during these periods. During the quiescent phase between E15.5 and E19.5 (day of birth), due to the naturally low apoptotic activity of the gonocytes, phthalates did not significantly reduce their number. This can be compared with the low sensitivity to retinoic acid (Livera *et al.*, 2000) and transforming growth factor (TGFβ1) (Olaso *et al.*, 1998) of rat germ cells during their quiescent period. These results are in accordance with the few studies that have previously focused on the modification of the number of gonocytes by phthalates in other species, with the mouse being particularly sensitive: in the rat *in vivo*, the number of gonocytes is reduced at E21.5 when the animals are treated from E13.5, but not when they are treated later (Ferrara *et al.*, 2006); *in vitro* MEHP reduces the number of gonocytes in the fetal testis of E14.5 (Chauvigné *et al.*, 2009) and of PN3 neonate (Li and Kim, 2003); lastly, phthalates have been shown recently to reduce

the number of gonocytes in human fetal testis *in vitro* at an early stage (Lambrot *et al.*, 2009).

Apoptosis could result from a direct effect of phthalates on the gonocytes or from an indirect effect via other cell types. The fact that the deleterious effects of MEHP on the gonocytes are similar whether it stimulates or inhibits steroidogenesis suggests that they do not involve antiandrogenic effects as postulated in the rat (Chauvigné *et al.*, 2009; Scott *et al.*, 2007; Skakkebaek *et al.*, 2001). This hypothesis was strengthened by the fact that MEHP reduces the number of gonocytes in a similar fashion in Tfm mice. The gonocytes of ERβKO mice were also similarly affected, demonstrating that MEHP does not also act via ER, directly, or indirectly via the modulation of aromatase. Therefore, even though we demonstrated previously that ERβ and AR pathways physiologically control fetal germ cell number (Delbes *et al.*, 2004; Merlet *et al.*, 2007), we can definitively eliminate a direct or indirect effect of MEHP on fetal germ cells via antiandrogenic/estrogenic effects.

In the rat, germ cell apoptosis could also be induced by Sertoli cells (Fisher *et al.*, 2003; Kleymenova *et al.*, 2005) which are targets of MEHP because, although their proliferation is not affected, they exhibit a reduction in AMH content. Various nonexclusive pathways have been postulated to be involved in MEHP-induced germ cell apoptosis, such as the TRAIL (McKee *et al.*, 2006) and NF-kB (Rasoulpour and Boekelheide, 2005) pathways and the Fas/FasL system (Giammona *et al.*, 2002; Yao *et al.*, 2008). Lastly, our *in vitro* observations in the mouse that phthalates disturb gonocyte migration and increase the occurrence of MNGs in late fetal life are in agreement with *in vivo* observations in the mouse (Gaido *et al.*, 2007) and rat (Ferrara *et al.*, 2006; Kleymenova *et al.*, 2005).

The phthalate-induced alteration in the number of gonocytes at the fetal and neonatal stage could compromise subsequent reactivation of germ cell proliferation and development during puberty as in the rat (Ferrara *et al.*, 2006) and lead to altered fertility in adulthood. Furthermore, the impairment of differentiation of gonocytes is to some extent reminiscent of the failure of fetal gonocyte differentiation that is thought to lead to formation of CIS (carcinoma *in situ*) cells and subsequent testicular germ cell tumors in humans (Rajpert-De Meyts, 2006). Further studies are needed to establish if *in utero* exposure to phthalates leads, in the mouse, to altered fertility or occurrence of tumors in adulthood.

CONCLUSION

The present data define specific critical periods of sensitivity of fetal mouse testis to MEHP and clearly establish that the effects of phthalates on steroidogenesis are unrelated to those on gonocyte development. We demonstrated that the strong deleterious effect of phthalates on the germ cell lineage that is observed in various mammalian species is independent of any

antiandrogenic effects and that the phthalates are neither estrogenic nor antiandrogenic because their effects do not involve, directly or indirectly, ER or AR.

FUNDING

Université Paris Diderot, by the Institut National de la Santé et de la Recherche Médicale; Commissariat à l'Énergie Atomique; Agence Nationale de la Recherche Project Santé-Environnement et Santé-Travail (ANR2007SEST01901); and Ministère de l'Éducation Nationale de la Recherche et de la Technologie (237132006) fellowship to A.L.

ACKNOWLEDGMENTS

We thank Prof. P. Chambon for providing transgenic mice (Institut de Génétique et Biologie Moléculaire et Cellulaire, Illkirch, France) and Prof A. Payne (Stanford University Medical Center, CA) for providing the anti-3 β HSD antibody. We also thank C. Chamailard for expert technical assistance, V. Neuville, S. Leblay, and S. Rodrigues for animal care and A. Gouret for secretarial assistance.

REFERENCES

- Baker, P. J., and O'Shaughnessy, P. J. (2001). Role of gonadotrophins in regulating numbers of Leydig and Sertoli cells during fetal and postnatal development in mice. *Reproduction* **122**, 227–234.
- Baker, P. J., Pakarinen, P., Huhtaniemi, I. T., Abel, M. H., Charlton, H. M., Kumar, T. R., and O'Shaughnessy, P. J. (2003). Failure of normal Leydig cell development in follicle-stimulating hormone (FSH) receptor-deficient mice, but not FSH β -deficient mice: Role for constitutive FSH receptor activity. *Endocrinology* **144**, 138–145.
- Barlow, N. J., and Foster, P. M. (2003). Pathogenesis of male reproductive tract lesions from gestation through adulthood following in utero exposure to Di(n-butyl) phthalate. *Toxicol. Pathol.* **31**, 397–410.
- Bhattacharya, N., Dufour, J. M., Vo, M. N., Okita, J., Okita, R., and Kim, K. H. (2005). Differential effects of phthalates on the testis and the liver. *Biol. Reprod.* **72**, 745–754.
- Boulogne, B., Habert, R., and Levacher, C. (2003). Regulation of the proliferation of cocultured gonocytes and Sertoli cells by retinoids, triiodothyronine, and intracellular signaling factors: Differences between fetal and neonatal cells. *Mol. Reprod. Dev.* **65**, 194–203.
- Chauvigné, F., Menuet, A., Lesné, L., Chagnon, M.-C., Chevrier, C., Régnier, J.-F., Angerer, J., and Jégou, B. (2009). Time- and Dose-related effects of di-(2-ethylhexyl) phthalate and its main metabolites on the function of the rat fetal testis in vitro. *Environ. Health Perspect.* **117**, 515–521.
- Corton, J. C., and Lapinskas, P. J. (2005). Peroxisome proliferator-activated receptors: Mediators of phthalate ester-induced effects in the male reproductive tract? *Toxicol. Sci.* **83**, 4–17.
- Culty, M., Thuillier, R., Li, W., Wang, Y., Martinez-Arguelles, D. B., Benjamin, C. G., Triantafilou, K. M., Zirkin, B. R., and Papadopoulos, V. (2008). In utero exposure to di-(2-ethylhexyl) phthalate exerts both short-term and long-lasting suppressive effects on testosterone production in the rat. *Biol. Reprod.* **78**, 1018–1028.
- Delbes, G., Duquenne, C., Szenker, J., Taccoen, J., Habert, R., and Levacher, C. (2007). Developmental changes in testicular sensitivity to estrogens throughout fetal and neonatal life. *Toxicol. Sci.* **99**, 234–243.
- Delbes, G., Levacher, C., Duquenne, C., Racine, C., Pakarinen, P., and Habert, R. (2005). Endogenous estrogens inhibit mouse fetal Leydig cell development via estrogen receptor alpha. *Endocrinology* **146**, 2454–2461.
- Delbes, G., Levacher, C., and Habert, R. (2006). Estrogen effects on fetal and neonatal testicular development. *Reproduction* **132**, 527–538.
- Delbes, G., Levacher, C., Pairault, C., Racine, C., Duquenne, C., Krust, A., and Habert, R. (2004). Estrogen receptor {beta}-mediated inhibition of male germ cell line development in mice by endogenous estrogens during perinatal life. *Endocrinology* **145**, 3395–3403.
- Dupont, S., Krust, A., Gansmuller, A., Dierich, A., Chambon, P., and Mark, M. (2000). Effect of single and compound knockouts of estrogen receptors α (ER α) and β (ER β) on mouse reproductive phenotypes. *Development* **127**, 4277–4291.
- Ferrara, D., Hallmark, N., Scott, H., Brown, R., McKinnell, C., Mahood, I. K., and Sharpe, R. M. (2006). Acute and long-term effects of in utero exposure of rats to di(n-butyl) phthalate on testicular germ cell development and proliferation. *Endocrinology* **147**, 5352–5362.
- Fisher, J. S. (2004). Environmental anti-androgens and male reproductive health: Focus on phthalates and testicular dysgenesis syndrome. *Reproduction* **127**, 305–315.
- Fisher, J. S., Macpherson, S., Marchetti, N., and Sharpe, R. M. (2003). Human 'testicular dysgenesis syndrome': A possible model using in-utero exposure of the rat to dibutyl phthalate. *Hum. Reprod.* **18**, 1383–1394.
- Foster, P. M. (2006). Disruption of reproductive development in male rat offspring following in utero exposure to phthalate esters. *Int. J. Androl.* **29**, 140–147; discussion 181–185.
- Foster, P. M., Mylchreest, E., Gaido, K. W., and Sar, M. (2001). Effects of phthalate esters on the developing reproductive tract of male rats. *Hum. Reprod. Update* **7**, 231–235.
- Gaido, K. W., Hensley, J. B., Liu, D., Wallace, D. G., Borghoff, S., Johnson, K. J., Hall, S. J., and Boelkeheide, K. (2007). Fetal mouse phthalate exposure shows that Gonocyte multinucleation is not associated with decreased testicular testosterone. *Toxicol. Sci.* **97**, 491–503.
- Ge, R. S., Chen, G. R., Dong, Q., Akingbemi, B., Sottas, C. M., Santos, M., Sealfon, S. C., Bernard, D. J., and Hardy, M. P. (2007). Biphasic effects of postnatal exposure to diethylhexylphthalate on the timing of puberty in male rats. *J. Androl.* **28**, 513–520.
- Giammona, C. J., Sawhney, P., Chandrasekaran, Y., and Richburg, J. H. (2002). Death receptor response in rodent testis after mono-(2-ethylhexyl) phthalate exposure. *Toxicol. Appl. Pharmacol.* **185**, 119–127.
- Gray, L. E., Jr., Wilson, V. S., Stoker, T., Lambricht, C., Furr, J., Noriega, N., Howdeshell, K., Ankley, G. T., and Guillette, L. (2006). Adverse effects of environmental antiandrogens and androgens on reproductive development in mammals. *Int. J. Androl.* **29**, 96–104; discussion 105–108.
- Gunnarsson, D., Leffler, P., Ekwurtzel, E., Martinsson, G., Liu, K., and Selstam, G. (2008). Mono-(2-ethylhexyl) phthalate stimulates basal steroidogenesis by a cAMP-independent mechanism in mouse gonadal cells of both sexes. *Reproduction* **135**, 693–703.
- Habert, R., and Picon, R. (1984). Testosterone, dihydrotestosterone and estradiol 17 β levels in maternal and fetal plasma and in fetal testes in the rat. *J. Steroid. Biochem.* **21**, 193–198.
- Hallmark, N., Walker, M., McKinnell, C., Mahood, I. K., Scott, H., Bayne, R., Coutts, S., Anderson, R. A., Greig, I., Morris, K., et al. (2007). Effects of monobutyl and di(n-butyl) phthalate in vitro on steroidogenesis and Leydig cell aggregation in fetal testis explants from the rat: Comparison with effects in vivo in the fetal rat and neonatal marmoset and in vitro in the human. *Environ. Health Perspect.* **115**, 390–396.

- Harris, C. A., Henttu, P., Parker, M. G., and Sumpter, J. P. (1997). The estrogenic activity of phthalate esters in vitro. *Environ. Health Perspect.* **105**, 802–811.
- Henley, D. V., and Korach, K. S. (2006). Endocrine-disrupting chemicals use distinct mechanisms of action to modulate endocrine system function. *Endocrinology* **147**, S25–S32.
- Hu, G. X., Lian, Q. Q., Ge, R. S., Hardy, D. O., and Li, X. K. (2009). Phthalate-induced testicular dysgenesis syndrome: Leydig cell influence. *Trends Endocrinol. Metab.* **20**, 139–145.
- Jost, A., and Magre, S. (1993). Sexual differentiation. In *Reproduction in Mammals and Man* (C. Thibault, M. Levasseur, and R. H. F. Hunter, Eds.), pp. 197–226. Ellipses, Paris.
- Kleymenova, E., Swanson, C., Boekelheide, K., and Gaido, K. W. (2005). Exposure in utero to di(n-butyl) phthalate alters the vimentin cytoskeleton of fetal rat Sertoli cells and disrupts Sertoli cell-gonocyte contact. *Biol. Reprod.* **73**, 482–490.
- Lague, E., and Tremblay, J. J. (2008). Antagonistic effects of testosterone and the endocrine disruptor mono-(2-ethylhexyl) phthalate on INSL3 transcription in Leydig cells. *Endocrinology* **149**, 4688–4694.
- Lambrot, R., Muczynski, V., Lecureuil, C., Angenard, G., Coffigny, H., Pairault, C., Moison, D., Frydman, R., Habert, R., and Rouiller-Fabre, V. (2009). Phthalates impair germ cell development in the human fetal testis in vitro without change in testosterone production. *Environ. Health Perspect.* **117**, 32–37.
- Latini, G. (2005). Monitoring phthalate exposure in humans. *Clin. Chim. Acta* **361**, 20–29.
- Lehmann, K. P., Phillips, S., Sar, M., Foster, P. M., and Gaido, K. W. (2004). Dose-dependent alterations in gene expression and testosterone synthesis in the fetal testes of male rats exposed to di (n-butyl) phthalate. *Toxicol. Sci.* **81**, 60–68.
- Li, H., and Kim, K. H. (2003). Effects of mono-(2-ethylhexyl) phthalate on fetal and neonatal rat testis organ cultures. *Biol. Reprod.* **69**, 1964–1972.
- Lin, H., Ge, R. S., Chen, G. R., Hu, G. X., Dong, L., Lian, Q. Q., Hardy, D. O., Sottas, C. M., Li, X. K., and Hardy, M. P. (2008). Involvement of testicular growth factors in fetal Leydig cell aggregation after exposure to phthalate in utero. *Proc. Natl. Acad. Sci. U. S. A.* **105**, 7218–7222.
- Livera, G., Delbes, G., Pairault, C., Rouiller-Fabre, V., and Habert, R. (2006). Organotypic culture, a powerful model for studying rat and mouse fetal testis development. *Cell Tissue Res.* **324**, 507–521.
- Livera, G., Rouiller-Fabre, V., Durand, P., and Habert, R. (2000). Multiple effects of retinoids on the development of Sertoli, germ and Leydig cells of fetal and neonatal rat testis in culture. *Biol. Reprod.* **62**, 1303–1314.
- Lovekamp-Swan, T., and Davis, B. J. (2003). Mechanisms of phthalate ester toxicity in the female reproductive system. *Environ. Health Perspect.* **111**, 139–145.
- Mahood, I. K., Hallmark, N., McKinnell, C., Walker, M., Fisher, J. S., and Sharpe, R. M. (2005). Abnormal Leydig Cell aggregation in the fetal testis of rats exposed to di (n-butyl) phthalate and its possible role in testicular dysgenesis. *Endocrinology* **146**, 613–623.
- McKee, C. M., Ye, Y., and Richburg, J. H. (2006). Testicular germ cell sensitivity to TRAIL-induced apoptosis is dependent upon p53 expression and is synergistically enhanced by DR5 agonistic antibody treatment. *Apoptosis* **11**, 2237–2250.
- McKinnell, C., Sharpe, R. M., Mahood, K., Hallmark, N., Scott, H., Ivell, R., Staub, C., Jegou, B., Haag, F., Koch-Nolte, F., and Hartung, S. (2005). Expression of insulin-like factor 3 protein in the rat testis during fetal and postnatal development and in relation to cryptorchidism induced by in utero exposure to di (n-Butyl) phthalate. *Endocrinology* **146**, 4536–4544.
- Merlet, J., Racine, C., Moreau, E., Moreno, S. G., and Habert, R. (2007). Male fetal germ cells are targets for androgens that physiologically inhibit their proliferation. *Proc. Natl. Acad. Sci. U. S. A.* **104**, 3615–3620.
- Mylchreest, E., Sar, M., Cattley, R. C., and Foster, P. M. (1999). Disruption of androgen-regulated male reproductive development by di(n-butyl) phthalate during late gestation in rats is different from flutamide. *Toxicol. Appl. Pharmacol.* **156**, 81–95.
- Nagano, R., Tabata, S., Nakanishi, Y., Ohsako, S., Kurohmaru, M., and Hayashi, Y. (2000). Reproliferation and relocation of mouse male germ cells (gonocytes) during prespermatogenesis. *Anat. Rec.* **258**, 210–220.
- O’Shaughnessy, P. J., Baker, P. J., and Johnston, H. (2005). Neuroendocrine regulation of Leydig cell development. *Ann. N. Y. Acad. Sci.* **1061**, 109–119.
- Olaso, R., and Habert, R. (2000). Genetic and cellular analysis of male germ cell development. *J. Androl.* **21**, 497–511.
- Olaso, R., Pairault, C., Boulogne, B., Durand, P., and Habert, R. (1998). Transforming growth factor β 1 and β 2 reduce the number of gonocytes by increasing apoptosis. *Endocrinology* **139**, 733–740.
- Parks, L. G., Ostby, J. S., Lambright, C. R., Abbott, B. D., Klinefelter, G. R., Barlow, N. J., and Gray, L. E., Jr. (2000). The plasticizer diethylhexyl phthalate induces malformations by decreasing fetal testosterone synthesis during sexual differentiation in the male rat. *Toxicol. Sci.* **58**, 339–349.
- Rajpert-De Meyts, E. (2006). Developmental model for the pathogenesis of testicular carcinoma in situ: Genetic and environmental aspects. *Hum. Reprod. Update* **12**, 303–323.
- Rasoulpour, R. J., and Boekelheide, K. (2005). NF-kappaB is activated in the rat testis following exposure to mono-(2-ethylhexyl) phthalate. *Biol. Reprod.* **72**, 479–486.
- Scott, H. M., Hutchison, G. R., Mahood, I. K., Hallmark, N., Welsh, M., De Gendt, K., Verhoeven, G., O’Shaughnessy, P., and Sharpe, R. M. (2007). Role of androgens in fetal testis development and dysgenesis. *Endocrinology* **148**, 2027–2036.
- Sharpe, R. M. (2006). Pathways of endocrine disruption during male sexual differentiation and masculinization. *Best Pract. Res. Clin. Endocrinol. Metab.* **20**, 91–110.
- Sharpe, R. M., and Skakkebaek, N. E. (2008). Testicular dysgenesis syndrome: Mechanistic insights and potential new downstream effects. *Fertil. Steril.* **89**, e33–e38.
- Silva, M. J., Reidy, J. A., Herbert, A. R., Preau, J. L., Jr., Needham, L. L., and Calafat, A. M. (2004). Detection of phthalate metabolites in human amniotic fluid. *Bull. Environ. Contam. Toxicol.* **72**, 1226–1231.
- Skakkebaek, N. E., Rajpert-De Meyts, E., and Main, K. M. (2001). Testicular dysgenesis syndrome: An increasingly common developmental disorder with environmental aspects. *Hum. Reprod.* **16**, 972–978.
- Takeuchi, S., Iida, M., Kobayashi, S., Jin, K., Matsuda, T., and Kojima, H. (2005). Differential effects of phthalate esters on transcriptional activities via human estrogen receptors alpha and beta, and androgen receptor. *Toxicology* **210**, 223–233.
- Vergouwen, R. P., Jacobs, S. G., Huiskamp, R., Davids, J. A., and de Rooij, D. G. (1991). Proliferative activity of gonocytes, Sertoli cells and interstitial cells during testicular development in mice. *J. Reprod. Fertil.* **93**, 233–243.
- Wang, R. A., Nakane, P. K., and Koji, T. (1998). Autonomous cell death of mouse male germ cells during fetal and postnatal period. *Biol. Reprod.* **58**, 1250–1256.
- Ward, J. M., Peters, J. M., Perella, C. M., and Gonzalez, F. J. (1998). Receptor and nonreceptor-mediated organ-specific toxicity of di(2-ethylhexyl)phthalate (DEHP) in peroxisome proliferator-activated receptor alpha-null mice. *Toxicol. Pathol.* **26**, 240–246.
- Welsh, M., Saunders, P. T., Fiskens, M., Scott, H. M., Hutchison, G. R., Smith, L. B., and Sharpe, R. M. (2008). Identification in rats of a programming window for reproductive tract masculinization, disruption of which leads to hypospadias and cryptorchidism. *J. Clin. Invest.* **118**, 1479–1490.
- Wilhelm, D., Palmer, S., and Koopman, P. (2007). Sex determination and gonadal development in mammals. *Physiol. Rev.* **87**, 1–28.

Yao, P. L., Lin, Y. C., and Richburg, J. H. (2008). TNF Alpha-mediated disruption of spermatogenesis in response to Sertoli cell injury in rodents is partially regulated by MMP2. *Biol. Reprod* **80**, 581–589.

Yoshimizu, T., Obinata, M., and Matsui, Y. (2001). Stage-specific tissue and cell interactions play key roles in mouse germ cell specification. *Development* **128**, 481–490.