

# Expected and unexpected adverse effects H1N1 vaccination for health care workers in a university hospital.

Aurelien Dinh, Christine Lawrence, Jérôme Salomon, Alexis Descatha

### ▶ To cite this version:

Aurelien Dinh, Christine Lawrence, Jérôme Salomon, Alexis Descatha. Expected and unexpected adverse effects H1N1 vaccination for health care workers in a university hospital. Vaccine, 2010, 28 (9), pp.2063. 10.1016/j.vaccine.2009.12.030 . inserm-00452802

### HAL Id: inserm-00452802 https://inserm.hal.science/inserm-00452802

Submitted on 3 Feb 2010

**HAL** is a multi-disciplinary open access archive for the deposit and dissemination of scientific research documents, whether they are published or not. The documents may come from teaching and research institutions in France or abroad, or from public or private research centers.

L'archive ouverte pluridisciplinaire **HAL**, est destinée au dépôt et à la diffusion de documents scientifiques de niveau recherche, publiés ou non, émanant des établissements d'enseignement et de recherche français ou étrangers, des laboratoires publics ou privés.

## Expected and Unexpected adverse effects H1N1 vaccination for health care workers in a University Hospital

Aurélien Dinh <sup>1</sup>, Christine Lawrence <sup>2</sup>, Jérôme Salomon <sup>1,2</sup>, Alexis Descatha <sup>3</sup>

- 1. AP-HP, UVSQ, Infectiology Unit, Poincare Teaching Hospital, Garches, France
- 2. AP-HP, Hygiene Operational Team, Poincare Teaching Hospital, Garches, France
- 3. AP-HP, UVSQ, Occupational Health Unit-INSERM U687, Poincare Teaching Hospital, Garches, France.

### **Correspondence and reprints:**

Alexis Descatha, Unite de pathologie professionnelle, CHU Poincare, 104 bd R poincare, 92380 Garches, France.

Tel (33) 1 47 10 77 54; Fax (33) 1 47 10 77 68. E-mail: alexis.descatha@rpc.aphp.fr

- (i) All authors declare that the answer to the questions on your competing interest relating to the study are all no and therefore have nothing to declare
- (ii) The paper and the data **have not previously been published**, either in whole or in part, and that no similar paper is in press or under review elsewhere. The manuscript has been read and approved by all the authors.
- (iii) All authors believe that manuscript is an honest work and **concur with the submitted version of the manuscript and with the listing of the authors**.
- (iv) All authors have participated to the study, writing and corrections.
- (v) Type: letter (or short communication), 373 words, 6 references

Sir,

We aimed at evaluating the potential adverse effects of a massive vaccination in a short period of time in order to inform correctly suspicious health care workers (HCW).

Taking into account the recommendations of Centers for Disease Control & Prevention,[1, 2] most of governments are starting vaccination programs for HCW. In France, it has started on October the 20<sup>th</sup>, 2009 in university hospitals. Vaccination team in our university hospital was gathered around infectiologists, hygienists, occupational physicians, nurses and managers (n=12). Subjects working in units with "high risk unit" of flu were asked to be vaccinated first (target units, such as infectious disease units, pediatric units, intensive care units, emergency units).

During the first week of vaccination, the vaccination team worked 22 hours in 8 target units around 10 sessions including night shift. The Pandemrix© vaccine was used (Influenza A H1N1 2009 Monovalent AS03-Adjuvanted Vaccine, GSK). Only 129 HCW were vaccinated this first week among 1870 HCW of the target population (6.9%). A self-administered questionnaire was filled out by this population about the vaccine adverse effects. We received 95 answers (73.6%). We observed an important local reaction (redness or important pain) in 50 cases (52.6%), 25 systemic reactions without fever (muscle or joint aches, 26.3%) and 6 with fever (>38°C, 6.3%). No severe allergic reaction has been observed. The mean and median was at 2 days [1;7]. Even though we observed more often adverse effects than the usual flu vaccine especially for systemic reaction [3, 4], reactions were short and not severe. We also observed unexpected side effects among our vaccination team. Vaccination team had to face with suspicious HCW whom felt reticent about it. HCW not only feared long term side effects of a 'new vaccine' [5, 6], but expressed doubt and irrational fear of long term effects.

However, taking into account the personal accomplishment associated with the importance of task, the consequence in our team was minimal (evaluate with a Malasch Burn out Inventory, adapted for vaccination).

In conclusion, all side effects of H1N1 vaccination should be anticipated including reactions to vaccine (immunological reaction and irrational behavior) and their consequences on vaccination team. We could also hope that better information based on these evaluations may decrease the fear of HCW and increase the vaccination cover.

### Acknowledgments

The authors would like to thank all members of our vaccination team, especially Dr Giroult and Ms Aouci, Combaud, Jannet, Lacheny, Tripon, Tesorière, Valin, and Olivia Laborde, an English Supervisor in our University, who helps us to improve our English.

No potential conflict of interest between authors and the present study.

### References

- [1] CDC. Use of Influenza A (H1N1) 2009 Monovalent Vaccine. Recommendations of the Advisory Committee on Immunization Practices (ACIP), 2009.
- [2] Jordan R, Hayward A. Should healthcare workers have the swine flu vaccine? BMJ 2009;339:b3398
- [3] Jefferson TO, Rivetti D, Di Pietrantonj C, Rivetti A, Demicheli V. Vaccines for preventing influenza in healthy adults. Cochrane.Database.Syst.Rev. 2007:CD001269
- [4] Bridges CB, Thompson WW, Meltzer MI, et al. Effectiveness and cost-benefit of influenza vaccination of healthy working adults: A randomized controlled trial. JAMA 2000;284:1655-63.
- [5] Pareek M, Clark T, Dillon H, Kumar R, Stephenson I. Willingness of healthcare workers to accept voluntary stockpiled H5N1 vaccine in advance of pandemic activity. Vaccine 2009;27:1242-7.
- [6] Chor JS, Ngai KL, Goggins WB, et al. Willingness of Hong Kong healthcare workers to accept pre-pandemic influenza vaccination at different WHO alert levels: two questionnaire surveys. BMJ 2009;339:b3391