

HTLV-2 in Central Africa: HTLV-2 subtype B strains similar to those found in Amerindian tribes are endemic in Bakola Pygmies from south Cameroon but not in surrounding Bantus and Baka Pygmies

Philippe Maucière, Laurent Meertens, Philippe Afonso, Sabine Plancoulaine, Claudia Filippone, Edouard Betsem, Sara Calattini, Alain Froment, Monique Van Beveren, Guy De Thé, et al.

► **To cite this version:**

Philippe Maucière, Laurent Meertens, Philippe Afonso, Sabine Plancoulaine, Claudia Filippone, et al.. HTLV-2 in Central Africa: HTLV-2 subtype B strains similar to those found in Amerindian tribes are endemic in Bakola Pygmies from south Cameroon but not in surrounding Bantus and Baka Pygmies. *Retrovirology*, BioMed Central, 2011, 8 (Suppl 1), pp.A82. <inserm-00599545>

HAL Id: inserm-00599545

<http://www.hal.inserm.fr/inserm-00599545>

Submitted on 10 Jun 2011

HAL is a multi-disciplinary open access archive for the deposit and dissemination of scientific research documents, whether they are published or not. The documents may come from teaching and research institutions in France or abroad, or from public or private research centers.

L'archive ouverte pluridisciplinaire **HAL**, est destinée au dépôt et à la diffusion de documents scientifiques de niveau recherche, publiés ou non, émanant des établissements d'enseignement et de recherche français ou étrangers, des laboratoires publics ou privés.



MEETING ABSTRACT

Open Access

HTLV-2 in Central Africa: HTLV-2 subtype B strains similar to those found in Amerindian tribes are endemic in Bakola Pygmies from south Cameroon but not in surrounding Bantus and Baka Pygmies

Philippe Mauclère^{1,2*}, Laurent Meertens¹, Philippe Afonso¹, Sabine Plancoulaine^{1,3}, Claudia Filippone¹, Edouard Betssem^{1,4}, Sara Calattini¹, Alain Froment⁵, Monique Van Beveren¹, Guy de Thé¹, Luis Quintana-Murci⁶, Renaud Mahieux¹, Antoine Gessain^{1*}

From 15th International Conference on Human Retroviruses: HTLV and Related Viruses
Leuven and Gembloux, Belgium. 5-8 June 2011

Background

Presence and origin of endemic foci of HTLV-2 infection in Africa remain a matter of debate.

Material and methods

To better appreciate the epidemiological and molecular determinants of HTLV-2 infection in Central Africa, we performed a survey in 3903 inhabitants of a South Cameroon forest area, including 1051 Bakola Pygmies, 815 Baka Pygmies and 2037 Bantus living in their neighboring. HTLV-1 and HTLV-2 infection was determined by both specific serological (IFA and WB) and molecular (different generic and specific PCR) methods.

Results

HTLV-1/2 prevalence was of 3% (117/3903) with 90 HTLV-1 (2.3%) and 27 HTLV-2 (0.7%). Surprisingly, HTLV-2 infection was restricted to Bakola Pygmies (27/1051 2.5%) with no HTLV-2 infection in any of the 2852 Baka or Bantus individuals. In Bakola Pygmies, HTLV-2 seroprevalence increased with age, reaching 6.5% in the elder persons. Ongoing intrafamilial HTLV-2 transmission was evidenced. Lymphoid T cell lines (CD8+ or CD4+, CD25+) producing HTLV-2 antigens, were established from PBMCs cultures of HTLV-2

infected individuals. Sequences of a 672 nucleotide LTR fragment, obtained from 7 HTLV-2 samples, showed a very high degree of homologies among samples (< 1% nucleotide divergence) but also surprisingly with Amerindian HTLV-2 B strains. Complete sequence (8954 bp) of one isolate confirmed a typical HTLV-2 B strain.

Conclusion

This study demonstrates clearly a HTLV-2 endemic population, with ongoing transmission, in Central Africa. Furthermore, it gives insights into several central questions regarding the origin and evolution rate of HTLV-2 and the migrations of infected populations.

Author details

¹Unité EPVO, CNRS URA 3015, Institut Pasteur, Paris, France. ²Centre Pasteur du Cameroun, Yaoundé, Cameroun. ³Laboratoire GHMI, INSERM, U.550, Paris, France. ⁴Faculté de Médecine et des Sciences Biomédicales, Université de Yaoundé, Cameroun. ⁵IRD, Musée de l'Homme, Paris, France. ⁶HEG Unit, CNRS URA 3012, Institut Pasteur, Paris, France.

Published: 6 June 2011

doi:10.1186/1742-4690-8-S1-A82

Cite this article as: Mauclère *et al.*: HTLV-2 in Central Africa: HTLV-2 subtype B strains similar to those found in Amerindian tribes are endemic in Bakola Pygmies from south Cameroon but not in surrounding Bantus and Baka Pygmies. *Retrovirology* 2011 **8**(Suppl 1):A82.

* Correspondence: agessain@pasteur.fr

¹Unité EPVO, CNRS URA 3015, Institut Pasteur, Paris, France

Full list of author information is available at the end of the article