

PD1/PD-L1 dependent immunosuppression by huiPS-derived cell population

Jonathan Pini, Arieh Moussaieff, Emilie Murriss, Reuven Shalom-Feuerstein,
Isabelle Petit, Daniel Aberdam, Matthieu Rouleau

► **To cite this version:**

Jonathan Pini, Arieh Moussaieff, Emilie Murriss, Reuven Shalom-Feuerstein, Isabelle Petit, et al..
PD1/PD-L1 dependent immunosuppression by huiPS-derived cell population. 6th european workshop
on immune-mediated inflammatory diseases, Nice, France. BioMed Central, 9 (Suppl 2), pp.P31, 2011.
<inserm-00643967>

HAL Id: inserm-00643967

<http://www.hal.inserm.fr/inserm-00643967>

Submitted on 23 Nov 2011

HAL is a multi-disciplinary open access archive for the deposit and dissemination of scientific research documents, whether they are published or not. The documents may come from teaching and research institutions in France or abroad, or from public or private research centers.

L'archive ouverte pluridisciplinaire **HAL**, est destinée au dépôt et à la diffusion de documents scientifiques de niveau recherche, publiés ou non, émanant des établissements d'enseignement et de recherche français ou étrangers, des laboratoires publics ou privés.



POSTER PRESENTATION

Open Access

PD1/PD-L1 dependent immunosuppression by huiPS-derived cell population

Jonathan Pini¹, Arieh Moussaieff¹, Emilie Murriss¹, Reuven Shalom-Feuerstein^{1,2}, Isabelle Petit^{1,2}, Daniel Aberdam^{1,2}, Matthieu Rouleau^{1*}

From 6th European Workshop on Immune-Mediated Inflammatory Diseases
Nice, France. 23-25 November 2011

Background

Human induced pluripotent stem cells (huiPS) are produced by *in vitro* reprogramming of somatic cells. They offer new perspectives in cell therapy due to their ability to differentiate in any cell types. Nevertheless, used either in allotransplantation (different donor and recipient), or in autologous transplantation (patient's derived iPS), iPS derived cells can induce rejecting immune responses [1]. It appears, therefore, crucial to better characterize their immunogenicity.

Methods

Applying to human iPS cells an *in vitro* ectodermal differentiation protocol (cellular matrix and BMP4 treatment) – initially designed in the laboratory for embryonic stem cells [2] – we isolated a cell population that can serve to analyse immunogenicity of iPS-derived cells.

Results

We demonstrate that these cells are unable to directly activate human lymphocytes, but instead display an immunosuppressive activity on allogenic activated T lymphocytes as shown by a strong inhibition of allogenic CD4- and CD8-positive T cell proliferation and reduction in IL-2 and IFN-gamma production. While immunosuppressive soluble factors such as IL-10 or TGF-beta are not involved, we demonstrate, with the use of blocking antibodies, that this immunosuppressive activity rely on the inhibitory interaction between the PD1 receptor (upregulated on T cells) and its ligand PD-1L strongly expressed by our human iPS –derived population.

Finally, we demonstrate that this cell population contains mesenchymal stem cells, as shown by immunophenotyping,

cytokines production and *in vitro* differentiation into adipocytes, osteoblasts and chondrocytes.

Conclusions

These results suggest that during *in vitro* human iPS cell differentiation into defined cell populations with strong interest in cellular therapy, some mesenchymal stem cells with immunosuppressive activity may differentiate and limit the immune process suppose to induce their rejection. Interestingly, the mechanisms involved might *in fine* participate in establishing a long term immune tolerance.

Acknowledgements

Financial support was provided by AFM (#14272), INSERM and University of Nice Sophia Antipolis.

Author details

¹U898 INSERM, University of Nice Sophia Antipolis, Nice, France. ²INSERTECH, Bruce Rappaport Institute of the Technion, Haifa, Israel.

Published: 23 November 2011

References

1. Zhao T, Zhang ZN, Rong Z, Xu Y: Immunogenicity of induced pluripotent stem cells. *Nature* 2011, **474**:212-215.
2. Aberdam E, Barak E, Rouleau M, Laforest SD, Berrih-Aknin S, Suter DM, Krause KH, Amit M, Itskovitz-Eldor J, Aberdam D: A Pure Population of Ectodermal Cells Derived from Human Embryonic Stem Cells. *Stem Cells* 2008, **26**:440-444.

doi:10.1186/1479-5876-9-S2-P31

Cite this article as: Pini et al.: PD1/PD-L1 dependent immunosuppression by huiPS-derived cell population. *Journal of Translational Medicine* 2011 **9**(Suppl 2):P31.

¹U898 INSERM, University of Nice Sophia Antipolis, Nice, France
Full list of author information is available at the end of the article