



**HAL**  
open science

## New insights into microtubule elongation mechanisms.

Odile Valiron

► **To cite this version:**

Odile Valiron. New insights into microtubule elongation mechanisms.. Communicative and Integrative Biology, 2011, 4 (1), pp.10-3. 10.4161/cib.4.1.13581 . inserm-00644013

**HAL Id: inserm-00644013**

**<https://inserm.hal.science/inserm-00644013>**

Submitted on 23 Nov 2011

**HAL** is a multi-disciplinary open access archive for the deposit and dissemination of scientific research documents, whether they are published or not. The documents may come from teaching and research institutions in France or abroad, or from public or private research centers.

L'archive ouverte pluridisciplinaire **HAL**, est destinée au dépôt et à la diffusion de documents scientifiques de niveau recherche, publiés ou non, émanant des établissements d'enseignement et de recherche français ou étrangers, des laboratoires publics ou privés.

## **New insights into microtubule elongation mechanisms**

**Odile VALIRON**

Institut National de la Santé et de la Recherche Médicale Unité 836, Institut des Neurosciences de Grenoble ; Université Joseph Fourier, 38042 Grenoble, Cedex 9, France

**Address correspondence to:** Odile VALIRON, Grenoble Institut des Neurosciences, Bâtiment Edmond J. Safra, Université Joseph Fourier, Site Santé La Tronche, BP 170, 38042 Grenoble Cedex 9, France. Tel: (+33) 4 56 52 05 37. Fax: (+33) 4 56 52 06 57. E-mail: [odile.valiron@ujf-grenoble.fr](mailto:odile.valiron@ujf-grenoble.fr)

**Key words:** microtubules, tubulin, nucleotide, GTP hydrolysis, tubulin structure, microtubule assembly, microtubule dynamics.

**Acknowledgements:** I thank Dr I. Arnal and Dr. D. Job for critical reading of the manuscript, and Dr. I. Arnal for figure 1A, B.

**Abbreviations:** GTP-tub, tubulin associated to GTP on exchangeable site; GDP-tub, tubulin associated to GDP on exchangeable site; GMPCPP, guanylyl-( $\alpha,\beta$ )-methylene-diphosphonate; GMPCPP-microtubules, microtubules assembled in the presence of GMPCPP; GDP-microtubules, microtubules assembled in the presence of GDP, followed by GTP hydrolysis in the polymer wall.

**Abstract:** Microtubules are cytoskeletal structures in the cytoplasm of eukaryotic cells, and their highly dynamic properties are essential to perform a wide variety of vital functions in cells. Microtubule growth proceeds through the endwise addition of nucleotide-bound tubulin molecules. It has largely been assumed that only tubulin dimers can incorporate into microtubules, and that the chemical state of the nucleotide is crucial for the incorporation. Recent observations reveal that both tubulin dimers and oligomers can add to microtubule ends and that the chemical state of the nucleotide is not decisive for tubulin addition. Together with structural studies of tubulin, these results show

tubulin assembly polymorphism, which could play a crucial role in microtubule-dependent cellular functions.

**Running title:** New insights into microtubule elongation mechanism

## Text

### Introduction

Microtubules are key components of the eukaryotic cell cytoskeleton and they have important roles in many cellular processes such as intracellular transport, cell motility, cell morphogenesis, meiosis and mitosis. The basic structural unit of microtubules is the  $\alpha\beta$ -tubulin heterodimer. Tubulin heterodimers bind head to tail to form protofilaments. In the general case, 13 protofilaments associate to give rise to a cylindrical microtubule, in which each tubulin molecule has lateral and longitudinal interactions with its neighbours in the polymer wall. Microtubules can switch between growing and shortening phases, a phenomenon known as dynamic instability.<sup>1,2</sup> During the growing phase, tubulin molecules are added at the end of polymers. Microtubules shorten through tubulin loss from their tips. In cells, dynamic instability is highly regulated by microtubule associated proteins (for reviews see references 3-4). However, this dynamic behaviour is an intrinsic property of microtubules themselves, observable also in microtubules assembled from purified tubulin suspensions. Actually, our knowledge of how microtubules grow and shrink is principally based on in vitro studies performed with pure tubulin (Figure 1A, B).

$\alpha$ - and  $\beta$ -tubulin monomers carry nucleotide binding sites in their N-terminal domain. The  $\alpha$ -subunit is constitutively associated to a non-exchangeable GTP, buried at the monomer-monomer interface. In contrast, the  $\beta$ -tubulin subunit comprises a binding site exposed at the protein surface and at which the nucleotide is exchangeable (E-site). Following the addition at microtubule end of a tubulin dimer containing GTP on E-site (GTP-tub), the E-site GTP is hydrolyzed and the newly generated GDP becomes non-exchangeable. As a result the microtubule wall is mainly composed of tubulin subunits containing GDP on E-site (GDP-tub).

A recent study has shown that, irrespective of their nucleotide binding state (GTP, GDP or a slowly hydrolysable analogue of GTP, GMPCPP), the majority of tubulin dimers in solution form small oligomers.<sup>5</sup> Moreover, GDP-tub oligomers are released from shortening microtubules<sup>6,7</sup> (Figure 1B). Thus, a simple chemical system containing only tubulin and GTP in appropriate conditions of buffer and temperature will ultimately be composed of multiple forms of tubulin molecules, GTP-tub

and GDP-tub dimers and oligomers, and microtubules. This review focuses on how these various tubulin subunits and polymer intermediates contribute to, and modulate, microtubule growth and dynamics.

### **Do tubulin oligomers participate to microtubule elongation?**

In recent experiments, individual microtubule elongation has been tracked at molecular scale. Two sets of experiments using optical tweezers showed that the microtubule length increases by step during polymer elongation.<sup>7,8</sup> However length increment measurements were not similar in both studies and these observations led to different conclusions. One report shows that small tubulin oligomers (20-30 nm i.e. 2 to 4 associated tubulin dimers) are able to add directly to microtubule tips<sup>8</sup> whereas the other study indicates that only tubulin dimers (8 nm) are incorporated.<sup>7</sup> Electron microscopy experiments suggest also that microtubules can elongate through tubulin oligomers addition.<sup>5</sup>

Previous electron microscopy studies have shown that growing microtubule ends form intermediate sheet structures that subsequently close into tubes<sup>9</sup> (Figure 1A). An implication of this finding is that tubulin molecules may add either at protofilament tips or within sheets by lateral interactions with tubulin subunits forming the sheet edge. Tubulin addition at the end of protofilament directly increases microtubule length and is detected by optical microscopy measurements. In contrast, the incorporation of tubulin molecules in the sheet does not participate to immediate microtubule length increase<sup>5</sup> and is not revealed using optical microscopy. The incorporation of tubulin as dimers or oligomers is therefore still a matter of debate.

### **GTP-tubulin remnants in microtubules**

Microtubules are able to polymerize in the presence of GMPCPP (GMPCPP-microtubules).<sup>10</sup> Structural differences have been detected in the lattice of GMPCPP-microtubules and of microtubules assembled from GTP, thereby composed of GDP-tub (GDP-microtubules).<sup>11</sup> Recently, an antibody was selected that recognizes specifically GMPCPP-microtubules but not GDP-microtubules.<sup>12</sup> In cells, the antibody stains growing microtubule ends, and also short fragments in the middle of microtubules.

The authors suggest that GTP-tubulin remnants could be present in microtubule lattice, indicating that GTP hydrolysis is sometimes incomplete during polymerization.<sup>12</sup> Interestingly, the position of these internal sites seems to correlate in cells with transitions from microtubule shrinkage to growth. GTP-remnants would then have a structural state more stable than the whole microtubule lattice. This new state remains to be structurally characterized. This study shows that GTP-tub can also be included in microtubules without GTP hydrolysis step, providing intrinsic chemical variability to microtubule lattice.

### **GDP-tub is directly incorporated in microtubules**

The possibility of a direct incorporation of GDP-tub into growing microtubules has been raised at early stages of microtubule research, and has remained a matter of controversy. Essentially based on turbidimetry measurements, an increase of polymerized tubulin was reported in suspension containing tubulin and GTP in the presence of GDP-tub.<sup>13-15</sup> These results suggested that GDP-tub was directly assembled into microtubules, although GDP-tub incorporation was not directly measured. Alternatively, other studies demonstrated that the addition of GDP-tub to microtubule suspension does not participate significantly to microtubule elongation.<sup>16-19</sup> The effect of GDP-tub on microtubule dynamics has been studied and results also exhibited discrepancies. Reports have shown that GDP-tub stabilizes microtubules<sup>17-18</sup> while in another study the principal effect of GDP-tub is an increase of catastrophe and rescue frequencies, with the occurrence of growth and shortening irregularities.<sup>20</sup>

Recently, the question of direct GDP-tub incorporation in microtubules was re-addressed using a minimal tubulin assembly system composed of nucleotide bound tubulin dimers, in the absence of excess free nucleotide.<sup>13, 21-24</sup> Such a system has two main advantages: (1) the proportion of GTP-tub and GDP-tub added in solution can be controlled at will and (2) measurements of direct incorporation of both nucleotide-bound tubulin dimers can be performed.<sup>24</sup> Results demonstrate that substantial amounts of GDP-tub can be directly incorporated into growing microtubules. Moreover microtubules assembled from GTP-tub and GDP-tub mixtures display modified dynamic behavior. Video-microscopy experiments performed with microtubules nucleated on centrosomes demonstrated that growth and shrinkage rates decreased by half in the presence of 30% GDP-tub in the starting mix.

The decrease of the shrinkage rate was of particular interest because microtubule shrinkage seems to arise from a zero order reaction, governed by intrinsic factors reflecting the structural state of the microtubule lattice.

Previous works have established that GTP hydrolysis is not necessary for tubulin assembly, but rather for microtubule disassembly.<sup>10, 25</sup> When tubulin is assembled in the presence of GMPCPP, resulting GMPCPP-microtubules are stable compared with GDP-microtubules.<sup>10</sup> This strongly suggests that much of the energy released by GTP hydrolysis is stored in the microtubule lattice and released during microtubule disassembly. The direct incorporation of GDP-tub into microtubule lattice is obviously not followed by hydrolysis of the bound nucleotide. Therefore it could be suggested that GDP may function as a natural non-hydrolyzable analog of GTP, with resulting impairment of the disassembly properties of microtubules. Detectable modifications have been found in the lattice organization of GMPCPP-microtubules compared to GDP-microtubules.<sup>11</sup> No structural modification was observed in microtubules that incorporated directly GDP-tub. It is possible that lattice modification would be blurred by the co-incorporation of GTP-tub and GDP-tub, compared to the homogeneity of GMPCPP-microtubules.

### **Structural changes in tubulin**

Structural studies of tubulin heterodimer in various nucleotide and polymerization states have been reported in the past few years. They have led to two challenging models for the relationship between GTP and conformational changes of tubulin molecule. A first set of reports indicate that GTP-tub has a straight conformation,  $\alpha$  and  $\beta$  tubulin subunits being aligned, while GDP-tub adopts a more bent conformation, with a kink between both tubulin subunits.<sup>26-27</sup> These results lead to a model in which curved GDP-tub would then be unable to form the lateral contacts necessary to fit the microtubule lattice, therefore would be unable to participate to microtubule elongation. Following nucleotide exchange, GTP-tub would straighten and lateral contacts between adjacent tubulin molecules could become possible.<sup>26-27</sup> More recently, another study suggests that free GTP-tub and GDP-tub in solution may be similarly bent. Major structural rearrangements of tubulin would not occur in response to GTP binding. Tubulin would straighten only after incorporation into growing

microtubules, as a consequence of the straightening of protofilaments during tube closure.<sup>28</sup> GTP only tunes the strength of contacts in the microtubule lattice. This model is compatible with a direct incorporation of GDP-tub within microtubule lattice, together with GTP-tub. Actually GDP-tub alone is not able to assemble into microtubules.<sup>13, 19, 24</sup>

These models only consider the addition of tubulin dimers into microtubule lattice. Results reported above suggest that tubulin oligomers are also able to add either into terminal sheet at microtubule tip or directly at the end of protofilaments. Within these oligomers, it seems likely that a proportion of tubulin-bound GTP is hydrolyzed so that oligomers are composed of both GTP-tub and GDP-tub. Such addition of macromolecular assembly of tubulin dimers remains to be structurally investigated.

## **Conclusions**

Multiple macromolecular tubulin assemblies can be formed from purified tubulin molecules, depending on nucleotide bound and oligomerization states. Following the classical textbook model, microtubules elongate exclusively by addition of GTP-tub. In contrast with this model, it has been shown that tubulin dimers as well as these structural intermediates may be used in microtubule elongation that therefore shows a high degree of plasticity. The recent reports of GDP-tub incorporation and GTP-tub remnants suggest that the tubulin subunits constituting the microtubule wall are in different structural states according to their initial nucleotide bound state, with resulting variations in intrinsic microtubule dynamic properties (Figure 1C). These findings reveal a novel form of microtubule “structural plasticity”.<sup>29</sup> Moreover, the polymorphism of microtubule assembly could reflect mechanisms that tune the interactions between microtubules and cellular factors.<sup>30</sup> To go further, we will need probes that recognize different structural states of microtubules in cells. An antibody is already available that recognizes specific structural states.<sup>12</sup> May be, other binding factors, either antibodies or drugs or proteins, will prove to be useful reporters of the microtubule structural status, which could facilitate in vivo investigation of microtubule structural plasticity.

## References

1. Desai A, Mitchison TJ. Microtubule polymerization dynamics. *Annu Rev Cell Dev Biol* 1997; 13:83-117.
2. Valiron O, Caudron N, Job D. Microtubule dynamics. *Cell Mol Life Sci* 2001; 58:2069-2084.
3. Amos LA, Schlieper D. Microtubules and maps. *Adv Protein Chem* 2005; 71:257-298.
4. Akhmanova A, Steinmetz MO. Tracking the ends: a dynamic protein network controls the fate of microtubule tips. *Nat Rev Mol Cell Biol* 2008; 9:309-322.
5. Mozziconacci J, Sandblad L, Wachsmuth M, Brunner D, Karsenti E. Tubulin dimers oligomerize before their incorporation into microtubules. *PLoS ONE* 2008; 3(11):e3821. doi:10.1371/journal.pone.0003821
6. Mandelkow EM, Mandelkow E, Milligan RA. Microtubule dynamics and microtubule caps: a time-resolved cryo-electron microscopy study. *J Cell Biol* 1991; 114:977-991.
7. Schek HT, Gardner MK, Cheng J, Odde DJ, Hunt AJ. Microtubule assembly dynamics at the nanoscale. *Curr Biol* 2007; 17:1445-1455.
8. Kerssemakers JW, Munteanu EL, Laan L, Noetzel TL, Janson ME, Dogterom M. Assembly dynamics of microtubules at molecular resolution. *Nature* 2006; 442:709-712.
9. Chrétien D, Fuller SD, Karsenti E. Structure of growing microtubule ends: two-dimensional sheets close into tubes at variable rates. *J Cell Biol* 1995; 129:1311-1328.
10. Hyman AA, Salser S, Drechsel DN, Unwin N, Mitchison TJ. Role of GTP hydrolysis in microtubule dynamics: information from a slowly hydrolyzable analogue, GMPCPP. *Mol Biol Cell* 1992; 3:1155-1167.
11. Hyman AA, Chrétien D, Arnal I, Wade RH. Structural changes accompanying GTP hydrolysis in microtubules: information from a slowly hydrolyzable analogue guanylyl-( $\alpha,\beta$ )-methylene-diphosphonate. *J. Cell Biol* 1995; 128:117-125.
12. Dimitrov A, Quesnoit M, Moutel S, Cantaloube I, Poüs C, Perez F. Detection of GTP-tubulin conformation in vivo reveals a role for GTP remnants in microtubules rescues. *Science* 2008; 322:1353-1356.

13. Carlier MF, Pantaloni D. Kinetic analysis of cooperativity in tubulin polymerization in the presence of guanosine di- or triphosphate nucleotides. *Biochemistry* 1978; 17:1908-1915.
14. Hamel E, Batra JK, Lin CM. Direct incorporation of guanine 5'-diphosphate into microtubules without guanosine 5'-triphosphate hydrolysis. *Biochemistry* 1986; 25:7054-7062.
15. Lin CL, Hamel E. Interrelationships of tubulin-GDP and tubulin-GTP in microtubule assembly. *Biochemistry* 1987; 26:7173-7182.
16. Jameson L, Caplow M. Effect of guanosine diphosphate on microtubule assembly and stability. *J Biol Chem* 1980; 255:2284-2292.
17. Zackroff RV, Weisenberg RC, Deery WJ. Equilibrium and kinetic analysis of microtubule assembly in the presence of guanosine diphosphate. *J Mol Biol* 1980; 139:641-677.
18. Caplow M, Reid R. Direct elongation model for microtubule GTP hydrolysis. *Proc Natl Acad Sci USA* 1985; 82:3267-3271.
19. Bayley PM, Butler FMM, Manser EJ. Control of nucleation in microtubule self-assembly. *FEBS Lett* 1986; 205:230-234.
20. Vandecandelaere A, Martin SR, Bayley PM. Regulation of microtubule dynamic instability by tubulin-GDP. *Biochemistry* 1995; 34:1332-1343.
21. O'Brien ET, Erickson HP. Assembly of pure tubulin in the absence of free GTP: effect of magnesium, glycerol, ATP, and the nonhydrolyzable GTP analogues. *Biochemistry* 1989; 28:1413-1422.
22. Carlier MF, Didry D, Pantaloni D. Hydrolysis of GTP associated with the formation of tubulin oligomers is involved in microtubule nucleation. *Biophys J* 1997; 73:418-427.
23. Caudron N, Valiron O, Usson Y, Valiron P, Job D. A reassessment of the factors affecting microtubule assembly and disassembly in vitro. *J Mol Biol* 2000; 297:211-220.
24. Valiron O, Arnal I, Caudron N, Job D. GDP-tubulin incorporation into growing microtubules modulates polymer stability. *J Biol Chem* 2010; 285:17507-17513.
25. Caplow M, Ruhlen RL, Shanks J. The free energy for hydrolysis of a microtubule-bound nucleotide triphosphate is near zero: all the free energy for hydrolysis is stored in the microtubule lattice. *J Cell Biol* 1994; 127:779-788.

26. Müller-Reichert T, Chrétien D, Severin F, Hyman AA. Structural changes at microtubule ends accompanying GTP hydrolysis: information from a slowly hydrolyzable analogue of GTP, guanylyl-(alpha, beta)methylenediphosphonate. *Proc Natl Acad Sci USA* 1998; 95:3661-3666.
27. Wang HW, Nogales E. Nucleotide-dependent bending flexibility of tubulin regulates microtubule assembly. *Nature* 2005; 435:911-915.
28. Rice LM, Montabana EA, Agard DA. The lattice as allosteric effector: structural studies of  $\alpha\beta$ - and  $\gamma$ -tubulin clarify the role of GTP in microtubule assembly. *Proc Natl Acad Sci USA* 2008; 105:5378-5383.
29. Kueh HY, Mitchison TJ. Structural plasticity in actin and tubulin polymer dynamics. *Science* 2009; 325:960-963.
30. Nogales E, Wang HW. Structural mechanisms underlying nucleotide-dependent self-assembly of tubulin and its relative. *Curr Opin Struct Biol* 2006; 16:1-9.

## Figure Legends

**Figure 1: Microtubule assembly.** (A) Cryo-electron microscopy image of a growing microtubule end showing an opened sheet (arrow). (B) Cryo-electron microscopy image a shrinking microtubule end showing protofilaments peeling into ring-like structures (arrows). Numerous tubulin oligomers coming from microtubule disassembly are visible in the background (arrowheads). Scale bar, 50 nm in both panels. (C) Schematic representation of a growing microtubule. The end displays a sheet-like conformation which later close into a tube. GTP-tub dimers, GDP-dimers and tubulin oligomers participate to microtubule elongation at the end of the polymer. GTP-tub,  $\alpha$ -tubulin and  $\beta$ -tubulin associated to GTP; GDP-tub,  $\alpha$ -tubulin and  $\beta$ -tubulin associated to GDP; \* indicate GTP-remnants integrated in the microtubule wall. Cryo-electron microscopy images are courtesy of Dr. I. Arnal, UMR 6026, Rennes, France.

Figure 1

