

**Unusual presentation of hepatitis B serological markers
in an Amerindian community of Venezuela with a
majority of occult cases.**

Nathalia Cardona, Carmen Loureiro, Domingo Garzaro, María Duarte, Daisy
García, Milian Pacheco, Isabelle Chemin, Flor Pujol

► **To cite this version:**

Nathalia Cardona, Carmen Loureiro, Domingo Garzaro, María Duarte, Daisy García, et al.. Unusual presentation of hepatitis B serological markers in an Amerindian community of Venezuela with a majority of occult cases.. *Virology Journal*, BioMed Central, 2011, 8 (1), pp.527. <10.1186/1743-422X-8-527>. <inserm-00657506>

HAL Id: inserm-00657506

<http://www.hal.inserm.fr/inserm-00657506>

Submitted on 6 Jan 2012

HAL is a multi-disciplinary open access archive for the deposit and dissemination of scientific research documents, whether they are published or not. The documents may come from teaching and research institutions in France or abroad, or from public or private research centers.

L'archive ouverte pluridisciplinaire **HAL**, est destinée au dépôt et à la diffusion de documents scientifiques de niveau recherche, publiés ou non, émanant des établissements d'enseignement et de recherche français ou étrangers, des laboratoires publics ou privés.

RESEARCH

Open Access

Unusual presentation of hepatitis B serological markers in an Amerindian community of Venezuela with a majority of occult cases

Nathalia E Cardona¹, Carmen L Loureiro², Domingo J Garzaro², María C Duarte¹, Daisy M García¹, Milian C Pacheco¹, Isabelle Chemin³ and Flor H Pujol^{2*}

Abstract

Background: Occult hepatitis B infection (OBI) is characterized by the presence of hepatitis B virus (HBV) DNA in the absence of HBsAg in the serum of patients. The aim of this study was to characterize HBV infection among a Piaroa community, an Amerindian group which exhibits significant evidence of exposure to HBV but relatively low presence of HBsAg, and to explore the presence of OBI in this population.

Results: Of 150 sera, with 17% anti-HBc and 1.3% HBsAg prevalence, 70 were tested for the presence of HBV DNA. From these, 25 (36%) were found positive for HBV DNA by PCR in the core region. Two of these 25 sera were HBsAg positive, indicating an overt infection. Of the remaining 68 sera tested, 23 exhibited OBI. Of these, 13 were HBV DNA out of 25 anti-HBc positive (52%) and 10 HBV DNA positive, out of 43 anti-HBc negative (23%), with a statistical significance of $p = 0.03$. Viral DNA and HBsAg were present intermittently in follow up sera of 13 individuals. Sequence analysis in the core region of the amplified DNA products showed that all the strains belonged to HBV genotype F3. The OBI isolates displayed 96-100% nucleotide identity between them. One isolate exhibited the co-circulation of a wild type variant with a variant with a premature stop codon at the core protein, and a variant exhibiting a deletion of 28 amino acids.

Conclusions: The frequency of OBI found in this Amerindian group warrants further studies in other communities exhibiting different degrees of HBV exposure.

Keywords: Hepatitis B virus, Occult infection, Amerindians

Background

Hepatitis B virus (HBV) infection is a significant health concern among Amerindians in the Americas with high exposure being documented in several Amerindian groups [1]. However, the prevalence of active HBV infection, defined as positivity for HBV surface antigen (HBsAg) is variable among different Amerindian communities, coexisting in the same geographic environment [2]. In a recent study in the Venezuelan Amazon, anti-HBc prevalence ranged from 17 to 70% [2].


Occult hepatitis B virus infection (OBI) is characterized by the presence of hepatitis B virus (HBV) DNA in

the absence of HBV surface antigen (HBsAg) [3,4]. OBI can lead to severe chronic manifestations including hepatocellular carcinoma (HCC) [5,6]. OBI has not been studied thoroughly in Amerindian populations and could be present in Amerindian populations exhibiting evidence of exposure to HBV without high prevalence of active infection. Indeed, OBI has been already described in Mexican Amerindians [7]. The aim of this study was to characterize HBV infection among a Piaroa community, an Amerindian group which exhibits significant evidence of exposure to HBV but relatively low presence of HBsAg [2], and to explore the presence of OBI in this population.

* Correspondence: fhpujol@gmail.com

²Laboratorio de Virología Molecular, Centro de Microbiología y Biología Celular, Instituto Venezolano de Investigaciones Científicas, Apdo 20632, Caracas 1020-A, Venezuela

Full list of author information is available at the end of the article



Results

A total of 150 sera from the Piaroa community Babilla de Pintao were analyzed (Figure 1). Total anticore antibodies (anti-HBc) prevalence was 17% (26/150) in this group and 31% (25/80) in individuals over 15 years of

age [2]. Only 2 sera (1.3%) were positive for HBsAg [2]. These 2 sera were negative for anti-HBc antibodies. A subset of 70 sera was analyzed for the presence of HBV DNA. Of these, 25 (36%) were positive for HBV DNA by PCR in the core region (Figure 1). All individuals

showed normal ALT levels. The 2 HBsAg sera were positive for HBV DNA. Of the remaining 23 sera, 13 were anti-HBc positive, and 10 were both anti-HBc and HBsAg negative. Among the HBsAg negative sera, 52% of the anti-HBc positive and 23% of the anti-HBc negative sera were HBV DNA positive, this difference being statistically significant ($p = 0.03$). HBV DNA was found even more frequently among anti-HBs positive individuals compared to anti-HBs negative ones ($p = 0.01$) (Figure 1). No difference was observed in the prevalence of OBI according to sex (9/25 of females and 16/41 of males had HBV DNA in their sera, $p = 0.99$), or to age (9/30 younger than 30 years vs. 12/25 older, $p = 0.26$).

Follow up sera were available for 13 individuals positive for HBV DNA. Viral DNA and HBsAg were present intermittently, as shown in Table 1. The two individuals presenting with an overt HBV infection at the beginning of the study, developed OBI later, since they carried HBV DNA in their sera for more than 2 years without the presence of HBsAg. The HBV genomic region that could readily be amplified was the core region, while the


S region could be amplified only in some sera (Table 1). From the sera collected from vaccinated subjects in 2009, 34/36 showed levels of anti-HBs antibodies higher than 10 mIU/ml.

Sequence analysis in the core region of the DNA amplified products showed that all the strains belonged to HBV genotype F3 (Figure 2). The OBI isolates displayed 96-100% nucleotide identity between them. The isolates were also closely related to sequences from HBV isolates circulating among other Piaroa, Yanomami and Yucpa Amerindians exhibiting overt infections and analyzed in previous studies [8]. One isolate, BP21, exhibited co-circulation of a wild type virus along with a variant harboring a premature stop codon at aa 42 of the core protein, and a variant exhibiting a deletion of 28 aas (aa 78-105) (Figure 3). A partial S genomic sequence was also available for 8 specimens. The sequences in the S region indicate the presence of HBV genotype F3, subtype adw4, although the length of the genomic region analyzed did not permit firm subgenotyping. Mutations associated with escape from antibody

Table 1 HBV DNA in sera from Piarao Amerindians

Serum	Collected April 2002			Collected March 2003			Collected August 2004		
	Serological status ¹	Core ² DNA	S ² DNA	Serological status ¹	Core ² DNA	S ² DNA	Serological status ¹	Core ² DNA	S ² DNA
BP131	S +, AC -	+	-	S -, AC-	+	+	S +, AC -	+	-
BP132	S +, AC	+	-	S -, AC-	+	-	S -, AC+	+	-
BP11	S -, AC+	+							
BP14	S -, AC+	+							
BP19	S -, AC+	+					S -, AC+	+	+
BP29	S -, AC+	+	+						
BP31	S -, AC+	+							
BP43	S -, AC+	+							
BP88	S -, AC+	+	-						
BP89	S -, AC+	+							
BP97	S -, AC+	+	+	S -, AC+	+				
BP113	S -, AC+	+							
BP117	S -, AC+	+							
BP152	S -, AC+	+							
BP168	S -, AC+	+							
BP21	S -, AC-	+	-				S -, AC-	+	+
BP74	S -, AC-	+	-				S -, AC-	+	-
BP92	S -, AC-	+	-				S -, AC-	+	-
BP134	S -, AC-	+	-	S -, AC-	+				
BP136	S -, AC-	+	-	S -, AC-	+	-			
BP147	S -, AC-	+	+	S+, AC+	+	+	S -, AC+	+	-
BP150	S -, AC-	+	-						
BP154	S -, AC-	+	-	S -, AC-	+	-	S +, AC-	+	-
BP156	S -, AC-	+	-	S -, AC-	+	+	S -, AC-	+	-
BP164	S -, AC-	+	+	S +, AC-	+		S -, AC-	+	-

1: Serological status: S (HBsAg), AC (anti-HBc). 2: PCR of the core (C) or surface antigen (S) region. Blank cells mean not determined




neutralization were not observed (data not shown). All the OBI strains were genetically related (Figure 2). Interestingly, 19/25 specimens of OBI shared at least one parent exhibiting OBI.

Discussion

HBV DNA was analyzed in an Amerindian population exhibiting moderate prevalence of infection (17% anti-

HBc), compared to other Venezuelan Amerindian populations, such as that of the Yanomami (58% anti-HBc). As described previously, this community showed a lower rate of acquisition of anti-HBc antibodies (1.4% in individuals less than 15 years old), compared to the Yanomami for example (38% in individuals less than 15 years old) [2]. The lower prevalence of HBV exposure and infection in this Amerindian community may be



genotype A was present at a similar prevalence in overt and OBI infections [18]. Altogether, these studies suggest that OBI appears not to be restricted to a particular genotype. In our study, one subject was infected by a wild type virus with variants coding for core defective proteins, a situation already described in Venezuelan blood donors with OBI [9]. Most of the subjects with OBI were related, suggesting that familiar transmission might have played a role in this situation. However, the number of samples analyzed and the short genomic sequence available for study did not allow testing of this hypothesis.

There is accumulating evidence of a pathogenic role for OBI [19]. OBI may contribute to the progression of liver fibrosis and HCC development [20], thus the potential benefits of antiviral treatment is in debate [6,21]. As shown in this study and in others, vaccination of those populations at risk for OBI should be undertaken as it may bring some benefits to these communities [22].

Conclusions

A high frequency of unusual HBV presentation was found in this Piaroa population. All the individuals were infected with HBV genotype F3. The OBI isolates displayed a restrained variability and were similar to the isolates causing overt HBV infection in other Venezuelan Amerindian groups. The frequency of OBI found in this Piaroa population warrants further studies in other Amerindian communities exhibiting different degrees of HBV exposure.

Methods

Population group

The Piaroa community of Babilla de Pintao (Amazon State, Venezuela) consists of 169 inhabitants, and 150 sera were analyzed for the presence of HBV serological markers, with informed consent and under approval of Bioethical Committees of CAICET and IVIC [2]. Individuals were also vaccinated during this period. Testing was performed between 2002 and 2004, and a subset of sera ($n = 36$) were collected in 2009 to evaluate anti-HBs antibodies.

Serological assays

Sera were tested for HBV markers with commercial assays: HBcAb DIMA™ (DIMA Diagnostika C.A., Venezuela), Murex HBsAg Version 3 (ABBOTT, Murex Biotech Limited, UK), Bioelisa anti-HBs (Biokit, S.A., Spain) and IgM anti-HBc by ETI-CORE-IGMK-2 (DiaSorin Ltda., Italy). A sample was considered anti-HBs positive if the levels of anti-HBs antibodies were higher than 10 UI/ml. ALT were also determined with a commercial assay (Wiener Lab, Argentina).

PCR and sequencing

A total of 70 sera (2 HBsAg positive, 25 anti-HBc positive and 43 remaining randomly selected) were analyzed by nested PCR of the core region [9]. A sample was considered positive if it repeated positive after a second extraction of viral DNA. When enough serum was available, samples were also amplified by nested PCR in the S region [8]. Purified PCR fragments were sent to CESAAN (Centro de Secuenciación y Análisis de Ácidos Nucleicos, IVIC, Caracas, Venezuela), for sequencing. Sequences obtained from the Venezuelan isolates were compared with different reference strains from GenBank and used for phylogenetic analysis. Sequence alignment and phylogenetic analysis by the Neighbor Joining method (1,000 bootstrap replicas, genetic distances evaluated with Kimura 2 parameters corrections) were conducted using DNAMAN 5.2.2 (Lynnon Bio Soft, Canada). Nucleotide sequence data have been deposited in GenBank database under the accession numbers JN255220-JN255243.

Statistical analysis

Statistical differences were evaluated by the Chi-Squares test with Yates correction, according to a computerized Epi Info program, version 3.3.2 (Centers for Disease Control and Prevention, Atlanta, GA).

List of abbreviations

HBV: Hepatitis B virus; OBI: Occult hepatitis B virus infection; HCC: Hepatocellular carcinoma; HBsAg: HBV surface antigen; anti-HBc: Anticore antibodies

Acknowledgements

This work was supported by Grant G-200001493 from FONACIT, Venezuela, and Projet ECOS-Nord France-Venezuela: V09S02.

Author details

¹Servicio Autónomo: Centro Amazónico para la Investigación y Control de enfermedades Tropicales, Simón Bolívar "CAICET", Puerto Ayacucho, Venezuela. ²Laboratorio de Virología Molecular, Centro de Microbiología y Biología Celular, Instituto Venezolano de Investigaciones Científicas, Apdo 20632, Caracas 1020-A, Venezuela. ³INSERM U1052 CRCL, Lyon I University, Villeurbanne, France.

Authors' contributions

NEC, CLL and DJG carried out the molecular genetic studies, and participated in the sequence alignment. NEC, IC and FHP drafted the manuscript. DMG and MCP carried out the immunoassays. MCD participated in the clinical and epidemiological study. NEC and FHP participated in the design of the study and performed the statistical analysis. All authors read and approved the final manuscript.

Competing interests

The authors declare that they have no competing interests.

Received: 2 September 2011 Accepted: 9 December 2011
Published: 9 December 2011

References

1. Devesa M, Pujol FH: Hepatitis B virus genetic diversity in Latin America. *Virus Res* 2007, **127**:177-184.

2. Duarte MC, Cardona N, Poblete F, González K, García M, Pacheco M, Botto C, Pujol FH, Williams JR: **A comparative epidemiological study of hepatitis B and hepatitis D virus infections in Yanomami and Piara Amerindians of Amazonas State, Venezuela.** *Trop Med Int Health* 2010, **15**:924-933.
3. Said ZN: **An overview of occult hepatitis B virus infection.** *World J Gastroenterol* 2011, **17**:1927-1938.
4. Hollinger FB, Sood G: **Occult hepatitis B virus infection: a covert operation.** *J Viral Hepat* 2010, **17**:1-15.
5. Lledó JL, Fernández C, Gutiérrez ML, Ocaña S: **Management of occult hepatitis B virus infection: an update for the clinician.** *World J Gastroenterol* 2011, **17**:1563-1568.
6. Owiredu WK, Kramvis A, Kew MC: **Hepatitis B virus DNA in serum of healthy black African adults positive for hepatitis B surface antibody alone: possible association with recombination between genotypes A and D.** *J Med Virol* 2001, **64**:441-454.
7. Roman S, Tanaka Y, Khan A, Kurbanov F, Kato H, Mizokami M, Panduro A: **Occult hepatitis B in the genotype H-infected Nahuas and Huichol native Mexican population.** *J Med Virol* 2010, **82**:1527-1536.
8. Devesa M, Loureiro CL, Rivas Y, Monsalve F, Cardona N, Duarte MC, Poblete F, Gutierrez MF, Botto C, Pujol FH: **Subgenotype diversity of hepatitis B virus American genotype F in Amerindians from Venezuela and the general population of Colombia.** *J Med Virol* 2008, **80**:20-26.
9. Gutiérrez C, Devesa M, Loureiro CL, León G, Liprandi F, Pujol FH: **Molecular and serological evaluation of surface antigen negative hepatitis B virus infection in blood donors from Venezuela.** *J Med Virol* 2004, **73**:200-207.
10. Minuk GY, Sun DF, Uhanova J, Zhang M, Caouette S, Nicolle LE, Gutkin A, Doucette K, Martin B, Giulivi A: **Occult hepatitis B virus infection in a North American community-based population.** *J Hepatol* 2005, **42**:480-485.
11. Martin CM, Welge JA, Shire NJ, Rouster SD, Shata MT, Sherman KE, Blackard JT: **Genomic variability associated with the presence of occult hepatitis B virus in HIV co-infected individuals.** *J Viral Hepat* 2010, **17**:588-597.
12. Mphahlele MJ, Lukhwareni A, Burnett RJ, Moropeng LM, Ngobeni JM: **High risk of occult hepatitis B virus infection in HIV-positive patients from South Africa.** *J Clin Virol* 2006, **35**:14-20.
13. Larrubia JR: **Occult hepatitis B virus infection: a complex entity with relevant clinical implications.** *World J Gastroenterol* 2011, **17**:1529-1530.
14. Reina-San-Martin B, Cosson A, Minoprio P: **Lymphocyte polyclonal activation: a pitfall for vaccine design against infectious agents.** *Parasitol Today* 2000, **16**:62-67.
15. Morsica G, Ancarani F, Bagaglio S, Maracci M, Cicconi P, Cozzi Lepri A, Antonucci G, Bruno R, Santantonio T, Tacconi L, Baldelli F, Piscopo R, Santoro D, Lazzarin A, D'Arminio Monforte A, HepalCONA and the ICONA Study Groups: **Occult hepatitis B virus infection in a cohort of HIV-positive patients: correlation with hepatitis C virus coinfection, virological and immunological features.** *Infection* 2009, **37**:445-449.
16. Pinarbasi B, Onel D, Cosan F, Akyuz F, Dirlík N, Cakaloglu Y, Badur S, Besisik F, Demir K, Okten A, Kaymakoglu S: **Prevalence and virological features of occult hepatitis B virus infection in female sex workers who work uncontrolled in Turkey.** *Liver Int* 2009, **29**:227-230.
17. Weinberger KM, Bauer T, Böhm S, Jilg W: **High genetic variability of the group-specific a-determinant of hepatitis B virus surface antigen (HBsAg) and the corresponding fragment of the viral polymerase in chronic virus carriers lacking detectable HBsAg in serum.** *J Gen Virol* 2000, **81**:1165-1174.
18. Pourkarim MR, Lemey P, Amini-Bavil-Olyaei S, Houspie L, Verbeeck J, Rahman M, Maes P, Vanwijngaerden E, Nevens F, Van Ranst M: **Molecular characterization of hepatitis B virus strains circulating in Belgian patients co-infected with HIV and HBV: overt and occult infection.** *J Med Virol* 2011, **83**:1876-1884.
19. Chemin I, Trépo C: **Evolution of Hepatitis B and C serum markers: a still challenging issue.** *Liver Int* 2011, **31**:905-907.
20. Shi Y, Wu YH, Wei W, Zhang WJ, Yang J, Chen Z: **Association between occult hepatitis B infection and the risk of hepatocellular carcinoma: a meta-analysis.** *Liver Int* 2001.
21. Chemin I, Trépo C: **Clinical impact of occult HBV infections.** *J Clin Virol* 2005, **34**:S15-S21.
22. Pereira JS, Gonçalves NS, Silva C, Lazarini MS, Pavan MH, Fais VC, Gonçalves FL Jr: **HBV vaccination of HCV-infected patients with occult HBV infection and anti-HBc-positive blood donors.** *Braz J Med Biol Res* 2006, **39**:525-531.

doi:10.1186/1743-422X-8-527

Cite this article as: Cardona et al.: Unusual presentation of hepatitis B serological markers in an Amerindian community of Venezuela with a majority of occult cases. *Virology Journal* 2011 8:527.

Submit your next manuscript to BioMed Central and take full advantage of:

- Convenient online submission
- Thorough peer review
- No space constraints or color figure charges
- Immediate publication on acceptance
- Inclusion in PubMed, CAS, Scopus and Google Scholar
- Research which is freely available for redistribution

Submit your manuscript at
www.biomedcentral.com/submit

