

Impact of obesity on global and regional systolic function in children: a CMR study

Magalie Viallon, Julie Wacker, Patrick Clarysse, Maurice Beghetti, Nathalie Farpour-Lambert, Pierre Croisille

► **To cite this version:**

Magalie Viallon, Julie Wacker, Patrick Clarysse, Maurice Beghetti, Nathalie Farpour-Lambert, et al.. Impact of obesity on global and regional systolic function in children: a CMR study. 2011 SCMR/Euro CMR Joint Scientific Sessions, Feb 2011, Nice, France. BioMed Central, Abstracts of the 2011 SCMR/Euro CMR Joint Scientific Sessions, 13 (Suppl 1), pp.P310, 2011, Journal of Cardiovascular Magnetic Resonance. <10.1186/1532-429X-13-S1-P310>. <inserm-00663862>

HAL Id: inserm-00663862

<http://www.hal.inserm.fr/inserm-00663862>

Submitted on 27 Jan 2012

HAL is a multi-disciplinary open access archive for the deposit and dissemination of scientific research documents, whether they are published or not. The documents may come from teaching and research institutions in France or abroad, or from public or private research centers.

L'archive ouverte pluridisciplinaire **HAL**, est destinée au dépôt et à la diffusion de documents scientifiques de niveau recherche, publiés ou non, émanant des établissements d'enseignement et de recherche français ou étrangers, des laboratoires publics ou privés.

POSTER PRESENTATION

Open Access

Impact of obesity on global and regional systolic function in children: a CMR study

Magalie Viallon^{1*}, Julie Wacker², Patrick Clarysse³, Maurice Beghetti², Nathalie Farpour-Lambert², Pierre Croisille³

From 2011 SCMR/Euro CMR Joint Scientific Sessions
Nice, France. 3-6 February 2011

Introduction

Early detection of subclinical manifestations of obesity is important to initiate early effective treatment aiming at reversing the process. Recent studies using echocardiography tissue doppler or strain rate imaging have suggested the potential of motion tracking methods to characterize subtle systolic changes that may be associated with obesity in children (1,2).

Purpose

The aim of the present study was to investigate LV global and intra-myocardial regional systolic mechanics using CMR tagging in obese and non-obese adolescents in order to better characterize the subclinical early changes associated with obesity in children.

Methods

Standard CMR global and regional LV function study and cine-tagging with complementary myocardial tagging (CSPAMM) were performed in 25 obese children and 25 lean controls. Inclusion criteria were: no previous diagnosis of hypertension and treatment with anti-hypertensive/anti-diabetic drugs or drugs known to affect glucose and lipid metabolism; no history of familial hypertension/dyslipidemia; absence of diabetes or other chronic disease. The study protocol has been approved by the Mother&Child Ethics Committee. Magnitude CSPAMM images were processed using *InTag* post-processing toolbox (CreatiS, Lyon, France) implemented in OsiriX software (Geneva, Switzerland) to perform quantitative myocardial strain analysis. Motion estimation is based on the *Sine Wave Modeling* approach(3). Regional and global peak circumferential (E_{cc}), longitudinal(E_{ll}) strains as well as peak rotation and torsion were calculated.

Results

Gender distribution was not different between the two groups but obese children were older (14.1 ± 1 vs 12.9 ± 1.7 yrs, $p < 0.01$). BMI (30.4 ± 5.4 vs 18.9 ± 1.9), Z-score (2.58 vs 0.02 , $p < 0.0001$), systolic (112 ± 13 vs 101 ± 10 mmHg, $p < 0.01$) and diastolic blood pressure (55 ± 6 vs 52 ± 7 mmHg, $p < 0.05$) were greater whereas heart rate (71 ± 9 vs 77 ± 9 bpm, $p < 0.05$) was lower in the obese group. Measurements of LV mass (97.24 ± 21.2 vs 80.5 ± 20.2 g, $p < 0.0001$) and EDV (117.6 ± 32.6 vs 101.3 ± 23.7 ml, $p < 0.01$) were significantly greater in the obese children. Although global systolic function was not different among the two groups, obese children exhibited a significantly increased stroke volume (60.44 ± 6.74 vs 58.06 ± 5.7 ml, $p < 0.01$). In obese children, E_{cc} displayed only a trend to lower values in basal level (-0.15 ± 0.2 vs $-0.16 \pm 0.2\%$, $p = NS$) similarly to what was observed for E_{ll} (-0.14 ± 0.2 vs 0.15 ± 0.1 , $p = NS$). Torsion was also not significantly increased in obese children ($19.5 \pm 3.8^\circ$ vs $18.0 \pm 4.3^\circ$, $p = NS$).

Conclusion

If our data confirm that obese children have early morphological changes in LV mass, EDV and SV, regional circumferential and longitudinal strains are not yet altered from normal. The trend to increased torsion and increased wall thickness may be an adaptation to increased LV ED volume and may lead to early impairment of LV diastolic function.

Author details

¹Radiology department, Hôpital Universitaire de Genève, Geneva, Switzerland. ²Pediatric Cardiology Unit, Child and adolescents department, Hôpital Universitaire de Genève, Geneva, Switzerland. ³CREATIS-LRMN; CNRS UMR5220; INSERM U630; INSA-Lyon; Cardiology Hospital L. Pradel, Lyon University, Lyon, France.

Published: 2 February 2011

¹Radiology department, Hôpital Universitaire de Genève, Geneva, Switzerland
Full list of author information is available at the end of the article

References

1. Di Salvo G: *EurHeartJ* 2006.
2. Lorch SM: *JCardiometabSyndr* 2007.
3. Arts T: *IEEETransMedImaging* 2010.

doi:10.1186/1532-429X-13-S1-P310

Cite this article as: Viallon et al.: Impact of obesity on global and regional systolic function in children: a CMR study. *Journal of Cardiovascular Magnetic Resonance* 2011 **13**(Suppl 1):P310.

**Submit your next manuscript to BioMed Central
and take full advantage of:**

- Convenient online submission
- Thorough peer review
- No space constraints or color figure charges
- Immediate publication on acceptance
- Inclusion in PubMed, CAS, Scopus and Google Scholar
- Research which is freely available for redistribution

Submit your manuscript at
www.biomedcentral.com/submit

