

## **Polymorphisms in the interleukin 4, interleukin 13 and corresponding receptor genes are not associated with Systemic Sclerosis and do not influence gene expression**

Jca Broen, Philippe Dieude, M Vonk, L Beretta, B Rueda, A Herrick, J Worthington, N Hunzelmann, G Riemekasten, H Kiener, et al.

► **To cite this version:**

Jca Broen, Philippe Dieude, M Vonk, L Beretta, B Rueda, et al.. Polymorphisms in the interleukin 4, interleukin 13 and corresponding receptor genes are not associated with Systemic Sclerosis and do not influence gene expression. 5th European Workshop on Immune-Mediated Inflammatory Diseases, Dec 2010, Sitges-Barcelona, Spain. BioMed Central, 8 (Suppl 1), pp.P47, 2010, Journal of Translational Medicine. <10.1186/1479-5876-8-S1-P47>. <inserm-00663872>

**HAL Id: inserm-00663872**

**<http://www.hal.inserm.fr/inserm-00663872>**

Submitted on 27 Jan 2012

**HAL** is a multi-disciplinary open access archive for the deposit and dissemination of scientific research documents, whether they are published or not. The documents may come from teaching and research institutions in France or abroad, or from public or private research centers.

L'archive ouverte pluridisciplinaire **HAL**, est destinée au dépôt et à la diffusion de documents scientifiques de niveau recherche, publiés ou non, émanant des établissements d'enseignement et de recherche français ou étrangers, des laboratoires publics ou privés.



POSTER PRESENTATION

Open Access

# Polymorphisms in the interleukin 4, interleukin 13 and corresponding receptor genes are not associated with Systemic Sclerosis and do not influence gene expression

JCA Broen<sup>1\*</sup>, P Dieude<sup>2</sup>, M C Vonk<sup>1</sup>, L Beretta<sup>3</sup>, B Rueda<sup>4</sup>, A Herrick<sup>5</sup>, J Worthington<sup>5</sup>, N Hunzelmann<sup>6</sup>, G Riemekasten<sup>7</sup>, H Kiener<sup>8</sup>, R Scorza<sup>3</sup>, C P Simeon<sup>9</sup>, V Fonollosa<sup>9</sup>, P Carreira<sup>10</sup>, N Ortego-Centeno<sup>11</sup>, M A Gonzalez-Gay<sup>12</sup>, P Airò<sup>13</sup>, MJH Coenen<sup>14</sup>, A Aliprantis<sup>15</sup>, J Martin<sup>4†</sup>, Y Allanore<sup>16†</sup>, TRDJ Radstake<sup>1†</sup>

From 5th European Workshop on Immune-Mediated Inflammatory Diseases  
Sitges-Barcelona, Spain. 1-3 December 2010

## Aim

Polymorphisms in the interleukin 4 (IL4), interleukin 13 (IL13) and their corresponding receptors have previously been found associated with systemic sclerosis (SSc). In this study we aim to validate these previous observations and scrutinize their effects on gene expression.

## Patients and methods

We genotyped a cohort consisting of 1902 systemic sclerosis patients and 1503 healthy controls, derived from France, The Netherlands, Spain, United Kingdom, Italy and Germany. Taqman assays were used for genotyping three SNPs correlating with IL-4 and receptor; interleukin 4 alpha receptor Q576R (rs1801275), interleukin 4 RI75V (rs1805010), and -590C/T (rs2243250). In the IL-13 gene the following SNPs were genotyped; R130Q (rs20541), (-1112C/T), rs1800925 and rs6646259 (base 43163:G/A). In addition, we investigated the effect of these polymorphisms on corresponding gene expression with RT-PCR in B cells, T cells, plasmacytoid dendritic cells, monocytes and myeloid dendritic cells.

## Results

None of these polymorphisms was found to be enriched in the SSc population or in any SSc clinical subtype. In addition, we did not observe an effect on expression levels in the cell subtypes.

## Conclusions

Our data show that these polymorphisms do not play a role in SSc and do not influence gene expression levels.

## Author details

<sup>1</sup>Dept. of Rheumatology, Radboud University Nijmegen Medical Center, The Netherlands. <sup>2</sup>Université Diderot Paris 7, Service de Rhumatologie, Hôpital Bichat Claude Bernard, Paris, France. <sup>3</sup>Referral Center for Systemic Autoimmune Diseases, University of Milan, Italy. <sup>4</sup>Instituto de Parasitología y Biomedicina, CSIC, Granada, Spain. <sup>5</sup>Rheumatic Diseases Centre, University of Manchester, Salford Royal NHS Foundation Trust, UK. <sup>6</sup>Dept. of Dermatology, University of Cologne, Germany. <sup>7</sup>Dept. of Rheumatology and Clinical Immunology, Charité University Hospital and German Rheumatism Research Centre, a Leibniz institute. <sup>8</sup>Dept. of Internal Medicine, Division of Rheumatology, University of Vienna, Austria. <sup>9</sup>Servicio de Medicina Interna, Hospital Vall d'Hebron, Barcelona, Spain. <sup>10</sup>Servicio de Reumatología, Hospital 12 de Octubre, Madrid, Spain. <sup>11</sup>Servicio de Medicina Interna, Hospital Xeral-Calde, Lugo, Spain. <sup>12</sup>Servicio de Reumatología, Hospital Marques de Valdecillas, Santander, Spain. <sup>13</sup>Servizio di Reumatologia ed Immunologia Clinica, Spedali Civili, Brescia, Italia. <sup>14</sup>Dept. of Human Genetics, Radboud University Nijmegen Medical Center, The Netherlands. <sup>15</sup>Dept. of Immunology and Infectious Diseases, Harvard School of Public Health, Boston, MA, USA. <sup>16</sup>Université Paris Descartes, INSERM U781, Hôpital Necker, Paris, France and Université Paris Descartes, Service de Rhumatologie A, Hôpital Cochin, Paris, France.

Published: 25 November 2010

doi:10.1186/1479-5876-8-S1-P47

Cite this article as: Broen et al.: Polymorphisms in the interleukin 4, interleukin 13 and corresponding receptor genes are not associated with Systemic Sclerosis and do not influence gene expression. *Journal of Translational Medicine* 2010 **8**(Suppl 1):P47.

† Contributed equally

<sup>1</sup>Dept. of Rheumatology, Radboud University Nijmegen Medical Center, The Netherlands

Full list of author information is available at the end of the article