

Accelerated immune senescence in HIV infection

Victor Appay

► **To cite this version:**

Victor Appay. Accelerated immune senescence in HIV infection. 16th International Symposium on HIV and Emerging Infectious Diseases, Mar 2010, Marseille, France. pp.I30, 10.1186/1742-4690-7-S1-I30 . inserm-00664019

HAL Id: inserm-00664019

<https://www.hal.inserm.fr/inserm-00664019>

Submitted on 28 Jan 2012

HAL is a multi-disciplinary open access archive for the deposit and dissemination of scientific research documents, whether they are published or not. The documents may come from teaching and research institutions in France or abroad, or from public or private research centers.

L'archive ouverte pluridisciplinaire **HAL**, est destinée au dépôt et à la diffusion de documents scientifiques de niveau recherche, publiés ou non, émanant des établissements d'enseignement et de recherche français ou étrangers, des laboratoires publics ou privés.



INVITED SPEAKER PRESENTATION

Open Access

Accelerated immune senescence in HIV infection

Victor Appay

From 16th International Symposium on HIV and Emerging Infectious Diseases
Marseille, France. 24-26 March 2010

Aim

The prospect that immune activation or inflammation may be directly related to the increased incidence in HIV infected donors of manifestations that are reminiscent of the human aging process (i.e. cardiovascular disease, malignancies, osteoporosis, cognitive impairment, depression and frailty) is raising increasing concerns. On the immunological side, in addition to promoting viral replication as well as CD4⁺ T cell apoptosis, HIV associated immune activation may also lead to an accelerated decline of immune competence resembling the phenomenon of immunosenescence. Our aim is to explore further the potential relationship between immune activation, HIV disease progression and immunosenescence.

Materials and methods

We performed a comparative analysis of immunologic markers in different groups of HIV infected donors and healthy controls. The initial investigation of blood lymphocyte populations led us rapidly to turn our attention onto the lymphocyte primary immune resources, i.e. the CD34⁺ hematopoietic progenitor cells (studied directly from the blood of patients).

Results

Human aging and HIV-1 infection exhibit a number of parallels with regards to immunological attributes, that are evocative of premature immunosenescence in HIV-1 infected patients and reflect a reduced production of lymphocytes. Analyses of CD34⁺ hematopoietic progenitor cell number, phenotype and clonogenic potential underline a manifest impairment of primary immune resources with age or HIV-1 infection. Systemic immune activation emerges as a major correlate of altered lymphopoiesis, which can be partially reversed with prolonged antiretroviral therapy. Poor CD4⁺ T cell count

recovery despite successful virological response on anti-retroviral treatment is associated with persistent damage to the lymphopoietic system.

Discussion

Our findings provide new insights into the consequences of persistent immune activation in HIV-1 infection, and demonstrate the importance of primary hematopoietic resources in HIV pathogenesis and the response to antiretroviral treatments, but also more generally in the development of immunosenescence.

Published: 11 May 2010

doi:10.1186/1742-4690-7-S1-I30

Cite this article as: Appay: Accelerated immune senescence in HIV infection. *Retrovirology* 2010 **7**(Suppl 1):I30.

Submit your next manuscript to BioMed Central
and take full advantage of:

- Convenient online submission
- Thorough peer review
- No space constraints or color figure charges
- Immediate publication on acceptance
- Inclusion in PubMed, CAS, Scopus and Google Scholar
- Research which is freely available for redistribution

Submit your manuscript at
www.biomedcentral.com/submit

