

**Long-term cocaine use is associated with premature alterations in regional aortic strain and distensibility measured by magnetic resonance imaging**

Alban Redheuil, Chia-Ying Liu, Elie Mousseaux, David Bluemke, Joao Lima, Shenghan Lai

► **To cite this version:**

Alban Redheuil, Chia-Ying Liu, Elie Mousseaux, David Bluemke, Joao Lima, et al.. Long-term cocaine use is associated with premature alterations in regional aortic strain and distensibility measured by magnetic resonance imaging. 13th Annual SCMR Scientific Sessions, Jan 2010, Phoenix, France. BioMed Central, 12 (Suppl 1), pp.P146, 2010, Journal of Cardiovascular Magnetic Resonance. <10.1186/1532-429X-12-S1-P146>. <inserm-00668476>

**HAL Id: inserm-00668476**

**<http://www.hal.inserm.fr/inserm-00668476>**

Submitted on 9 Feb 2012

**HAL** is a multi-disciplinary open access archive for the deposit and dissemination of scientific research documents, whether they are published or not. The documents may come from teaching and research institutions in France or abroad, or from public or private research centers.

L'archive ouverte pluridisciplinaire **HAL**, est destinée au dépôt et à la diffusion de documents scientifiques de niveau recherche, publiés ou non, émanant des établissements d'enseignement et de recherche français ou étrangers, des laboratoires publics ou privés.

Poster presentation

Open Access

## Long-term cocaine use is associated with premature alterations in regional aortic strain and distensibility measured by magnetic resonance imaging

Alban Redheuil<sup>\*1</sup>, Chia-Ying Liu<sup>1</sup>, Elie Mousseaux<sup>2</sup>, David Bluemke<sup>3</sup>, Joao Lima<sup>1</sup> and Shenghan Lai<sup>4</sup>

Address: <sup>1</sup>The Johns Hopkins Hospital, Baltimore, USA, <sup>2</sup>University of Paris Descartes, HEGP and INSERM U678, Paris, France, <sup>3</sup>Department of Radiology and Imaging Sciences, National Institutes of Health, Bethesda, USA and <sup>4</sup>Department of Epidemiology, Johns Hopkins School of Hygiene and Public Health, Baltimore, USA

\* Corresponding author

from 13th Annual SCMR Scientific Sessions  
Phoenix, AZ, USA. 21-24 January 2010

Published: 21 January 2010

*Journal of Cardiovascular Magnetic Resonance* 2010, **12**(Suppl 1):P146 doi:10.1186/1532-429X-12-S1-P146

This abstract is available from: <http://jcmr-online.com/content/12/S1/P146>

© 2010 Redheuil et al; licensee BioMed Central Ltd.

### Introduction

Long-term cocaine use has been associated with regional systolic and diastolic left ventricular dysfunction, hypertension and aortic dissection. The repetitive cardiovascular stress induced by cocaine use may lead to premature arterial stiffening but data on this relationship are scarce and inconsistent.

### Purpose

We studied the relationship between regional aortic stiffness measured by MRI and long-term cocaine use.

### Methods

We enrolled 46 consecutive subjects from an addiction clinic: 33 long term cocaine users (13 men, 20 women, mean age:  $46 \pm 7$  yrs, mean years of cocaine use:  $15 \pm 8$ ) and 13 non-cocaine users (6 men, 7 women, mean age:  $43 \pm 9$  yrs). Aortic stiffness of the ascending and descending aorta was determined by MRI from aortic strain (AS: relative difference in cross-sectional area) and distensibility (AD: aortic strain normalized by pulse pressure) using an automated contours detection method applied to modulus images of a phase-contrast acquisition perpendicular to the ascending aorta (Art-Fun, INSERM). Blood pressure was measured by a brachial cuff during aortic MRI. Drug and smoking habitus was determined by a standardized questionnaire.

### Results

SBP was slightly higher in the cocaine group vs. non-users ( $130 \pm 18$  vs.  $123 \pm 37$  mmHg) but pulse pressure was comparable ( $47 \pm 12$  vs.  $46 \pm 10$  mmHg respectively) and hypertension was similarly distributed (30% vs. 38%). Total cholesterol was also slightly higher ( $174 \pm 36$  vs.  $168 \pm 35$  mg) and current cigarette smoking more prevalent in the cocaine group (88% vs. 30%). Only 3 subjects of the cocaine group were diabetics and no diabetes in non-users. Aortic strain and distensibility were lower in the cocaine group vs. non-users as summarized in Table 1. Univariate analysis showed a negative correlation between aortic strain and distensibility and duration of cocaine use. After further adjustment for age, gender, cholesterol, smoking and diabetes the duration of cocaine use was an independent predictor of descending aortic function with a significant average decrease in strain of 2.5% ( $p = 0.02$ ) and a trend for a decrease in distensibility of  $4.5 \times 10^{-3} \text{ kPa}^{-1}$  ( $p = 0.14$ ) for 1 year of cocaine use.

### Conclusion

Long-term cocaine use is associated with premature regional stiffening of the aorta. This association was stronger for the descending aorta for which duration of cocaine use was found to be an independent predictor of vascular function beyond the effects of age and traditional risk factors.

**Table 1: Comparison of the aortic ascending and descending strain and distensibility of the cocaine**

	Non-Users Controls	Cocaine Use
Ascending Aortic Strain, %	18 ± 4	17 ± 8
Ascending Aortic Distensibility, 10 <sup>-3</sup> .kPa <sup>-1</sup>	35 ± 13	29 ± 16
Descending Aortic Strain, %	18 ± 6	16 ± 5
Descending Aortic Distensibility, 10 <sup>-3</sup> .kPa <sup>-1</sup>	31 ± 14	27 ± 13

Publish with **BioMed Central** and every scientist can read your work free of charge

*"BioMed Central will be the most significant development for disseminating the results of biomedical research in our lifetime."*

Sir Paul Nurse, Cancer Research UK

Your research papers will be:

- available free of charge to the entire biomedical community
- peer reviewed and published immediately upon acceptance
- cited in PubMed and archived on PubMed Central
- yours — you keep the copyright

Submit your manuscript here:  
[http://www.biomedcentral.com/info/publishing\\_adv.asp](http://www.biomedcentral.com/info/publishing_adv.asp)

