

Activation of intestinal peroxisome proliferator-activated receptor- α increases high-density lipoprotein production.

Sophie Colin, Olivier Briand, Véronique Touche, Kristiaan Wouters, Morgane Baron, François Pattou, Rémy Hanf, Anne Tailleux, Giulia Chinetti, Bart Staels, et al.

► To cite this version:

Sophie Colin, Olivier Briand, Véronique Touche, Kristiaan Wouters, Morgane Baron, et al.. Activation of intestinal peroxisome proliferator-activated receptor- α increases high-density lipoprotein production.. European Heart Journal, Oxford University Press (OUP): Policy B, 2013, 34 (32), pp.2566-74. <10.1093/eurheartj/ehs227>. <inserm-00730530>

HAL Id: inserm-00730530

<http://www.hal.inserm.fr/inserm-00730530>

Submitted on 1 Jul 2013

HAL is a multi-disciplinary open access archive for the deposit and dissemination of scientific research documents, whether they are published or not. The documents may come from teaching and research institutions in France or abroad, or from public or private research centers.

L'archive ouverte pluridisciplinaire **HAL**, est destinée au dépôt et à la diffusion de documents scientifiques de niveau recherche, publiés ou non, émanant des établissements d'enseignement et de recherche français ou étrangers, des laboratoires publics ou privés.

Activation of intestinal peroxisome proliferator-activated receptor- α increases high-density lipoprotein production

Sophie Colin¹, Olivier Briand¹, Véronique Touche¹, Kristiaan Wouters¹, Morgane Baron¹, François Pattou^{2,3}, Rémy Hanf⁴, Anne Tailleux¹, Giulia Chinetti¹, Bart Staels^{1*}, Sophie Lestavel¹

¹ Récepteurs nucléaires, maladies cardiovasculaires et diabète EGID FR 3508 INSERM : U1011, Institut Pasteur de Lille, Université Lille II - Droit et santé, 1 rue du Prof Calmette 59019 Lille Cedex, FR

² Thérapie cellulaire du diabète INSERM : U859, Université Lille II - Droit et Santé, Faculté de Médecine 1 Place de Verdun Pole Recherche Lille 59045 Lille Cedex, FR

³ Service de Chirurgie Générale et Endocrinienne CHRU Lille, Hôpital Claude Huriez, 2 Avenue Oscar Lambret - 59037 Lille Cedex, FR

⁴ Genfit Entreprse Biopharmaceutique GENFIT Loos, Parc Eurasanté, 885 Avenue Eugène Avinée 59120 Loos, FR

* Correspondence should be addressed to: Bart Staels <bart.staels@pasteur-lille.fr >

Abstract

Aims

Peroxisome Proliferator-Activated Receptor (PPAR) α is a transcription factor controlling lipid metabolism in liver, heart, muscle and macrophages. PPAR α -activation increases plasma HDL-cholesterol and exerts hypotriglyceridemic actions via the liver. However, the intestine expresses PPAR α , produces HDL and chylomicrons and is exposed to diet-derived PPAR α ligands. Therefore, we examined the effects of PPAR α -activation on intestinal lipid and lipoprotein metabolism.

Methods and Results

The impact of PPAR α -activation was evaluated in term of HDL-related gene expression in mice, *ex-vivo* in human jejunal biopsies and in Caco-2/TC7 cells. ApoAI/HDL secretion, cholesterol esterification and trafficking were also studied *in-vitro*.

In parallel to improving plasma lipid profiles and increasing liver and intestinal expression of fatty-acid-oxidation genes, treatment with the dual PPAR α / δ -ligand GFT505 resulted in a more pronounced increase of plasma HDL compared to fenofibrate in mice. GFT505, but not fenofibrate, increased the expression of HDL-production genes such as apolipoprotein-AI and ATP-Binding-Cassette-A1 transporter in murine intestines. A similar increase was observed upon PPAR α -activation of human biopsies and Caco-2/TC7 cells. Additionally, HDL secretion by Caco-2/TC7 cells increased. Moreover, PPAR α -activation decreased the cholesterol-esterification capacity of Caco-2/TC7 cells, modified cholesterol trafficking and reduced apolipoprotein-B secretion.

Conclusions

PPAR α -activation reduces cholesterol esterification, suppresses chylomicron- and increases HDL-secretion by enterocytes. These results identify the intestine as a target organ of PPAR α -ligands with entero-hepatic tropism to reduce atherogenic dyslipidemia.

MESH Keywords Animals ; Apolipoproteins B ; metabolism ; Butyrates ; pharmacology ; Caco-2 Cells ; Cells, Cultured ; Chalcones ; pharmacology ; Enterocytes ; metabolism ; Esterification ; physiology ; Fatty Acids ; metabolism ; Female ; Humans ; Jejunum ; metabolism ; Lipoproteins, HDL ; metabolism ; Mice ; Mice, Knockout ; PPAR alpha ; antagonists & inhibitors ; physiology ; Phenylurea Compounds ; pharmacology ; Propionates ; pharmacology

Author Keywords PPAR α ; intestine ; HDL ; dyslipidemia

Introduction

High levels of low-density-lipoprotein (LDL)-cholesterol and low levels of high-density-lipoprotein (HDL)-cholesterol are risk factors for cardiovascular disease (1). In association with lifestyle intervention, LDL-cholesterol lowering therapy with statins is highly efficient to reduce cardiovascular risk, but has only modest effects on HDL-cholesterol (2). Furthermore, low HDL-cholesterol is an independent risk factor which contributes to the persistent residual cardiovascular risk in statin-treated patients (3; 4). Therefore, raising HDL may be a promising strategy to attenuate residual cardiovascular risk in statin-treated patients.

In enterocytes, dietary lipids are packaged with apolipoprotein-B48 (apoB48) into chylomicrons secreted into the lymph (5). Several studies demonstrated that post-prandial chylomicrons contribute to lipid accumulation in the arterial wall and to atherogenesis (6). Animal studies have shown that the intestine produces HDL (7). Indeed, intestinal-specific deletion of ATP-Binding-Cassette-A1 transporter (ABCA1) in mice results in a 30% decrease of HDL-cholesterol levels (8). Furthermore, exercise-training enhances HDL-cholesterol in rats in correlation with increased intestinal ABCA1 mRNA expression (9). Interestingly, human intestinal mucosa was shown to produce apolipoprotein-AI (apoAI) (10) and HDL (11).

Peroxisome Proliferator-Activated Receptor (PPAR) α is a transcription factor belonging to the nuclear receptor superfamily and activated by fibrates, eicosanoids and fatty-acids (12; 13). PPAR α controls lipid and glucose metabolism in several tissues and cell types including liver, heart, kidney, adipose tissue and macrophages (14–16). PPAR α -activation enhances hepatic fatty-acid-oxidation and peripheral triglyceride clearance, thus reducing hypertriglyceridemia (13; 17). PPAR α increases hepatic apoAI synthesis and ABCA1 expression, resulting in increased plasma HDL-cholesterol (18; 19). In addition, PPAR α -activation controls cholesterol trafficking in human macrophages and raises cholesterol efflux *via* ABCA1 contributing to increased reverse cholesterol transport (15; 20). Since diet-derived compounds (fatty-acids and phospholipids) can activate PPAR α and since the intestine is the site of dietary lipid absorption, it is highly exposed to PPAR α -ligands (21; 22). The role of PPAR α in the intestine has been addressed in only a few studies. Previous studies in rodents did not evidence pronounced effects of the PPAR α -ligand fenofibrate (FF) in the intestine contrary to the liver, possibly because of its rapid urinary excretion and hence relatively low intestinal exposure (24). However, a genome wide analysis of intestinal RNA of mice treated with Wy14643 revealed that PPAR α controls numerous metabolic pathways and in particular lipid metabolism (25), pointing to a role for PPAR α in this organ.

We thus investigated whether intestinal PPAR α -activation regulates HDL production. Treatment of mice with two different PPAR-agonists, FF, a pure PPAR α -agonist and GFT505 (26), a PPAR α / δ -modulator, which, unlike FF, undergoes extensive enterohepatic cycling, resulted in comparable hypotriglyceridemic effects. However, only GFT505 treatment increased apoAI and ABCA1 mRNA levels in the murine intestine, an effect which was associated with a more pronounced increase of plasma HDL levels. These effects were not observed in PPAR α -deficient mice. Using the human Caco-2/TC7 model cultured on porous filters, we found that PPAR α -activation reduces secretion of chylomicrons while enhancing HDL production probably by increasing ABCA1 expression, apoAI secretion and by reducing cholesterol esterification. The physiological relevance of these regulatory processes was verified in human jejunal biopsies. Collectively, our data suggest that the intestine is a target organ for entero-hepato-tropic PPAR α -ligands to increase HDL, an effect which may result in a reduction of residual cardiovascular risk.

Materials and methods

For details, see supplementary materials and methods (supp.M&M).

Animal study

Wild-type (+/+) and homozygous (-/-) PPAR α -deficient female mice in the apoE2-KI background (12 week-old) fed a western-diet were treated for 14d with GFT505 (10 or 30mpk) or FF (200mpk) or carboxy-methyl-cellulose (0.5%) (27).

Human intestine culture

Intestinal biopsies were recovered during gastric bypass surgery from obese patients enrolled in the ABOS study (ClinicalTrials.gov; NCT01129297).

Caco-2/TC7 cell culture and PPAR α knock-down

Caco-2/TC7 were grown as described (28). For stable PPAR α invalidation (29) and culture details, see supp.M&M.

HDL preparation

HDL (d=1.12–1.21g/ml) from human plasma and basolateral media of Caco-2/TC7 were prepared by sequential ultracentrifugations. For details in preparation, electron microscopy and apolipoprotein quantification, see supp.M&M.

Post-prandial micelle preparation

Synthetic micelles prepared as described (28).

Confocal microscopy

For details, see supp.M&M and (30).

Cholesterol esterification assay

Caco-2/TC7 were incubated with [3 H]-cholesterol-micelles (7.5 μ Ci/well), lipids were extracted and separated by TLC.

Macrophage cell culture and cholesterol efflux

Mononuclear cells were isolated from blood of healthy donors, cholesterol-loaded with [3 H]-cholesterol-AcLDL and efflux assays performed as described (15).

Statistics

For details, see supp.M&M.

Results

PPAR α activation in mice and in human jejunal biopsies increases the expression of genes involved in intestinal HDL production

To determine whether PPAR α -activation regulates HDL production in the small intestine, apoE2-KI mice, which display a similar lipid-response to PPAR α -agonists as humans (36), were fed a western-diet with daily oral administration of FF or the dual PPAR α/δ -agonist GFT505 for 14d. As reported (36), FF decreased plasma triglycerides (TG) and total-cholesterol (tab1). Similarly, GFT505 lowered plasma TG and total-cholesterol. Whereas plasma HDL-cholesterol increased approximately 2.5-fold upon treatment with FF, the increase of HDL-cholesterol was significantly more pronounced upon GFT505 (tab1). FF and GFT505 induced the hepatic expression of PPAR α target genes, such as Acyl-CoenzymeA-oxidase-1 (ACOX1) at a similar extent (supp.fig 1) indicating equipotent dosing. The effects of GFT505 on TG and HDL-cholesterol were abolished in PPAR α -deficient apoE2-KI mice, demonstrating that the effects are PPAR α -dependent (supp.tab1). Interestingly, a decrease, albeit less pronounced, of total and nonHDL-cholesterol was maintained in GFT505-treated PPAR α -deficient apoE2-KI mice, suggesting a contribution of its PPAR δ activity on these lipid parameters (supp.tab1). GFT505-treated mice displayed a PPAR α dependent higher ABCA1 and apoAI gene expression in the small intestine compared to untreated mice (fig 1A, fig 1B), whereas FF did not regulate ABCA1 nor apoAI mRNA expression (fig 1C, fig 1D). Treatment of mice with GW0742, a pure PPAR δ -agonist, did not modify ABCA1 nor ApoAI gene expression (not shown).

To determine whether PPAR α -activation also regulates these genes in human intestine, human jejunal biopsies were treated with GFT505 and a synthetic PPAR α -ligand, GW7647, for 18h. Both compounds significantly increased ABCA1 and apoAI mRNA expression (fig 1E, fig 1F).

PPAR α -activation increases the production of HDL in Caco-2/TC7

Based on observations in mice (8), we hypothesized that the up-regulation of ABCA1 and apoAI gene expression observed upon PPAR α -activation would translate in increased intestinal HDL production. To test this, we used an *in-vitro* model of the human intestinal barrier, polarized Caco-2/TC7 cells grown on porous filters (28; 32). PPAR α expression was knocked-down (ShPPAR α) or not (ShControl) by lentiviral infection and cells were subsequently incubated with GW7647. HDL secreted by Caco-2/TC7 were isolated by ultracentrifugation and analyzed by electron microscopy. PPAR α -activation increased the number of HDL secreted at the basolateral side of Caco-2/TC7 (fig 2A). ApoAI secreted in the HDL fraction increased when non-infected Caco-2/TC7 and ShControl-infected cells were treated with GW7647, whereas apoAI secretion was not modified upon treatment of ShPPAR α -infected cells by GW7647 (fig 2B), demonstrating that this effect is PPAR α -dependent. Interestingly, the quantity of apoAI in the chylomicron fraction was not modified (not shown). To evaluate the HDL functionality, we investigated their capacity to promote cholesterol efflux from macrophages. Human monocyte-derived macrophages were cholesterol-loaded with [³H]-cholesterol-AcLDL and subsequently incubated with HDL secreted by Caco-2/TC7 treated with GW7647 for 24h. Equivalent volumes of isolated HDL from non-infected or ShControl-infected Caco-2/TC7 treated with GW7647 enhanced macrophage cholesterol efflux compared to HDL isolated from vehicle-treated cells (fig 2C). HDL from ShPPAR α -infected cells treated with GW7647 did not show enhanced efflux capacity compared to HDL from untreated cells. ABCA1 gene expression increased upon GW7647 treatment in a PPAR α -dependent manner (fig 2D). A similar up-regulation was observed with GFT505 (supp.fig 2). In contrast to the *in-vivo* and *ex-vivo* response, apoAI mRNA was not induced by PPAR α -agonists in Caco-2/TC7 (not shown).

PPAR α -activation modulates cholesterol distribution, increases cytoplasmic lipid droplet formation and reduces apoB secretion in Caco-2/TC7

To test the effect of PPAR α -activation on intestinal cholesterol absorption known to occur in sequential steps, from uptake at the brush border membrane and transport into the cell to packaging and secretion of lipoproteins at the basolateral side of enterocytes (33), Caco-2/TC7 were incubated for 24h with [³H]-cholesterol-micelles and activated with GW7647 24h before and 24h after cholesterol-loading. Cholesterol content in the apical medium, the cellular compartment and the basolateral medium of Caco-2/TC7 was not modulated by GW7647 (not shown), suggesting that PPAR α regulates neither total cholesterol uptake nor efflux by enterocytes. Next, GW7647-treated Caco-2/TC7 were incubated with fluorescent-NBD-cholesterol-containing micelles for 4h, followed by non-fluorescent micelles for an additional 4h period. Quantitative analysis of fluorescence intensity at each optical slice showed that, without affecting the total amount of absorbed cholesterol (assessed by radioactive assay), GW7647 treatment modifies cholesterol distribution inducing retention of cholesterol in the subapical compartments (fig 3A). Interestingly, as shown by confocal microscopy (fig 3B), GW7647 also increased the size of cholesterol-containing lipid droplets in the apical and basolateral compartments.

Since PPAR α -activation in Caco-2/TC7 does not affect the total amount of cholesterol uptake and efflux, we hypothesized that PPAR α -activation may decrease apoB-containing lipoprotein secretion to counterbalance the increase of secreted HDL. Indeed, GW7647 decreased apoB secretion in the basolateral compartment of Caco-2/TC7 (fig 3C) as previously shown (34).

PPAR α -activation decreases cholesterol esterification capacity of Caco-2/TC7

Cholesterol trafficking from the plasma membrane to the endoplasmic reticulum is a limiting step for its esterification and subsequent secretion in chylomicrons (35). To address whether, by interfering with cholesterol trafficking, PPAR α reduces cholesterol esterification, Caco-2/TC7 were incubated for 24h with [3 H]-cholesterol-micelles and activated with GW7647 24h before and 24h after cholesterol loading. PPAR α -activation decreased the quantity of [3 H]-cholesteryl esters (CE) assayed by TLC both in the cells (fig 4A) and secreted in the basolateral compartment (fig 4B). The decreased CE production does not result only from a perturbation of cholesterol trafficking since ACAT-2 (acyl-coenzymeA:cholesterol-acyltransferase-2) gene expression was also decreased (fig 4C).

PPAR α -activation increases the expression of lipid droplet formation-genes in Caco-2/TC7 cells

To investigate the mechanism behind the increase of cytoplasmic lipid droplet size upon PPAR α -activation (fig 3B), we analyzed the expression of lipid droplet-associated proteins. PPAR α -activation increased mRNA levels of perilipin-2 (PLIN2), which coats lipid droplets with its partner ABHD5 (AB-hydrolase-domain-containing-5) (fig 5A) (36). In line, ABHD5 gene expression was also increased (fig 5B). Furthermore, the increase of PLIN2 was strongly reduced and the increase of ABHD5 was totally abolished in ShPPAR α -infected cells treated with GW7647 (supp.fig 3A, supp.fig 3B). In addition, PLIN2 and ABHD5 mRNA levels increased only in intestines of GFT505-treated PPAR α +/, but not PPAR α -/- apoE2-KI mice (supp.fig 3C, suppl.fig 3D). The increased PLIN2 and ABHD5 gene expression was also observed, albeit less pronounced, in human jejunal biopsies treated with GW7647 or GFT505 (fig 5C, fig 5D).

PPAR α -activation increases fatty-acid-oxidation genes in human enterocytes and murine intestines

In addition to the decreased apoB secretion (fig 3C) and in accordance with previous data showing that intestinal PPAR α -activation suppresses postprandial lipidemia by increasing fatty-acid-oxidation (25; 34), PPAR α ligands increased mRNA levels of fatty-acid-oxidation genes including Carnitine-Palmitoyl-Transferase-1 (CPT-1A), ACOX1 and long-chain Acyl-CoA synthetase family member 5 (ACSL5) (fig 6A, fig 6B, fig 6C) in Caco-2/TC7. CPT-1A and ACOX1 mRNA levels were increased in human biopsies treated with PPAR α -ligands (fig 6D, fig 6E). The effect of GW7647 on CPT-1A and ACSL5 was strongly reduced in treated ShPPAR α -infected cells and totally abolished for ACOX1 (supp.fig 4A, supp.fig 4B, supp.fig 4C). GFT505 treatment increased the intestinal expression of fatty-acid-oxidation genes in a PPAR α -dependent manner in apoE2-KI mice (supp.fig 4D, supp.fig 4E). The expression of fatty-acid-oxidation genes was induced in intestines of FF-treated mice, in contrast to the lack of regulation of HDL-related genes, suggesting that FF acts as a selective PPAR modulator in the intestine (supp.fig 4F, supp.fig 4G, supp.fig 4H).

Interestingly, PPAR α -ligands increased cAMP-responsive-element-binding protein-3-like-3 (CREB3L3) mRNA levels in Caco-2/TC7 (supp.fig 5A), a transcription factor involved in the regulation of TG metabolism for which mutations were recently associated with hypertriglyceridemia in humans (37). The induction of CREB3L3 was observed in human biopsies treated with GW7647 and GFT505 (supp.fig 5B). Furthermore, PPAR α -ligands did not regulate this gene in ShPPAR α -infected cells nor in PPAR α -deficient mice (supp.fig 5C, supp.fig 5D). Collectively, these results indicate that PPAR α -activation increases the fatty-acid-oxidation pathway, the transcription factor CREB3L3 and increases lipid droplet size.

Discussion

In addition to the liver, the intestine has emerged more recently as a central organ participating in lipid metabolism by regulating lipid absorption, trafficking and lipoprotein metabolism through its involvement in the production of chylomicrons (5; 32), HDL (8) and in trans-intestinal cholesterol excretion (38). Thus, both the liver and intestine are involved in the regulation of whole body lipoprotein metabolism and play a role in the pathophysiology of atherogenic dyslipidemia (39; 40). Fibrates, PPAR α -agonists, are often used in the treatment of dyslipidemia characterized by high TG and low HDL-cholesterol levels. Until now, the effects of PPAR α -agonists on the improvement of lipid profiles were essentially attributed to their action on the liver and the arterial wall (41; 42). However, PPAR α is also expressed in the intestine. In a genome wide analysis of intestinal RNA isolated from PPAR α -ligand-treated mice, intestinal PPAR α was found to regulate fatty-acid metabolism (25). Moreover, PPAR α -activation decreased the secretion of apoB by enterocytes (34). Therefore, we studied whether intestinal PPAR α -activation could contribute to the control of HDL metabolism.

Using a humanized mouse model of mixed dyslipidemia, the apoE2-KI mouse, we show in line with previous studies that FF and GFT505-treatments improve the lipid profile by decreasing TG, total cholesterol and by increasing HDL-cholesterol (27; 31). We show that PPAR α mediates the effects of GFT505 on TG and HDL-cholesterol levels. Although all clinical studies illustrate the hypotriglyceridemic efficiency of PPAR α -agonists, the effects on HDL levels are variable. Such discrepancies could be explained by

differences in population selection and heterogeneity or in the molecule used (43–45). In our study, the HDL-increasing effect of GFT505 was more pronounced than FF at concentrations (10mpk vs 100mpk) inducing similar expression levels of ACOX1 in the liver. Although FF did induce intestinal fatty-acid-oxidation-related genes, suggesting that FF can activate intestinal PPAR α , only GFT505 induced expression of intestinal HDL production-related genes such as apoAI and ABCA1. Thus, GFT505 and FF differ in the spectrum of regulated target genes, giving the rationale to design new selective PPAR modulators. Furthermore, the different activity of fenofibrate vs GFT505 on intestinal gene expression is due to the extensive enterohepatic recycling and fecal excretion of GFT505, whereas fenofibrate is excreted in the urine (Rubenstrunk A and al., manuscript in preparation).

Using the Caco-2/TC7 model (28), we confirmed that PPAR α -ligands decrease apoB secretion (34), suggesting a decrease of TG-rich lipoprotein secretion. Moreover, we showed that PPAR α -activation increased ABCA1 gene expression and apoAI secretion. Although the previously reported absence of apoAI mRNA regulation (46) demonstrates the limits of the Caco-2/TC7 model (apoAI mRNA was increased in PPAR α -agonist-treated human jejunal biopsies), PPAR α -activation increased the number of HDL secreted, which in turn enhanced cholesterol efflux from human monocyte-derived macrophages.

As we did not observe any difference in the levels of apical cholesterol uptake nor in the levels of cholesterol secreted at the basal side of treated Caco-2/TC7, we hypothesized that PPAR α -activation induces a shift of the absorbed cholesterol from the chylomicron to the HDL production pathway. This hypothesis is strengthened by the fact that less cholesterol was esterified in treated cells. The decrease of esterification may be explained by a decrease of ACAT-2 gene expression which results, as shown in human hepatic cells and macrophages (47; 48), in a decrease of ACAT-2 activity which is also under the control of substrate availability at its site of action, the endoplasmic reticulum (35). By investigating cholesterol trafficking using a fluorescent cholesterol derivative, we observed cholesterol retention in the sub-apical compartment of GW7647-treated Caco-2/TC7 and an increase of lipid droplet size concomitantly with an increase of PLIN2 expression, which is necessary for lipid droplet stabilization, and ABHD5, sequestered by PLIN2 at the lipid surface (49). We did not observe any regulation of Rab proteins which are required for vesicular trafficking, nor NPC1 and NPC2 (not shown) which contrasts to macrophages (47). Thus, the mechanism responsible for the effects of PPAR α -activation on intracellular cholesterol trafficking remains to be clarified. However, the decrease of ACAT-2 gene expression was concomitant to an increase of CPT-1A expression, which could decrease the availability of fatty-acid for cholesterol esterification, as shown in macrophages (47). In line, PPAR α -activation increased the esterase-arylacetamide-deacetylase (AADC) (not shown) and ACSL5 gene expression, both associated with a reduction of TG accumulation and an increase of fatty-acid-oxidation in rodent hepatocytes (50). Different mechanisms could thus act in concert to divert cholesterol from esterification: decrease of ACAT-2 gene expression, reduction of fatty-acid availability through an increase of CPT-1A expression and fatty-acid-oxidation enzymes and/or retention of cholesterol in lipid droplets.

It has been recently shown that a mutation of CREB3L3 in humans is associated with hypertriglyceridemia. CREB3L3-deficient mice are hypertriglyceridemic and display defective TG clearance (37). Interestingly, we observed a PPAR α -dependent up-regulation of CREB3L3 expression in GW7647-treated Caco-2/TC7 and in GFT505-treated mice. The importance of the regulation of this gene expression and its implication in the modulation of lipid homeostasis upon PPAR α -activation still remain to be established.

The results of our study collectively show that the increase of ABCA1 expression, the modification in cholesterol trafficking and the decreased cholesterol esterification capacity induced by PPAR α -activation potentially contribute to the overall effects on intestinal secretion of apoAI-containing HDL by Caco-2/TC7. The transcriptional regulation of HDL-related genes was observed in mice and in human jejunal biopsies. Therefore, the decrease of TG and the increase of HDL levels observed in PPAR α -agonist-treated patients could be due, in addition to hepatic actions, to effects in the intestine. Our work strengthens the idea that the intestine is an important target organ for strategies to increase HDL production via activation of intestinal PPAR α . Moreover, our data demonstrate the relevance of designing new specific PPAR α modulators with appropriate tissue and gene selective profiles.

Acknowledgements:

Funding:

This work was supported by grants from Université Lille 2, Région Nord/Pas-de-Calais, the FEDER and the “Fondation Leducq”.

We thank M. Tardivel, C. Allet and A. Loyens of the BICeL (IMPRT-IFR114) for access to confocal microscopy and electron microscopy platforms and technical advices.

Footnotes:

Disclosures: No conflicts of interest exist

Abbreviations

ABCA1 : ATP-Binding Cassette A1 transporter

ABHD5 : AB-Hydrolase Domain containing 5

ACAT-2 : Acyl-coenzyme A: Cholesterol Acyltransferase-2

ACOX1 : Acyl-Coenzyme A oxidase 1
 ACSL5 : Acyl-CoA Synthetase Long-chain family member 5
 apo : apolipoprotein
 CE : Cholesteryl Ester
 CPT-1 : Carnitine Palmitoyl Transferase I
 CREB3L3 : cAMP responsive element binding protein 3 like 3
 FC : Free Cholesterol
 FF : Fenofibrate
 HDL : High Density Lipoproteins
 LDL : Low Density Lipoproteins
 PLIN2 : Perilipin 2
 PPAR : Peroxisome proliferator-activated receptor

References:

Bibliography

- 1 . Castelli W , Garrison R , Wilson P , Abbott R , Kalousdian S , Kannel W . Incidence of coronary heart disease and lipoprotein cholesterol levels. the Framingham study . *JAMA* . 1986 ; 256 : 2835 - 2838
- 2 . McTaggart F , Jones P . Effects of statins on HDL: a potential contribution to cardiovascular benefit . *Cardiovasc Drugs Ther* . 2008 ; 22 : 321 - 338
- 3 . Baigent C , Keech A , Kearney P , Blackwell L , Buck G , Pollicino C , Kirby A , Sourjina T , Peto R , Collins R , Simes R , CTTC . Efficacy and safety of cholesterol-lowering treatment: prospective meta-analysis of data from 90,056 participants in 14 randomised trials of statins . *Lancet* . 2005 ; 366 : 1267 - 1278
- 4 . Barter P , Gotto A , LaRosa J , Maroni J , Szarek M , Grundy S , Kastelein J , Bittner V , Fruchart J . Treating to New Targets Investigators . HDL-cholesterol, very low levels of LDL-cholesterol, and cardiovascular events . *N Engl J Med* . 2007 ; 357 : 1301 - 1310
- 5 . Black D . Development and physiological regulation of intestinal lipid absorption. I. Development of intestinal lipid absorption: cellular events in chylomicron assembly and secretion . *Am J Physiol Gastrointest Liver Physiol* . 2007 ; 293 : G519 - 24
- 6 . Su J , Nzekwu M , Cabezas M , Redgrave T , Proctor S . Methods to assess impaired post-prandial metabolism and the impact for early detection of cardiovascular disease risk . *Eur J Clin Invest* . 2009 ; 39 : (9) 741 - 754
- 7 . Green P , Tall A , Glickman R . Rat intestine secretes discoid HDL . *J Clin Invest* . 1978 ; 61 : 528 - 534
- 8 . Brunham L , Kruit J , Pape T , Parks J , Kuipers F , Hayden M . Intestinal ABCA1 directly contributes to HDL biogenesis in-vivo . *J Clin Invest* . 2006 ; 116 : 1052 - 1062
- 9 . Khabazian B , Ghanbari-Niaki A , Safarzadeh-Golpordesari A , Ebrahimi M , Rahbarzadeh F , Abednazari H . Endurance training enhances ABCA1 expression in rat small intestine . *Eur J Appl Physiol* . 2009 ; 107 : 351 - 358
- 10 . Rachmilewitz D , Fainaru M . ApolipoproteinA-I synthesis and secretion by cultured human intestinal mucosa . *Metab Clin Exp* . 1979 ; 28 : 739 - 743
- 11 . Johansson C , Rössner S , Wallidius G . HDL secretion from human intestinal tract . *Lancet* . 1978 ; 8059 : 324 - 325
- 12 . Forman B , Chen J , Evans R . Hypolipidemic drugs, polyunsaturated fatty-acids, and eicosanoids are ligands for PPAR-alpha and delta . *Proc Natl Acad Sci US A* . 1997 ; 94 : 4312 - 4317
- 13 . Lefebvre P , Chinetti G , Fruchart J , Staels B . Sorting out the roles of PPAR-alpha in energy metabolism and vascular homeostasis . *J Clin Invest* . 2006 ; 116 : 571 - 580
- 14 . Braissant O , Foufelle F , Scotto C , Dauça M , Wahli W . Differential expression of peroxisome proliferator-activated receptors: tissue distribution of PPAR-alpha, -beta, and -gamma in the adult rat . *Endocrinology* . 1996 ; 137 : 354 - 366
- 15 . Chinetti G , Lestavel S , Bocher V , Remaley A , Neve B , Pineda Torra I , Teissier E , Minnich A , Jaye M , Duverger N , Brewer H , Fruchart J , Clavey V , Staels B . PPAR-alpha and PPAR-gamma activators induce cholesterol removal from human macrophage foam cells through stimulation of the ABCA1 pathway . *Nat Med* . 2001 ; 7 : 53 - 58
- 16 . Colin S , Bourguignon E , Boullay A , Tousaint J , Huet S , Caira F , Staels B , Lestavel S , Lobaccaro J , Delerive P . Intestine-specific regulation of PPAR-alpha gene transcription by LXR . *Endocrinology* . 2008 ; 149 : 5128 - 5135
- 17 . Watts G , Barrett P , Ji J , Serone A , Chan D , Croft K , Loehrer F , Johnson A . Differential regulation of lipoprotein kinetics by atorvastatin and fenofibrate in subjects with the metabolic syndrome . *Diabetes* . 2003 ; 52 : 803 - 811
- 18 . Berthou L , Duverger N , Emmanuel F , Langouët S , Auwerx J , Guillouzo A , Fruchart J , Rubin E , Denèfle P , Staels B , Branellec D . Opposite regulation of human versus mouse apolipoproteinA-I by fibrates in human apolipoproteinA-I transgenic mice . *J Clin Invest* . 1996 ; 97 : 2408 - 2416
- 19 . Hossain M , Tsujita M , Gonzalez F , Yokoyama S . Effects of fibrate drugs on expression of ABCA1 and HDL biogenesis in hepatocytes . *J Cardiovasc Pharmacol* . 2008 ; 51 : 258 - 266
- 20 . Chinetti G , Lestavel S , Fruchart J , Clavey V , Staels B . PPAR-alpha reduces cholesterol esterification in macrophages . *Circ Res* . 2003 ; 92 : 212 - 217
- 21 . Chakravarthy M , Lodhi I , Yin L , Malapaka R , Xu H , Turk J , Semenkovich C . Identification of a physiologically relevant endogenous ligand for PPAR-alpha in liver . *Cell* . 2009 ; 138 : (3) 476 - 488
- 22 . Delerive P , Furman C , Teissier E , Fruchart J , Duriez P , Staels B . Oxidized phospholipids activate PPAR-alpha in a phospholipase A2-dependent manner . *FEBS Lett* . 2000 ; 471 : 34 - 38
- 23 . Staels B , van Tol A , Andreu T , Auwerx J . Fibrates influence the expression of genes involved in lipoprotein metabolism in a tissue-selective manner in the rat . *Arterioscler Thromb* . 1992 ; 12 : 286 - 294
- 24 . Bünger M , van den Bosch H , van der Meijde J , Kersten S , Hooiveld G , Müller M . Genome-wide analysis of PPAR-alpha activation in murine small intestine . *Physiol Genomics* . 2007 ; 30 : (2) 192 - 204
- 25 . Cariou B , Zaïr Y , Staels B , Bruckert E . Effects of the new dual PPAR-alpha/delta agonist GFT505 on lipid and glucose homeostasis in abdominally obese patients with combined dyslipidemia or impaired glucose metabolism . *Diabetes Care* . 2011 ; 34 : 2008 - 2014
- 26 . Lalloyer F , Wouters K , Baron M , Caron S , Vallez E , Vanhoutte J , Baugé E , Shiri-Sverdlov R , Hofker M , Staels B , Tailleux A . PPAR-alpha gene level differently affects lipid metabolism and inflammation in apolipoprotein-E2 knock-in mice . *Arterioscler Thromb Vasc Biol* . 2011 ; 31 : 1573 - 1579
- 27 . Chateau D , Pauqui T , Delers F , Rousset M , Chambaz J , Demignot S . Lipid micelles stimulate the secretion of triglyceride-enriched apolipoprotein-B48-containing lipoproteins by Caco-2 cells . *J Cell Physiol* . 2005 ; 202 : 767 - 776
- 28 . Béaslas O , Cueille C , Delers F , Chateau D , Chambaz J , Rousset M , Carrière V . Sensing of dietary lipids by enterocytes: a new role for SR-BI/c1a-1 . *PLoS ONE* . 2009 ; 4 : e4271 - 4278
- 29 . Rasband W . ImageJ . US National Institutes of Health ; Bethesda, Maryland, USA <http://rsb.info.nih.gov/ij/> 1997-2009 ;
- 30 . Hennuyer N , Tailleux A , Torpier G , Mezdoor H , Fruchart J , Staels B , Fiévet C . PPAR-alpha, but not PPAR-gamma, activators decrease macrophage-laden atherosclerotic lesions in a nondiabetic mouse model of mixed dyslipidemia . *Arterioscler Thromb Vasc Biol* . 2005 ; 25 : 1897 - 1902
- 31 . Duval C , Touche V , Tailleux A , Fruchart J , Fievet C , Clavey V , Staels B , Lestavel S . NPC1L1 gene expression is down regulated by LXR activators in the intestine . *BBRC* . 2006 ; 340 : 1259 - 1263
- 32 . Iqbal J , Hussain M . Intestinal lipid absorption . *Am J Physiol Endocrinol Metab* . 2009 ; 296 : 1183 - 1194

- 33 . Kimura R , Takahashi N , Murota K , Yamada Y , Niiya S , Kanzaki N , Murakami Y , Moriyama T , Goto T , Kawada T . Activation of PPAR- α suppresses postprandial lipidemia through fatty-acid-oxidation in enterocytes . *Biochem Biophys Res Commun* . 2011 ; 410 : 1 - 6
- 34 . Field F , Watt K , Mathur S . Ezetimibe interferes with cholesterol trafficking from the plasma membrane to the endoplasmic reticulum in Caco-2 cells . *J Lipid Res* . 2007 ; 48 : 1735 - 1745
- 35 . Lee B , Zhu J , Wolins N , Cheng J , Buhman K . Differential association of adipophilin and TIP47 proteins with cytoplasmic lipid droplets in mouse enterocytes during dietary fat absorption . *Biochim Biophys Acta* . 2009 ; 1791 : (12) 1173 - 1180
- 36 . Lee J , Giannikopoulos P , Duncan S , Wang J , Johansen C , Brown J , Plutzky J , Hegele R , Glimcher L , Lee A . The transcription factor cyclic amp-responsive element-binding protein H regulates triglyceride metabolism . *Nat Med* . 2011 ; 17 : (7) 812 - 815
- 37 . van der Velde A , Brufau G , Groen A . Transintestinal cholesterol efflux . *Curr Opin Lipidol* . 2010 ; 21 : 167 - 171
- 38 . Weingärtner O , Lütjohann D , Böhm M , Laufs U . Relationship between cholesterol synthesis and intestinal absorption is associated with cardiovascular risk . *Atherosclerosis* . 2010 ; 210 : 362 - 365
- 39 . Xiao C , Lewis G . Regulation of chylomicron production in humans . *Biochim Biophys Acta* . 2011 ; 10.1016/j.bbaliip.2011.09.019
- 40 . Fruchart J , Duriez P . Mode of action of fibrates in the regulation of triglyceride and HDL-cholesterol metabolism . *Drugs Today* . 2006 ; 42 : 39 - 64
- 41 . Rigamonti E , Chinetti-Gbaguidi G , Staels B . Regulation of macrophage functions by PPAR-alpha, PPAR-gamma, and LXRs in mice and men . *Arterioscler Thromb Vasc Biol* . 2008 ; 28 : 1050 - 1059
- 42 . Keech A , Simes R , Barter P , Best J , Scott R , Taskinen M , Forder P , Pillai A , Davis T , Glasziou P , Drury P , Kesäniemi Y , Sullivan D , Hunt D , Colman P , d' Emden M , Whiting M , Ehnholm C , Laakso M , FIELD SI . Effects of long-term fenofibrate therapy on cardiovascular events in 9795 people with type 2 diabetes mellitus (the field study): randomised controlled trial . *Lancet* . 2005 ; 366 : 1849 - 1861
- 43 . Tenenbaum A , Motro M , Fisman E , Tanne D , Boyko V , Behar S . Bezafibrate for the secondary prevention of myocardial infarction in patients with metabolic syndrome . *Arch Intern Med* . 2005 ; 165 : 1154 - 1160
- 44 . The BIP Study Group . Secondary prevention by raising HDL-cholesterol and reducing triglycerides in patients with coronary artery disease: the bezafibrate infarction prevention study . *Circulation* . 2000 ; 102 : 21 - 27
- 45 . Dullens S , Mensink R , Mariman E , Plat J . Differentiated Caco-2 cells as an in-vitro model to evaluate de-novo apolipoproteinA-I production in the small intestine . *Eur J Gastroenterol Hepatol* . 2009 ; 21 : 642 - 649
- 46 . Chinetti-Gbaguidi G , Rigamonti E , Helin L , Mutka A , Lepore M , Fruchart J , Clavey V , Ikonen E , Lestavel S , Staels B . PPAR-alpha controls cellular cholesterol trafficking in macrophages . *J Lipid Res* . 2005 ; 46 : 2717 - 2725
- 47 . Pramfalk C , Angelin B , Eriksson M , Parini P . Cholesterol regulates ACAT2 gene expression and enzyme activity in human hepatoma cells . *Biochem Biophys Res Commun* . 2007 ; 364 : 402 - 409
- 48 . Granneman J , Moore H , Krishnamoorthy R , Rathod M . Perilipin controls lipolysis by regulating the interactions of AB-hydrolase containing 5 and adipose triglyceride lipase . *J Biol Chem* . 2009 ; 284 : 34538 - 34544
- 49 . Lo V , Erickson B , Thomason-Hughes M , Ko K , Dolinsky V , Nelson R , Lehner R . Arylacetamide deacetylase attenuates fatty-acid-induced triacylglycerol accumulation in rat hepatoma cells . *J Lipid Res* . 2010 ; 51 : 368 - 377
- 50 . Zhou Y , Abidi P , Kim A , Chen W , Huang T , Kraemer F , Liu J . Transcriptional activation of hepatic ACSL3 and ACSL5 by oncostatin m reduces hypertriglyceridemia through enhanced beta-oxidation . *Arterioscler Thromb Vasc Biol* . 2007 ; 27 : 2198 - 2205

Figure 1

PPAR α -activation with GFT505 *in-vivo* in mice and GFT505 and GW7647 *ex-vivo* in human jejunal biopsies increases the expression of genes involved in HDL production

Q-PCR analysis of ABCA1 (A, C, E) and apoAI (B, D, F) was performed on RNA isolated from scrapped intestinal mucosa (A, B, C, D) of GFT505 (30mpk) or FF (200mpk)-treated apoE2-KI mice for 14d and from human jejunal biopsies (E, F) treated with GW7647 (600nM) and GFT505 (1 μ M) for 18h. *In vivo* values are expressed as means \pm SEM and *ex vivo* values as means \pm SD. t test; *, P<0.05; ***,P<0.001 Vehicle vs treatment).




Figure 2

GW7647 increases the production of HDL by Caco-2/TC7

Caco-2/TC7 (non-infected, ShControl- or ShPPAR α -infected) were treated with GW7647 (600nM) for 48h (Q-PCR and ELISA analysis) or 72h (cholesterol efflux assay). (A) Number of HDL secreted by non-infected Caco-2/TC7 per equivalent volume. (B) ELISA analysis of apoAI in the HDL fraction (representative of 3 independent experiments, values are expressed as means \pm SEM). (C) Cholesterol-loaded primary human macrophages were incubated with HDL secreted by Caco-2/TC7 (representative of 2 independent experiments, values are expressed as means \pm SEM). (D) Q-PCR analysis of ABCA1. Values are expressed as means \pm SD. t-test; * P <.05; ** P <.01; *** P <.001 Vehicle vs treatment; # P <.05 treated ShControl- vs treated-ShPPAR α -infected cells.





Figure 3

PPAR α -activation modulates intracellular cholesterol trafficking, increases cytoplasmic lipid droplet formation and reduces apoB secretion by Caco-2/TC7

Caco-2/TC7 were incubated with NBD-cholesterol-micelles (50 μ g/well) for 4h and then for 4h with non-fluorescent micelles. GW7647 (600nM) was added 24h before and during the NBD-cholesterol-loading period (representative of 3 independent experiments). (A) NBD-fluorescence was quantified using Image J at each optical slice. (B) NBD-cholesterol (green), ZO-1 (red) and Hoechst (blue) were visualized by confocal microscopy. (C) ApoB in the basolateral medium was quantified by ELISA. Values are expressed as means \pm SEM. t-test; *P<.05 Vehicle vs treatment.

**Figure 4**

PPAR α -activation decreases cholesterol esterification in Caco-2/TC7

Caco-2/TC7 were incubated with [3 H]-cholesterol-micelles (7.5 μ Ci/well) for 48h. GW7647 (600nM) was added 24h before and during the [3 H]-cholesterol-loading. Intracellular lipids (A) and lipids secreted in the basolateral compartment (B) were extracted and separated by TLC. Results are expressed as means of CE percent of total cholesterol (\pm SEM). Q-PCR analysis of ACAT-2 (C) was performed on GW7647 (600nM)-treated Caco-2/TC7 for 24h. Values are expressed as means \pm SD. t-test; *P<.05; **P<.01 Vehicle vs treatment.




Figure 5

PPAR α -activation increases cytoplasmic lipid droplet formation-associated genes in Caco-2/TC7 and in human jejunal biopsies

Q-PCR analysis of PLIN2 (A, C) and ABHD5 (B, D) was performed on RNA isolated from Caco-2/TC7 (A, B) and human jejunal biopsies (C, D) treated with GW7647 (600nM) or GFT505 (1 μ M) for 24h or 18h respectively. Values are expressed as means \pm SD. t test; *P<.05; **P<.01; ***P<.001 Vehicle vs treatment.




Figure 6

PPAR α -activation increases fatty-acid oxidation genes in Caco-2/TC7 and in human jejunal biopsies

Q-PCR analysis of CPT-1 (A, D), ACOX1 (B, E) and ACSL5 (C) was performed on RNA isolated from Caco-2/TC7 (A, B, C) or human jejunal biopsies (D, E) treated with GW7647 (600nM) or GFT505 (1 μ M) for 24h or 18h respectively. Values are expressed as means \pm SD. t-test; *P<.05; **P<.01; ***P<.001 Vehicle vs treatment.




Table 1

Lipid profiles of apoE2-KI mice treated with FF or GFT505

ApoE2-KI mice on a western-diet were treated by oral gavage with FF (200mpk) or GFT505 (10mpk) or vehicle for 14d (n=6 mice/group).

	Vehicle	FF	GFT505
TG (mg/dL)	278 ± 76	156 ± 62 [*]	110 ± 50 [*]
Total cholesterol (mg/dL)	1006 ± 190	459 ± 201 ^{**}	384 ± 142 ^{***}
HDL-cholesterol (mg/dL)	26 ± 7	62 ± 11 ^{***}	71 ± 10 ^{***, #}
non HDL-cholesterol (mg/dL)	980 ± 196	397 ± 203 ^{**}	312 ± 142 ^{***}

Values are expressed as means ± SEM. ANOVA;

^{*} P<.05;^{**} P<.01;^{***} P<.001 vs untreated mice;[#] P<.05 vs FF-treated mice.