

A central role for Galectin 3 during renal epithelial cell morphogenesis after nephrectomy

Françoise Poirier, Amandine Viau, Jérémy Magescas, Martine Burtin, Fabiola Terzi, Delphine Delacour

► **To cite this version:**

Françoise Poirier, Amandine Viau, Jérémy Magescas, Martine Burtin, Fabiola Terzi, et al.. A central role for Galectin 3 during renal epithelial cell morphogenesis after nephrectomy. First International Cilia in Development and Disease Scientific Conference, 1 (Suppl 1), pp.P76, 2012. <inserm-00752967>

HAL Id: inserm-00752967

<http://www.hal.inserm.fr/inserm-00752967>

Submitted on 16 Nov 2012

HAL is a multi-disciplinary open access archive for the deposit and dissemination of scientific research documents, whether they are published or not. The documents may come from teaching and research institutions in France or abroad, or from public or private research centers.

L'archive ouverte pluridisciplinaire **HAL**, est destinée au dépôt et à la diffusion de documents scientifiques de niveau recherche, publiés ou non, émanant des établissements d'enseignement et de recherche français ou étrangers, des laboratoires publics ou privés.

POSTER PRESENTATION

Open Access

A central role for Galectin 3 during renal epithelial cell morphogenesis after nephrectomy

F Poirier^{1*}, A Viau², J Magescas¹, M Burtin², F Terzi², D Delacour¹

From First International Cilia in Development and Disease Scientific Conference (2012)
London, UK. 16-18 May 2012

Galectin-3, a member of the multigene family of beta-galactoside binding lectins, is mainly expressed in epithelial cells. The distribution of galectin-3 depends on cell type and cell differentiation. We previously showed that galectin-3 is transiently associated with the centrosome, and more specifically with the basal body at the basis of the primary cilium of MDCK cells. In the present study, we use *galectin-3* null (*Gal3*^{-/-}) mutant mice to investigate the functional consequences of our observations in the context of kidney regeneration after subtotal nephrectomy. Three months after surgery, renal functions were more severely affected in *Gal3*^{-/-} mutant than in wt animals. Upon sacrifice, we found that *Gal3*^{-/-} mutant kidneys were bigger and heavier, and this hypertrophy was associated with large tubular dilatations. Interestingly, in wt mice, galectin 3 intracellular distribution was altered in response to injury, displaying a distinct centrosomal localization in epithelial cells lining the collecting ducts close to the wound, whereas in cells away from the wound galectin-3 remained uniformly cytosolic. The impact of nephron reduction on primary cilium growth was examined by 3D reconstruction of wt and *Gal3*^{-/-} renal tubules stained for acetylated alpha-tubulin. This analysis revealed that mutant primary cilia are irregular, bent, twisted and longer than wt primary cilia, indicating that Galectin-3 is essential for primary cilium growth. This suggests that the failure in kidney regeneration observed in *Gal3*^{-/-} mutant mice may be due to defects in primary cilium biogenesis.

Author details

¹Institut Jacques Monod CNRS Paris Diderot, France. ²INSERM U845, France.

Published: 16 November 2012

* Correspondence: poirier@ijm.univ-paris-diderot.fr

¹Institut Jacques Monod CNRS Paris Diderot, France

Full list of author information is available at the end of the article

doi:10.1186/2046-2530-1-S1-P76

Cite this article as: Poirier et al.: A central role for Galectin 3 during renal epithelial cell morphogenesis after nephrectomy. *Cilia* 2012 1 (Suppl 1):P76.

Submit your next manuscript to BioMed Central and take full advantage of:

- Convenient online submission
- Thorough peer review
- No space constraints or color figure charges
- Immediate publication on acceptance
- Inclusion in PubMed, CAS, Scopus and Google Scholar
- Research which is freely available for redistribution

Submit your manuscript at
www.biomedcentral.com/submit

