

Delineation of CCDC39/CCDC40 mutation spectrum and associated phenotypes in primary ciliary dyskinesia

Marie Legendre, Sylvain Blanchon, Bruno Copin, Philippe Duquesnoy, Guy Montantin, Esther Kott, Florence Dastot, Ludovic Jeanson, Marine Cachanado, A Rousseau, et al.

► **To cite this version:**

Marie Legendre, Sylvain Blanchon, Bruno Copin, Philippe Duquesnoy, Guy Montantin, et al.. Delineation of CCDC39/CCDC40 mutation spectrum and associated phenotypes in primary ciliary dyskinesia. First International Cilia in Development and Disease Scientific Conference, May 2012, Londres, United Kingdom. BMC, 1 (Suppl 1), pp.P91, 2012. <inserm-00752969>

HAL Id: inserm-00752969

<http://www.hal.inserm.fr/inserm-00752969>

Submitted on 16 Nov 2012

HAL is a multi-disciplinary open access archive for the deposit and dissemination of scientific research documents, whether they are published or not. The documents may come from teaching and research institutions in France or abroad, or from public or private research centers.

L'archive ouverte pluridisciplinaire **HAL**, est destinée au dépôt et à la diffusion de documents scientifiques de niveau recherche, publiés ou non, émanant des établissements d'enseignement et de recherche français ou étrangers, des laboratoires publics ou privés.

POSTER PRESENTATION

Open Access

Delineation of *CCDC39/CCDC40* mutation spectrum and associated phenotypes in primary ciliary dyskinesia

M Legendre^{1*}, S Blanchon^{1,2}, B Copin¹, P Duquesnoy¹, G Montantin¹, E Kott¹, F Dastot¹, L Jeanson¹, M Cachanado³, A Rousseau³, JF Papon⁴, A Tamalet², AM Vojtek⁵, D Escalier¹, A Coste⁴, J de Blic⁶, A Clément², E Escudier¹, S Amselem¹

From First International Cilia in Development and Disease Scientific Conference (2012) London, UK. 16-18 May 2012

Background

CCDC39 and *CCDC40* genes have recently been implicated in primary ciliary dyskinesia (PCD) with inner dynein arms (IDA) defects and axonemal disorganization; their contribution to the disease is, however, unknown. With the aim to delineate *CCDC39/CCDC40* mutation spectrum and associated phenotypes, we screened a large cohort of patients with IDA defects, and accurately described their clinical and ciliary phenotypes.

Methods

All *CCDC39* and *CCDC40* exons and intronic boundaries were sequenced in 43 patients from 40 unrelated families. We recorded and compared clinical features (sex, origin, consanguinity, laterality defects, ages at first symptoms and evaluation, neonatal respiratory distress, airway infections, nasal polyposis, otitis media, bronchiectasis, infertility), ciliary beat frequency and quantitative ultrastructural analyses of cilia and sperm flagella.

Results

Biallelic *CCDC39* or *CCDC40* mutations were identified in 30/34 (88.2%) unrelated families with IDA defects and axonemal disorganization (22 and 8 families, respectively). Fourteen of the 28 identified mutations are novel. No mutation was found in the 6 families with isolated IDA defects. Patients with identified mutations shared a similar phenotype, in terms of both clinical features and ciliary

structure and function. The sperm flagellar ultrastructure, analyzed in 4/7 infertile males, evidenced abnormalities similar to the ciliary ones.

Conclusions

CCDC39 and *CCDC40* mutations represent the major cause of PCD with IDA defects and axonemal disorganization. Patients carrying *CCDC39* or *CCDC40* mutations are phenotypically indistinguishable. *CCDC39* and *CCDC40* analyses in selected patients ensure to find mutations with high probability, even if clinical or ciliary phenotypes cannot prioritize one analysis over the other.

Author details

¹INSERM, UMR_S933, UPMC Univ Paris 06; and AP-HP, Hôpital Armand-Trousseau, Service de Génétique et d'Embryologie Médicales, F-75012, Paris, France. ²AP-HP, Hôpital Armand-Trousseau, Unité de Pneumologie Pédiatrique, Centre National de Référence des Maladies Respiratoires Rares, F-75012, Paris, France. ³AP-HP, Hôpital Saint-Antoine, Unité de Recherche Clinique; and UPMC Univ Paris 06, Unité Fonctionnelle de Pharmacologie, F-75012, Paris, France. ⁴AP-HP, Hôpital Inter-Communal et Groupe Hospitalier Henri Mondor-Albert Chenevier, Service d'ORL et de Chirurgie Cervico-Faciale, F-94000, Créteil, France. ⁵Hôpital Inter-Communal, Service d'Anatomo-Pathologie, F-94000, Créteil, France. ⁶AP-HP, Groupe Hospitalier Necker-Enfants Malades, Service de Pneumologie et Allergologie Pédiatriques, F-75015, Paris, France.

Published: 16 November 2012

doi:10.1186/2046-2530-1-S1-P91

Cite this article as: Legendre *et al.*: Delineation of *CCDC39/CCDC40* mutation spectrum and associated phenotypes in primary ciliary dyskinesia. *Cilia* 2012 **1**(Suppl 1):P91.

* Correspondence: marie.legendre@trs.aphp.fr

¹INSERM, UMR_S933, UPMC Univ Paris 06; and AP-HP, Hôpital Armand-Trousseau, Service de Génétique et d'Embryologie Médicales, F-75012, Paris, France

Full list of author information is available at the end of the article