

How do T-type calcium channels control low-threshold exocytosis?

Norbert Weiss, Gerald Zamponi, Michel De Waard

► **To cite this version:**

Norbert Weiss, Gerald Zamponi, Michel De Waard. How do T-type calcium channels control low-threshold exocytosis?. *Communicative and Integrative Biology*, Taylor

Francis Open, 2012, 5 (4), pp.377-80. <10.4161/cib.19997>. <inserm-00759004>

HAL Id: inserm-00759004

<http://www.hal.inserm.fr/inserm-00759004>

Submitted on 29 Nov 2012

HAL is a multi-disciplinary open access archive for the deposit and dissemination of scientific research documents, whether they are published or not. The documents may come from teaching and research institutions in France or abroad, or from public or private research centers.

L'archive ouverte pluridisciplinaire **HAL**, est destinée au dépôt et à la diffusion de documents scientifiques de niveau recherche, publiés ou non, émanant des établissements d'enseignement et de recherche français ou étrangers, des laboratoires publics ou privés.

How do T-type calcium channels control low-threshold exocytosis?

Norbert Weiss,^{1,*} Gerald W. Zamponi¹ and Michel De Waard^{2,3}

¹Hotchkiss Brain Institute; Department of Physiology and Pharmacology; University of Calgary; Calgary, AB Canada; ²Grenoble Institute of Neuroscience; La Tronche, France; ³University Joseph Fourier; Grenoble, France

Low-voltage-activated T-type calcium channels act as a major pathway for calcium entry near the resting membrane potential in a wide range of neuronal cell types. Several reports have uncovered an unrecognized feature of T-type channels in the control of vesicular neurotransmitter and hormone release, a process so far thought to be mediated exclusively by high-voltage-activated calcium channels. However, the underlying molecular mechanisms linking T-type calcium channels to vesicular exocytosis have remained enigmatic. In a recent study, we have reported that Ca_v3.2 T-type channel forms a signaling complex with the neuronal Q-SNARE syntaxin-1A and SNAP-25. This interaction that relies on specific Ca_v3.2 molecular determinants, not only modulates T-type channel activity, but was also found essential to support low-threshold exocytosis upon Ca_v3.2 channel expression in MPC 9/3L-AH chromaffin cells. Overall, we have indentified an unrecognized regulation pathway of T-type calcium channels by SNARE proteins, and proposed the first molecular mechanism by which T-type channels could mediate low-threshold exocytosis.

Depolarization-evoked synaptic transmission relies on the calcium (Ca²⁺)-regulated release of quantal packets of neurotransmitters following fusion of synaptic vesicles with the presynaptic plasma membrane.¹ It is well established that neuronal voltage-gated Ca²⁺ channels, by converting electrical signals into intracellular Ca²⁺ concentration elevations, play a key role in triggering evoked neurotransmitter

release.^{2–4} Hence, Ca²⁺ entry through high-voltage-activated (HVA) channels (N-, P/Q-, and in some extent and particular cell populations, L- and R-type) into presynaptic nerve terminals in response to action potentials supports a transient Ca²⁺ microdomain⁵ essential for synaptic exocytosis. However, the observation that some neurons can release functionally significant amounts of neurotransmitter below the threshold of action potentials⁶ questioned the possible involvement of another source of Ca²⁺ ions, independent of HVA channels activation.

In contrast to HVA channels, low-voltage-activated (LVA) T-type Ca²⁺ channels activate in response to subthreshold membrane depolarizations between -65 mV and -50 mV and thus represent an important source of Ca²⁺ entry near the resting membrane potential. Hence, besides controlling important physiological processes by regulating neuronal excitability, pacemaker activity and post-inhibitory rebound burst firing, mounting evidences from various neuronal cell types suggest an efficient role of T-type channels in fast and low-threshold exocytosis.^{7–10} Until now, however, the mechanism whereby these channels support exocytosis events at the molecular level remained a mystery. In a recent study, we provided compelling evidence of the existence of a Ca_v3.2/syntaxin-1A molecular complex essential for T-type-dependent exocytosis.¹¹

In mammalian synapses, interaction of several members of the vesicle-docking / release machinery (including syntaxin-1A/1B and SNAP-25) onto a *synprint* (*synaptic protein interaction site*) domain

Keywords: calcium channel, T-type, Ca_v3.2 channel, SNARE, syntaxin-1A, SNAP-25, exocytosis, neuron, chromaffin cell, MPC cell

Submitted: 03/11/12

Accepted: 03/12/12

<http://dx.doi.org/10.4161/cib.19997>

*Correspondence to: Norbert Weiss;
Email: naweiss@ucalgary.ca
or norbert.weiss@yahoo.fr

Addendum to: Weiss N, Hameed S, Fernández-Fernández JM, Fablet K, Karmazinova M, Poillot C, et al. A Ca_v3.2/syntaxin-1A signaling complex controls T-type channel activity and low-threshold exocytosis. *J Biol Chem* 2012; 287:2810–8; PMID:22130660; <http://dx.doi.org/10.1074/jbc.M111.290882>

located within the intracellular loop between domains II and III of the $Ca_v2.1$ ¹² and $Ca_v2.2$ ¹³ channels (Fig. 1A, top panel) ensures a close localization of the secretory vesicles near the Ca^{2+} source. In turn, both syntaxin-1A/1B and SNAP-25 modulate calcium channel activity^{14,15} to fine tune Ca^{2+} entry and synaptic strength. However, in contrast to HVA channels, all of the three T-type channel members (i.e. $Ca_v3.1$, $Ca_v3.2$ and $Ca_v3.3$) lack the consensus *synprint* site, making the molecular understanding of the involvement of these channels in the exocytosis process quite difficult. Our observation that $Ca_v3.2$ channels associate with syntaxin-1A in central neurons prompted us to investigate the possible existence of specific $Ca_v3.2$ channel molecular determinants other than the consensus HVA *synprint* domain. Using biochemical and cellular trafficking approaches, we demonstrated that syntaxin-1A, as well as SNAP-25, interact with the C-terminal domain of $Ca_v3.2$ channel (Fig. 1A, lower panel). Moreover, using patch-clamp recordings performed on tsA-201 cells expressing $Ca_v3.2$ channels, we demonstrated that co-expression of a syntaxin-1A in its “closed” conformational state (i.e. the conformation adopted by the syntaxin-1A in isolation or in interaction with Munc18)^{16,17} potentially decreases $Ca_v3.2$ channel availability by shifting the voltage-dependence of inactivation toward more hyperpolarized membrane potentials, similarly to what was previously reported for N- and P/Q-type channels.^{14,15,18-20} Interestingly, this regulation was abolished upon co-expression of SNAP-25, and not observed with a constitutively “open” syntaxin-1A (i.e. the conformation adopted upon its association with SNAP-25)¹⁸ (Fig. 1B). Given that syntaxin-1A undergoes a conformational switch from a “closed” to an “open” conformation during the vesicle release cycle,^{16,21,22} this suggests that syntaxin-1A may be able to dynamically regulate T-type channel availability during various stages of exocytosis. Interestingly, although T-type channels utilize distinct molecular determinants to interact with SNARE proteins (the C-terminal domain vs. the classical *synprint* of the II-III linker), they are subjected to a similar SNARE

regulation. Does this observation question the molecular mechanism by which binding of syntaxin-1A produces changes in channel gating? Earlier reports have shown

that reorganization of intramolecular interactions among the main intracellular loops of Ca_v2 channels critically influence channel inactivation.²³⁻³⁰ Mapping the




Figure 1. SNARE proteins modulate high- and low-voltage-gated calcium channels via distinct molecular determinants. (A) Membrane topology of voltage-gated calcium channels highlighting the localization of the *synprint* site located within the intracellular linker between domains II and III of $Ca_v2.1/Ca_v2.2$ channels (top panel), and the “*synprint like*” domain of $Ca_v3.x$ channels (bottom panel) located within the C-terminal domain of the channel. (B) Voltage-dependence of $Ca_v3.x$ channel availability during the conformational switch of syntaxin-1A.

intramolecular interactions of T-type channels along with the characterization of the minimal sequence engaged in the interaction with SNARE proteins will provide important structural information on how syntaxin-1A modulates channel gating.

It is well known that direct interaction of SNARE proteins with Ca_v2.1 and Ca_v2.2 channels is critical for depolarization-evoked neurotransmitter release. Hence, disruption of the Ca²⁺ channel-SNARE proteins coupling by deletion of the *synprint* domain or by peptides derived from the *synprint* sequence, alters synaptic transmission.³¹⁻³⁴ We revealed that similarly to HVA channels, T-type channels-mediated exocytosis relies on a channel-SNARE protein interaction. Indeed, membrane capacitance recordings performed on MPC 9/3L-HA chromaffin cells expressing Ca_v3.2 channels revealed robust voltage-dependent exocytosis which was totally prevented by co-expression of the Ca_v3.2 C-terminal domain (i.e the synaptic protein interaction site of Ca_v3.2). Ablation of Ca_v3.2-dependent exocytosis most likely results from the specific uncoupling of the channel with SNARE proteins and not from a side

alteration of the exocytosis machinery by itself because no alteration was observed when exocytosis was induced by direct intracellular Ca²⁺ elevation. Hence, we showed that similarly to HVA channels, a physical coupling between SNARE proteins and T-type channels is critical for T-type-dependent exocytosis. Considering the relative small conductance of T-type channels³⁵ and the restricted diffusion of Ca²⁺ due to the high Ca²⁺ buffering capacity of neuronal cells,³⁶ it is conceivable that this interaction allows the close localization of the vesicle-docking / release machinery in close proximity to the Ca²⁺ source in order to efficiently sense Ca²⁺ elevation. However, we cannot exclude the possibility that interaction of T-type channels with SNARE proteins could form a macromolecular complex through which channel conformational changes following membrane depolarization would work as an on/off molecular switch of secretion by controlling the ultimate conformational change of the releasing complex as previously proposed for HVA channels.^{37,38} Although this concept still requires further investigation, the use on a non-conducting channel to investigate T-type-dependent secretion would definitively provide

interesting information about the functional importance of T-type channel interaction with SNARE proteins.

Overall, we revealed an unrecognized regulation of low-voltage-activated T-type Ca²⁺ channels by SNARE proteins, and provide the first evidence for a molecular mechanism by which these channels could mediate low-threshold exocytosis. We revealed that, although T-type Ca²⁺ channels differ from HVA channels by their molecular constituents, they possess the same ability to functionally interact with SNARE proteins, highlighting a key evolutionary mechanism for specialized fast and spatially delimited exocytosis.

Disclosure of Potential Conflicts of Interest

No potential conflicts of interest were disclosed.

Acknowledgments

N.W is supported by a postdoctoral fellowship from Alberta Innovates Health Solutions (AIHS) and Hotchkiss Brain Institute. G.W.Z is funded by the Canadian Institutes of Health Research, is a Canada Research Chair and AIHS Scientist.

References

- Edwards RH. The neurotransmitter cycle and quantal size. *Neuron* 2007; 55:835-58; PMID:17880890; <http://dx.doi.org/10.1016/j.neuron.2007.09.001>
- Linás R, Sugimori M, Silver RB. Microdomains of high calcium concentration in a presynaptic terminal. *Science* 1992; 256:677-9; PMID:1350109; <http://dx.doi.org/10.1126/science.1350109>
- Neher E, Sakaba T. Multiple roles of calcium ions in the regulation of neurotransmitter release. *Neuron* 2008; 59:861-72; PMID:18817727; <http://dx.doi.org/10.1016/j.neuron.2008.08.019>
- Weber AM, Wong FK, Tufford AR, Schlichter LC, Matveev V, Stanley EF. N-type Ca₂₊ channels carry the largest current: implications for nanodomains and transmitter release. *Nat Neurosci* 2010; 13:1348-50; PMID:20953196; <http://dx.doi.org/10.1038/nn.2657>
- Schneggenburger R, Neher E. Presynaptic calcium and control of vesicle fusion. *Curr Opin Neurobiol* 2005; 15:266-74; PMID:15919191; <http://dx.doi.org/10.1016/j.conb.2005.05.006>
- Ivanov AI, Calabrese RL. Intracellular Ca₂₊ dynamics during spontaneous and evoked activity of leech heart interneurons: low-threshold Ca currents and graded synaptic transmission. *J Neurosci* 2000; 20:4930-43; PMID:10864951
- Carabelli V, Marcantoni A, Comunanza V, de Luca A, Díaz J, Borges R, et al. Chronic hypoxia up-regulates alpha1H T-type channels and low-threshold catecholamine secretion in rat chromaffin cells. *J Physiol* 2007; 584:149-65; PMID:17690152; <http://dx.doi.org/10.1113/jphysiol.2007.132274>
- Pan ZH, Hu HJ, Perring P, Andrade R. T-type Ca²⁺ channels mediate neurotransmitter release in retinal bipolar cells. *Neuron* 2001; 32:89-98; PMID:11604141; [http://dx.doi.org/10.1016/S0896-6273\(01\)00454-8](http://dx.doi.org/10.1016/S0896-6273(01)00454-8)
- Egger V, Svoboda K, Mainen ZF. Mechanisms of lateral inhibition in the olfactory bulb: efficiency and modulation of spike-evoked calcium influx into granule cells. *J Neurosci* 2003; 23:7551-8; PMID:12930793
- Tang AH, Karson MA, Nagode DA, McIntosh JM, Uebele VN, Renger JJ, et al. Nerve terminal nicotinic acetylcholine receptors initiate quantal GABA release from perisomatic interneurons by activating axonal T-type (Cav3) Ca²⁺ channels and Ca²⁺ release from stores. *J Neurosci* 2011; 31:13546-61; PMID:21940446; <http://dx.doi.org/10.1523/JNEUROSCI.2781-11.2011>
- Weiss N, Hameed S, Fernández-Fernández JM, Fablet K, Karmazinova M, Poillot C, et al. A Ca_v3.2/syntaxin-1A signaling complex controls T-type channel activity and low-threshold exocytosis. *J Biol Chem* 2012; 287:2810-8; PMID:22130660; <http://dx.doi.org/10.1074/jbc.M111.290882>
- Rettig J, Sheng ZH, Kim DK, Hodson CD, Snutch TP, Catterall WA. Isoform-specific interaction of the alpha1A subunits of brain Ca²⁺ channels with the presynaptic proteins syntaxin and SNAP-25. *Proc Natl Acad Sci U S A* 1996; 93:7363-8; PMID:8692999; <http://dx.doi.org/10.1073/pnas.93.14.7363>
- Sheng ZH, Rettig J, Takahashi M, Catterall WA. Identification of a syntaxin-binding site on N-type calcium channels. *Neuron* 1994; 13:1303-13; PMID:7993624; [http://dx.doi.org/10.1016/0896-6273\(94\)90417-0](http://dx.doi.org/10.1016/0896-6273(94)90417-0)
- Bezprozvanny I, Scheller RH, Tsien RW. Functional impact of syntaxin on gating of N-type and Q-type calcium channels. *Nature* 1995; 378:623-6; PMID:8524397; <http://dx.doi.org/10.1038/378623a0>
- Zhong H, Yokoyama CT, Scheuer T, Catterall WA. Reciprocal regulation of P/Q-type Ca₂₊ channels by SNAP-25, syntaxin and synaptotagmin. *Nat Neurosci* 1999; 2:939-41; PMID:10526329; <http://dx.doi.org/10.1038/14721>
- Dulubova I, Sugita S, Hill S, Hosaka M, Fernandez I, Südhof TC, et al. A conformational switch in syntaxin during exocytosis: role of munc18. *EMBO J* 1999; 18:4372-82; PMID:10449403; <http://dx.doi.org/10.1093/emboj/18.16.4372>
- Brunger AT. Structure of proteins involved in synaptic vesicle fusion in neurons. *Annu Rev Biophys Biomol Struct* 2001; 30:157-71; PMID:11340056; <http://dx.doi.org/10.1146/annurev.biophys.30.1.157>
- Jarvis SE, Barr W, Feng ZP, Hamid J, Zamponi GW. Molecular determinants of syntaxin 1 modulation of N-type calcium channels. *J Biol Chem* 2002; 277:44399-407; PMID:12221094; <http://dx.doi.org/10.1074/jbc.M206902200>
- Wiser O, Bennett MK, Atlas D. Functional interaction of syntaxin and SNAP-25 with voltage-sensitive L- and N-type Ca²⁺ channels. *EMBO J* 1996; 15:4100-10; PMID:8861939
- Sutton KG, McRory JE, Guthrie H, Murphy TH, Snutch TP. P/Q-type calcium channels mediate the activity-dependent feedback of syntaxin-1A. *Nature* 1999; 401:800-4; PMID:10548106; <http://dx.doi.org/10.1038/44586>

21. Fiebig KM, Rice LM, Pollock E, Brunger AT. Folding intermediates of SNARE complex assembly. *Nat Struct Biol* 1999; 6:117-23; PMID:10048921; <http://dx.doi.org/10.1038/5803>
22. Richmond JE, Weimer RM, Jorgensen EM. An open form of syntaxin bypasses the requirement for UNC-13 in vesicle priming. *Nature* 2001; 412:338-41; PMID:11460165; <http://dx.doi.org/10.1038/35085583>
23. Restituito S, Cens T, Barrere C, Geib S, Galas S, De Waard M, et al. The [beta]2a subunit is a molecular groom for the Ca²⁺ channel inactivation gate. *J Neurosci* 2000; 20:9046-52; PMID:11124981
24. Geib S, Sandoz G, Cornet V, Mabrouk K, Fund-Saunier O, Bichet D, et al. The interaction between the I-II loop and the III-IV loop of Cav2.1 contributes to voltage-dependent inactivation in a beta -dependent manner. *J Biol Chem* 2002; 277:10003-13; PMID:11790766; <http://dx.doi.org/10.1074/jbc.M106231200>
25. Sandoz G, Lopez-Gonzalez I, Stamboulian S, Weiss N, Arnoult C, De Waard M. Repositioning of charged I-II loop amino acid residues within the electric field by beta subunit as a novel working hypothesis for the control of fast P/Q calcium channel inactivation. *Eur J Neurosci* 2004; 19:1759-72; PMID:15078550; <http://dx.doi.org/10.1111/j.1460-9568.2004.03216.x>
26. Agler HL, Evans J, Tay LH, Anderson MJ, Colecraft HM, Yue DT. G protein-gated inhibitory module of N-type (ca(v)2.2) ca²⁺ channels. *Neuron* 2005; 46:891-904; PMID:15953418; <http://dx.doi.org/10.1016/j.neuron.2005.05.011>
27. Raghil A, Bertaso F, Davies A, Page KM, Meir A, Bogdanov Y, et al. Dominant-negative synthesis suppression of voltage-gated calcium channel Cav2.2 induced by truncated constructs. *J Neurosci* 2001; 21:8495-504; PMID:11606638
28. Page KM, Hebllich F, Davies A, Butcher AJ, Leroy J, Bertaso F, et al. Dominant-negative calcium channel suppression by truncated constructs involves a kinase implicated in the unfolded protein response. *J Neurosci* 2004; 24:5400-9; PMID:15190113; <http://dx.doi.org/10.1523/JNEUROSCI.0553-04.2004>
29. Page KM, Hebllich F, Margas W, Pratt WS, Nieto-Rostro M, Chaggar K, et al. N terminus is key to the dominant negative suppression of Ca(V)2 calcium channels: implications for episodic ataxia type 2. *J Biol Chem* 2010; 285:835-44; PMID:19903821; <http://dx.doi.org/10.1074/jbc.M109.065045>
30. Bucci G, Mochida S, Stephens GJ. Inhibition of synaptic transmission and G protein modulation by synthetic CaV2.2 Ca²⁺ channel peptides. *J Physiol* 2011; 589:3085-101; PMID:21521766; <http://dx.doi.org/10.1113/jphysiol.2010.204735>
31. Mochida S, Sheng ZH, Baker C, Kobayashi H, Catterall WA. Inhibition of neurotransmission by peptides containing the synaptic protein interaction site of N-type Ca²⁺ channels. *Neuron* 1996; 17:781-8; PMID:8893034; [http://dx.doi.org/10.1016/S0896-6273\(00\)80209-3](http://dx.doi.org/10.1016/S0896-6273(00)80209-3)
32. Rettig J, Heinemann C, Ashery U, Sheng ZH, Yokoyama CT, Catterall WA, et al. Alteration of Ca²⁺ dependence of neurotransmitter release by disruption of Ca²⁺ channel/syntaxin interaction. *J Neurosci* 1997; 17:6647-56; PMID:9254677
33. Harkins AB, Cahill AL, Powers JF, Tischler AS, Fox AP. Deletion of the synaptic protein interaction site of the N-type (CaV2.2) calcium channel inhibits secretion in mouse pheochromocytoma cells. *Proc Natl Acad Sci U S A* 2004; 101:15219-24; PMID:15471993; <http://dx.doi.org/10.1073/pnas.0401001101>
34. Keith RK, Poage RE, Yokoyama CT, Catterall WA, Meriney SD. Bidirectional modulation of transmitter release by calcium channel/syntaxin interactions in vivo. *J Neurosci* 2007; 27:265-9; PMID:17215385; <http://dx.doi.org/10.1523/JNEUROSCI.4213-06.2007>
35. Perez-Reyes E. Molecular physiology of low-voltage-activated t-type calcium channels. *Physiol Rev* 2003; 83:117-61; PMID:12506128
36. Foehring RC, Zhang XF, Lee JC, Callaway JC. Endogenous calcium buffering capacity of substantia nigral dopamine neurons. *J Neurophysiol* 2009; 102:2326-33; PMID:19675297; <http://dx.doi.org/10.1152/jn.00038.2009>
37. Atlas D. Signaling role of the voltage-gated calcium channel as the molecular on/off-switch of secretion. *Cell Signal* 2010; 22:1597-603; PMID:20388539; <http://dx.doi.org/10.1016/j.cellsig.2010.04.003>
38. Weiss N. Control of depolarization-evoked presynaptic neurotransmitter release by Cav2.1 calcium channel: old story, new insights. *Channels (Austin)* 2010; 4:431-3; PMID:20935476; <http://dx.doi.org/10.4161/chan.4.6.13613>