

Multistability in large scale models of brain activity

Mathieu Golos, Viktor Jirsa, Emmanuel Dauce

► **To cite this version:**

Mathieu Golos, Viktor Jirsa, Emmanuel Dauce. Multistability in large scale models of brain activity. Gennady S Cymbalyuk and Astrid A Prinz. the Twenty Second Annual Computational Neuroscience Meeting: CNS*2013, Jul 2013, paris, France. Biomed Central, 14 (Suppl 1), pp.P84, 2013. <inserm-00842317>

HAL Id: inserm-00842317

<http://www.hal.inserm.fr/inserm-00842317>

Submitted on 8 Jul 2013

HAL is a multi-disciplinary open access archive for the deposit and dissemination of scientific research documents, whether they are published or not. The documents may come from teaching and research institutions in France or abroad, or from public or private research centers.

L'archive ouverte pluridisciplinaire **HAL**, est destinée au dépôt et à la diffusion de documents scientifiques de niveau recherche, publiés ou non, émanant des établissements d'enseignement et de recherche français ou étrangers, des laboratoires publics ou privés.

POSTER PRESENTATION

Open Access

Multistability in large scale models of brain activity

Mathieu Golos¹, Viktor K Jirsa¹, Emmanuel Dauce^{1,2*}

From Twenty Second Annual Computational Neuroscience Meeting: CNS*2013
Paris, France. 13-18 July 2013

Rich spatiotemporal dynamical patterns, observed in the brain at rest, reveal several large-scale functional networks, presumably involved in different brain functions. In parallel, structural networks obtained by Diffusion Spectrum Imaging ("connectome") identify several

interconnected sub-networks that overlap with the functional networks [3]. Neural mass model simulations aim at developing realistic models of the brain activity. In particular, multiple fixed-point attractors can be identified and spontaneous alternation between several brain




Figure 1 Number of attractors obtained for different $(\Omega, ||W||)$ couples with $\gamma = 30$. Attractor patterns isodensity lines are indicated in black

* Correspondence: Emmanuel.Dauce@centrale-marseille.fr

¹Inserm, Aix-Marseille Université, INS UMR 1106, Marseille 13005, France

Full list of author information is available at the end of the article

“states” can be obtained [2]. The number and the organization of the “fixed-point” attractors strongly vary depending of the parameters used in the model.

In order to provide a general view of connectome-based dynamical systems, we developed a simplified neural mass model inspired by the continuous Hopfield network [4], to which we add a stochastic component, and a dynamic threshold (eq.2) that prevents reaching the two trivial attractors of system (eq.1).

$$\tau \frac{dx_i}{dt} = -x_i + \sum_{j=1}^N W_{ij} \phi(x_j, \gamma, \theta) + \sigma^2 \eta_i(t) \quad (1)$$

$$\tau_\theta \frac{d\theta}{dt} = -\theta + \frac{\Omega}{N} \sum_{i=1}^N \phi(x_i, \gamma, \theta) \quad (2)$$

$$\phi(x_i, \gamma, \theta) = \frac{1}{2} (1 + \tanh(\gamma(x_i - \theta))) \quad (3)$$

where W is the $N = 66$ nodes human connectome [3], x is the node potential, ϕ is the node output, θ is the adaptive threshold, $\eta(t)$ is a white noise and σ the diffusion parameter. Three main parameters (excitation strength $\|W\|$, inhibition strength Ω and node excitability γ) are systematically varied in order to identify the fixed points attractors of the dynamics (Figure 1).

The adaptive threshold plays the role of a global inhibition and allows stabilizing many attractors, where the $\Omega / \|W\|$ ratio controls the sparsity of the activity. Then multistable dynamical systems are obtained in a large region of the parameter space, allowing to identify many different attractor patterns. The number of attractors is found to increase with the value of gamma (node excitability), at the expense of their intrinsic stability. Itinerant dynamics is obtained when a significant noise level is introduced in the system. Then, several “attractive” patterns can be reached on a single trajectory, where the duration of the visit reflects the stability of the pattern.

The simple dynamical system we have implemented allows exploring a large region of the parameters space, but is also capable of reproducing some aspects of the brain’s spontaneous activity (switching between different attractors). The set of attractors we find in some regions of the parameters space share similarities with the different functional modes observed in the resting-state activity [1].

Author details

¹Inserm, Aix-Marseille Université, INS UMR 1106, Marseille 13005, France.

²Ecole Centrale Marseille, Marseille 13013, France.

Published: 8 July 2013

References

1. Damoiseaux JS, Rombouts S, Barkhof F, Scheltens P, Stam C, Smith P, Beckmann C: **Consistent resting-state networks across healthy subjects.** *PNAS* 2006.
2. Deco G, Jirsa VK: **Ongoing cortical activity at rest: criticality, multistability and ghost attractors.** *J Neuroscience* 2012.
3. Hagmann P, Cammoun L, Gigandet X, Meuli R, Honey C, Wedeen V, Sporns O: **Mapping the structural core of human cerebral cortex.** *PLoS Biol* 2008.
4. Hopfield JJ: **Neural networks and physical systems with emergent collective computational abilities.** *Proc Natl Acad Sci USA* 1982.

doi:10.1186/1471-2202-14-S1-P84

Cite this article as: Golos et al.: Multistability in large scale models of brain activity. *BMC Neuroscience* 2013 **14**(Suppl 1):P84.

Submit your next manuscript to BioMed Central and take full advantage of:

- Convenient online submission
- Thorough peer review
- No space constraints or color figure charges
- Immediate publication on acceptance
- Inclusion in PubMed, CAS, Scopus and Google Scholar
- Research which is freely available for redistribution

Submit your manuscript at
www.biomedcentral.com/submit

