

## **Estrogenic and anti-estrogenic activity of 23 commercial textile dyes.**

Ingrid Bazin, Aziza Ibn Hadj Hassine, Yosra Haj Hamouda, Wissem Mnif, Ahgheb Bartegi, Miguel Lopez-Ferber, Michel De Waard, Catherine Gonzalez

► **To cite this version:**

Ingrid Bazin, Aziza Ibn Hadj Hassine, Yosra Haj Hamouda, Wissem Mnif, Ahgheb Bartegi, et al.. Estrogenic and anti-estrogenic activity of 23 commercial textile dyes.. *Ecotoxicology and Environmental Safety*, Elsevier, 2012, 85, pp.131-6. <10.1016/j.ecoenv.2012.08.003>. <inserm-00843006>

**HAL Id: inserm-00843006**

**<http://www.hal.inserm.fr/inserm-00843006>**

Submitted on 10 Jul 2013

**HAL** is a multi-disciplinary open access archive for the deposit and dissemination of scientific research documents, whether they are published or not. The documents may come from teaching and research institutions in France or abroad, or from public or private research centers.

L'archive ouverte pluridisciplinaire **HAL**, est destinée au dépôt et à la diffusion de documents scientifiques de niveau recherche, publiés ou non, émanant des établissements d'enseignement et de recherche français ou étrangers, des laboratoires publics ou privés.

# 1 **Estrogenic and anti-estrogenic activity of 23 commercial** 2 **textile dyes**

3  
4 Ingrid Bazin<sup>a,\*</sup>, Aziza Ibn Hadj Hassine<sup>a</sup>, Yosra Haj Hamouda<sup>b</sup>, Wissem Mnif<sup>b</sup>, Ahgleb  
5 Bartegi<sup>c</sup>, Miguel Lopez-Ferber<sup>a</sup>, Michel De Waard<sup>d,e</sup>, Catherine Gonzalez<sup>a</sup>  
6

7 <sup>a</sup> Ecole des Mines d'Alès, LGEI center, 6 avenue de Clavière 30319 Alès Cedex, France.

8 <sup>b</sup> Laboratory of Biochemistry, Institute of Biotechnology, Monastir, Tunisia

9 <sup>c</sup> Department of Biology, Faculty of Sciences - King Faisal University P.O. Box 1759 31982  
10 Al Hassa, Saudi Arabia

11 <sup>d</sup> Inserm U836, Grenoble Institut of Neurosciences, Site Santé de la Tronche, 38042 Grenoble,  
12 Cedex 09, France

13 <sup>e</sup> Université Joseph Fourier, Grenoble, France

14  
15 \*Corresponding author: [ingrid.bazin@mines-ales.fr](mailto:ingrid.bazin@mines-ales.fr)

---

## 16 **Abstract**

17 The presence of dyes in wastewater effluent of textile industry is well documented. In  
18 contrast, the endocrine disrupting effects of these dyes and wastewater effluent have been  
19 poorly investigated. Herein, we studied twenty-three commercial dyes, usually used in the  
20 textile industry, and extracts of blue jean textile wastewater samples were evaluated for their  
21 agonistic and antagonistic estrogen activity. Total estrogenic and anti-estrogenic activities  
22 were measured using the Yeast Estrogen Screen bioassay (YES) that evaluates estrogen  
23 receptor binding-dependent transcriptional and translational activities. The estrogenic  
24 potencies of the dyes and wastewater samples were evaluated by dose-response curves and  
25 compared to the dose-response curve of 17 $\beta$ -estradiol (E2), the reference compound. The  
26 dose-dependent anti-estrogenic activities of the dyes and wastewater samples were  
27 normalized to the known antagonistic effect of 4-hydroxytamoxifen (4-OHT) on the induction  
28 of the lac Z reporter gene by E2. About half azo textile dyes have anti-estrogenic activity with  
29 the most active being Blue HFRL. Most azo dyes however have no or weak estrogenic  
30 activity. E2/dye or E2/waste water ER competitive binding assays show activity of Blue  
31

32 HFRL, benzopurpurine 4B, Everzol Navy Blue FBN, direct red 89 BNL 200% and waste  
33 water samples indicating a mechanism of action common to E2. Our results indicate that  
34 several textile dyes are potential endocrine disrupting agents. The presence of some of these  
35 dyes in textile industry wastewater may thus impact the aquatic ecosystem.

36

37 *Keywords:* textile dyes, estrogenic activity, anti-estrogenic activity, industrial textile effluent.

---

38

## 39 **1. Introduction**

40

41 Dyes are widely used in most industries such as those manufacturing papers, plastics, food,  
42 cosmetics, textiles or leathers. These dyes are useful to colour the final products. Dyes are  
43 classified depending on their colors, their chemical structures and/or their origin (natural or  
44 synthetic). Natural dyes most frequently originate from plants (such as riboflavin or  $\beta$ -  
45 carotene). Determination of the chemical structures of natural dyes and the accomplishment of  
46 their synthesis allow the gradual replacement of these natural dyes by their synthetic  
47 counterparts. Approximately 10.000 commercial dyes are used in the coloring industry. More  
48 than 10% of dyestuff used during the coloring processes does not bind to the fibers and  
49 therefore these excess dyes are released into the environment producing serious  
50 environmental pollution (Pearce et al., 2003; Rajeswari et al., 2011). The presence of these  
51 dyes in wastewater and subsequently in water resources, even at very low concentrations, is  
52 easy to observe visually as the result of textile industry activities. They may increase effluent  
53 toxicity and lead to environmental damage (Robinson et al., 2002). In addition, many  
54 synthetic dyes are poorly biodegradable. In some dyehouse effluents, dye concentration can  
55 reach up to 400 mg/l (O'Neill et al., 1999). A specific study has even demonstrated that these  
56 concentrations can exceed 600 mg/L in Nigeria (Yusuff and Sonibare, 2004). Their presence  
57 in water reduces light penetration and has a negative impact on photosynthesis. Moreover, the  
58 dyeing process itself generally also contributes to the water body contamination by chromium,  
59 zinc and copper which are all toxic to aquatic plants and fish below 1.0 mg/L (Eremektar et  
60 al., 2007; Sharma et al., 2007; Verma, 2008). The research on textile wastewater toxicity,  
61 which has been carried out so far, shows how the action of toxic dyes occurs at different  
62 levels of the food chain, from food supply (i.e. algae and plants) to consumers (i.e.  
63 crustaceans and fishes) (Manu et al., 2003; Robinson et al., 2002; Sharma et al., 2007; Soni et  
64 al., 2006; Tigini et al., 2011). Most research effort has been devoted towards elucidating or

65 improving the degradation mechanisms of textile dyes with the hope to reduce their toxicity  
66 levels (Phugare et al., 2011a,b). Environmental regulations in most countries (i.e. EU  
67 directive 91/271) have prioritized in wastewater dye decontamination in order to minimize  
68 environmental damage (Robinson et al., 2001). Although the Environmental Protection  
69 Agency (EPA) has emitted new practice guidelines for environmental management, the  
70 presence of pollution dyes remains a serious environmental issue specifically for small textile  
71 industries in various countries (e.g. China, India, Taiwan) where working conditions and low  
72 economic status does not allow an efficient wastewater treatment before disposal into water  
73 sources (Mathur et al., 2005; Gregory et al., 2007; You et al., 2009). Since the mid nineties,  
74 the links between dyes, environmental impact and cancer emergence have been the subject of  
75 considerable interest both from researchers and from the general public. However, the causal  
76 relationship between dyes and certain types of cancer is difficult to establish. Several studies  
77 have been conducted on the toxicity, mutagenicity and genotoxicity of textile dyes (Bakshi et  
78 al., 2003; Ben Mansour et al., 2007; Dogan et al., 2005; Durnev et al., 1995; Mathur et al.,  
79 2007; Schneider et al., 2004). In the case of azo dyes, the increase in bladder cancer  
80 incidence, observed among textile industry workers, has been linked to prolonged exposure to  
81 these dyes. A report of experts from the “Cosmetic Ingredient Review Committee” confirmed  
82 that some anthraquinone dyes, like the disperse Blue 7 dye, used in the cosmetic industry as  
83 hair colorant and in textile, induce genotoxicity in bacteria (Cosmetic ingredient, 2007). Since  
84 then, research on these dyes confirmed their carcinogenic effects for humans and animals  
85 (Tsuda et al., 2001). This toxic effect has been linked to their presence in the environment  
86 (Dogan et al., 2005; Chou et al., 2007; Tigini et al., 2011). From an environmental point of  
87 view, toxicity, genotoxicity and mutagenicity of industrial effluents have been demonstrated  
88 (Alves de Lima et al., 2007; Grinevicius et al., 2009; Tigini et al., 2011). However, these  
89 effects have generally not been linked to the presence of dyes. Chou and collaborators  
90 associated the dioxin-like activity of some dyeing wastewater with the presence of specific  
91 anthraquinone dyes (Chou et al., 2006, 2007). In the same studies, these authors demonstrated  
92 that disperse blue 56 can bind the aryl hydrocarbon receptor (AhR), which is involved in  
93 many physiological functions such as cell regulation and reproduction. Two Brazilian studies  
94 have identified that the mutagenic activity of the Cristais River, a drinking water source of  
95 São Paulo, is caused by the presence of three blue dyes; C.I. Disperse Blue 37 contributing to  
96 55% of this effect (De Aragão et al., 2005).

97 Up to now, the estrogenic activity of dyes has seldom been studied. Rare studies  
98 mainly report the effects of food colors, such as tartazine (E102) and erythrosin B (E127).  
99 Both compounds affect chromosome structure and increase Estrogen Receptor (ER) site-  
100 specific DNA binding to Estrogen Response Element (ERE) in HTB 133 cells (Roychoudhury  
101 et al., 1989) and in the E-screen test (Datta et al., 2008). The present manuscript investigates  
102 both the estrogenic and anti-estrogen activities of twenty three pure commercial textile dyes  
103 using the Yeast Estrogen Screen (YES). This *in vitro* assay has been developed for the  
104 detection of endocrine disrupting compounds (EDCs). In order to indentify estrogen and anti-  
105 estrogen activity of textile dyes, the YES assay has been used to probe two different modes of  
106 action. First, a reporter gene assay measures the impact of dye binding on ER on its ability to  
107 promote binding-dependent transcriptional and translational activity. Second, an ER  
108 competitive binding assay measures how a dye competes with 17 $\beta$ -estradiol (E2)-dependent  
109 ER activation. In addition, the endocrine activity (estrogenic and anti-estrogenic activity) has  
110 been assessed on a textile effluent coming from a blue jean manufacturing Tunisian industry  
111 in an attempt to correlate the endocrine activity observed for commercial compounds and that  
112 obtained for dyeing wastewater.

113

## 114 2. Materials and methods

115

### 116 2.1. Selected dyes

117

118 All the dyes selected (23) in this study are used in the textile industry (Table 1 and 2).  
119 Only, 17 on 23 dyes have a known or communicated structure. The structure of the 6 other  
120 dyes were confidential and not available in the chemical abstracts service. However, the  
121 safety and chemical information certificate of analysis usually indicates the nature of the dye  
122 (azoic or anthraquinone-type chemical structure). For each dye, a stock solution (10 g/L) was  
123 prepared by dissolving in distilled water, followed by filtration through Whatmann No. 5 filter  
124 paper. All these dyes appeared perfectly soluble in water at this concentration as assessed by  
125 the absence of precipitate. For each dye, the endocrine activity was assessed at four  
126 concentrations in the range of  $1 \times 10^{-5}$  g/L to 1 g/L. The natural fluorescence of the dyes and  
127 their interference on the fluorescence emitted by yeast was determined beforehand in order to  
128 avoid spurious signals generated by the YES test in our experimental conditions. The  
129 fluorescence of each dye has been measured in the absence of yeast cells and subtracted from

130 the data obtained for estrogenic tests. The quenching potential of the dyes was also assessed  
131 on the yeast fluorescence emission during the antiestrogenic test as follows. The fluorescence  
132 of yeast cell has been measured after 6 h of incubation of the reference compound E2 (from  
133 Sigma-Aldrich, St Quentin-Fallavier, France). Then, each dye was added to the medium at  
134 various concentrations and the fluorescence quenching was measured immediately.  
135 Compounds inducing more than 10% of fluorescence quenching were not investigated further  
136 for anti-estrogenic activity (noted by an asterisk in Table 2). The cytotoxicity of each dye was  
137 evaluated by measuring yeast growth at an O.D. of 600 nm.

138

## 139 2.2. Wastewater sampling and extraction

140

141 Wastewater samples (1 L) were collected from the textile effluent of an industrial blue  
142 jeans factory in Tunisia. Raw water samples were centrifuged (2000 g, 15 min) to eliminate  
143 solids in suspension. Supernatants were then extracted by solid-phase extraction as described  
144 by Pillon et al. (2005). Briefly, aqueous samples were concentrated on reverse-phase C18 (5  
145 g, 20 mL) cartridges (Sigma-Aldrich, St Quentin-Fallavier, France) preconditioned with  
146 methanol. Compounds elution from the column was triggered using methanol followed by  
147 hexane. Eluates were dried at 37°C in a rotary evaporator and residues were taken up in 2 mL  
148 methanol (concentration factor: 500).

149

## 150 2.3. Yeast estrogen screen assay (YES)

151

### 152 2.3.1. Assay of estrogenic and anti-estrogenic activity

153

154 Both agonist and antagonist activities of chemical dyes were examined using the Yeast  
155 strain BY4741 (Euroscarf, Frankfurt, Germany). This recombinant yeast strain carries the  $\beta$ -  
156 galactosidase reporter gene under the control of the ERE and contains the human ER cloned  
157 into the constitutive yeast expression vector pAAH5 (García-Reyero et al., 2001). The test  
158 measures  $\beta$ -galactosidase activity (fluorescence at 460 nm and excitation at 355 nm) with a  
159 fluorimeter (Fluoroskan Twinkle LB 970, BERTHOLD Technologies) after 6 h of exposure to  
160 the compounds to evaluate. Tests were performed in 96-well plates. To determine the estrogen  
161 agonist activity of dyes and effluent samples, E<sub>2</sub> was used as a positive control and distilled  
162 water was used as a negative control. Four dye concentrations were tested in the range of

163  $1 \times 10^{-5}$  g/L to 1 g/L. For the effluent samples, a series of dilutions (1 to 1000 of concentrated  
164 extract) were tested for their estrogenic activity. To determine the estrogen antagonist activity  
165 of the dyes, 4-OHT was used as a positive control and E<sub>2</sub> (1 nM) was used as a negative  
166 control. In the antagonist test, the ability of the dye to compete with E<sub>2</sub> for binding to ER  
167 and/or inhibit the receptor functions was evaluated. The tested dyes or 4-OHT, the positive  
168 control, were combined with 1 nM E<sub>2</sub> before the start of the assay. All experiments were  
169 performed in triplicate. For estrogen agonist activity, the half maximal effective concentration  
170 (EC<sub>50</sub>) was calculated based on the sigmoidal dose–effect curve of E<sub>2</sub>. For estrogen antagonist  
171 activity, the half-maximal effective concentration (AC<sub>50</sub>) was calculated based on the  
172 sigmoidal dose–effect curve of 4-OHT.

173

#### 174 2.3.2. Competitive binding assay

175

176 Competition between dyes and/or sample effluent and E<sub>2</sub> was measured at various  
177 concentrations of E<sub>2</sub>. Textile dye concentration used is 0.01 g/L and the textile dyeing effluent  
178 sample at a dilution factor 5 of the concentrated extract. Thus any decrease of β-galactosidase  
179 activity after 6 h of exposure indicates that the dye induces a decrease in E<sub>2</sub> binding.

180

#### 181 2.4. UV/Visible spectrum deconvolution

182

183 The absorbency spectrum of a water sample can be decomposed into a few numbers of  
184 spectra (reference spectra). The shape of the UV spectrum can be considered as a linear  
185 combination of defined spectra (REF<sub>1</sub>,..., REF<sub>p</sub>) related to potential compounds present in  
186 studied water sample (Thomas et al., 1996).  $S_w = \sum a_i \times REF_i \pm r$ , where  $S_w$  is the final  
187 spectra,  $a_i$  and  $r$  are the coefficient of the  $i^{\text{th}}$  reference spectra and the admitted error,  
188 respectively. The Secomam company (Alès, France) has developed the UVPro software based  
189 on advanced UV spectral deconvolution (UV PRO, 2000) which allows creating dedicated  
190 models and determination of reference spectra from a set of studied wastewater UV spectra.  
191 The UVPro software has been applied to the textile effluent of an industrial blue jeans factory  
192 of Tunisia. The reference spectra used for the deconvolution are obtained from the 23 selected  
193 textiles dyes (Table 1). Using the deconvolution model, it is possible to assess the  
194 contribution of studied dyes in the wastewater spectrum.

195

196 **3. Results / Discussion**

197

198 3.1. Estrogenicity / anti-estrogenicity studies of individual dyes

199

200 YES allows a fast determination of both endocrine agonist and antagonist actions. The  
201 relative estrogenicity of each dye has been estimated using YES by comparison to the  
202 estrogenicity of E2 as reference compound. Typical dose-response curves for E2 have been  
203 established (data not shown). In the present study, we have essentially analyzed azo dyes.  
204 These dyes are extensively used for dyeing cotton in textile industries. By using YES assay in  
205 the present work, we have analyzed the degree of interference of textile dyes with the  
206 endocrine system considering both the potential agonist and antagonist actions. A great  
207 variability between dyes is observed. The YES assay data illustrate a dose-dependent estrogen  
208 agonist activity from  $1 \times 10^{-5}$  g/L to 1 g/L for three dyes (Yellow Flavina CXL, Reactive dye  
209 red 3BS and Solvent yellow 56) (Figure 1). All these dyes have weak estrogenic effects since  
210 the maximum activity is obtained at 1 g/L. In addition, the estrogenic effect of the dyes did  
211 not saturate precluding the determination of the EC<sub>50</sub> value, contrary to the positive control E2  
212 (EC<sub>50</sub> value at 1 ng/L) (Figure 1). Higher dye concentrations were not investigated, since  
213 water body dye concentration in the environment never exceeds 1 g/l. Table 2 also  
214 summarizes the fact that the twenty other dyes had no estrogenic effect by themselves. These  
215 results indicate that these dyes are not xenoestrogens that should highly impact the  
216 environment, as compared to E2 or other known xenoestrogens (e.g. bisphenol A, paraben...)  
217 (Routledge et al., 1998).

218 In addition, some textile dyes have an anti-estrogenic activity. The anti-estrogenic  
219 reference compound 4-OHT was used to demonstrate efficient anti-estrogenic activity when  
220 incubated in the presence of 1 nM E2. The AC<sub>50</sub> value of 4-OHT was 0.5 μM or 0.02 mg/L  
221 (Table 2). Similar experiments were conducted with the dyes. Only some dyes have been  
222 tested (Figure 2), the other ones, marked by an asterisks, could not be tested because they  
223 quenched the β-galactosidase fluorescence. At a concentration of 1 g/L, antagonistic activities  
224 of these dyes vary from 100% to 10% inhibition according to their colour (Table 2 and Figure  
225 2A). Blue and red dyes show the highest inhibition potential. Everzol navy blue FBN, blue  
226 HFRL, Direct Red 89 BNL 200% and Benzopurpurine 4B (Red 4B) are the most potent  
227 inhibitors tested. Their inhibitory effects remain however inferior to that of 4-OHT. At 1 g/L,  
228 these dyes are the only ones that induce a complete inhibition of the reporter gene in yeast



229 cells. Among these four dyes, Blue HFRL is the strongest antagonist; its anti-estrogenic  
230 potency being approximately 750-fold less than that of 4-OHT. A reduction of 50% of  $\beta$ -  
231 galactosidase activity is observed for concentrations as low as 15 mg/L. These concentration  
232 values are thus considerably lower than the total color level found in the environment (400  
233 mg/L). The antagonist effect of three other dyes (Benzopurpurine 4B (Red 4B), Everzol Navy  
234 Blue FBN, and Direct Red 89 BNL) were, respectively, 3500-, 2500-, and 1250-fold less than  
235 that of 4-OHT. In contrast, all other dyes were either weak ER inhibitors or non-ER  
236 inhibitors. For instance, Everzol Blue ED, Red alpacide 3BL, Direct Blue 71, Blue ED 250,  
237 Direct Black VSF and Blue DERF were weak ER inhibitors. The anti-estrogenic activity for  
238 Blue DERF and Direct Blue 71 at 1 g/l, are 10% and 22%, respectively. Among these dyes,  
239 most are of the azo class with sulfonated aniline (Benzopurpurine 4B, Direct Red 89 BNL  
240 200%) and one is an anthraquinone dye, Everzol Blue ED. The Direct Black VSF AZO-  
241 FREE, Blue ED 250, both being polyazo dyes, Everzo Yellow ED and Red ED (for which no  
242 structures can be disclosed) induce inhibition of  $\beta$ -galactosidase expression by E2 with  $AC_{50}$   
243  $> 1$  g/L.

244 Competition between the textile dyes and E2 was also measured by varying the  
245 concentration of E2 and maintaining a constant dye concentration (0.01 g/L, a concentration  
246 that does not fully inhibit E2 effect, Figure 2A). Benzopurpurine 4B, Blue HFRL and Everzol  
247 Navy Blue FBN can compete with E2 for binding to ER (Figure 2B). These dyes induces a  
248 slight inhibition of E2 estrogenic activity when E2 is used at concentrations equal or lower  
249 than  $1 \cdot 10^{-8}$  M. No effect is observed for higher concentrations of E2. For example, this  
250 activity decreased from 200000 RLU without dye to 100000 RLU in presence of the Blue  
251 HFRL at  $10^{-8}$  M E2. This study also shows that the Red Alpacide 3BL induces a 44%  
252 inhibition of estrogenic activity and the metal-complexed azo dye yellow 4G induces a 13%  
253 inhibition (Table 2). The competitive binding assay in the YES assay showed that these dyes  
254 can bind specifically to ER in order to induce these antagonist effects.

255

### 256 3.2. Estrogenicity / anti-estrogenicity studies of textile effluent

257

258 The textile effluent sample analyzed presents a weak estrogenic effect at the maximum  
259 concentration tested (5-fold dilution). At higher concentrations of the sample, the sample  
260 compounds reduce cell viability (data not show) precluding the use of the sample for the  
261 estrogenic assay. For the dilution range (5- to 10000-fold), results from the YES bioassay

262 indicate that the industrial textile effluent presents a low (below 15%) but significant  
263 estrogenic activity (Figure 3A). Also, the effluent sample inhibits 60% of estrogenic activity  
264 (Figure 3B). In competition experiments, in which E2 concentration is varied and a single  
265 dilution dose of the effluent sample is used (5-fold dilution), the anti-estrogenic effect is  
266 observable until  $10^{-5}$  M of E2 (Figure 3C). The competition effect of the sample is most  
267 significant at  $10^{-8}$  M.

268         The composition of the effluent sample in dyes has not been fully determined. Spectral  
269 analyses indicate that most of its components are blue dyes. Indeed, UV/Visible  
270 deconvolution of the spectrum of a blue jeans manufacturer's effluent of Tunisia indicates that  
271 the sample contains mainly Blue DERF, Direct Blue 71, Everzol navy Blue FBN, and Grey  
272 GGL dye according to the spectrum of each individual dye (Figure 4). The deconvolution of  
273 the sample spectrum did not predict the presence of other dyes. Interestingly, our studies have  
274 demonstrated that at least three of these dyes presented anti-estrogenic activity with the YES  
275 assay. Most blue and navy blue dyes are derived from the Reactive Black 5 azo dye, which is  
276 classified Xn (harmful). It is also suspected to be mutagenic and to be associated to bladder  
277 cancer development (You et al., 2009). Nevertheless, without a complete characterization of  
278 the nature of the dyes being released in the local environment of this factory, the anti-  
279 estrogenic effect of the effluent sample can't be unequivocally be attributed to these blue  
280 dyes. Since this effluent is essentially blue, and that some blue dyes are among those  
281 presenting the highest anti-estrogenic effect, suspicion remains however high that this  
282 Tunisian factory releases harmful components in the environment. In addition, it is worth  
283 noting that the most active dyes have anti-estrogenic  $AC_{50}$  values (between 15 mg/L and 70  
284 mg/L) that are well below the dye concentration that can be detected in some water bodies  
285 (400 mg/L). If these same dyes are indeed present in the environment, they would present a  
286 real anti-estrogenic activity to local water consumers.

287

#### 288 **4. Conclusions**

289

290         Our results on the endocrine effects of textile dyes come in complement of those  
291 already published on the carcinogenic and mutagenic effects of dyes. They point to the fact  
292 that some dyes, mainly blue and red dyes, may be endocrine disruptor compounds. These  
293 observations raise two issues. Concerning environmental pollution, some of these dyes may  
294 be present at concentrations high enough that they may indeed affect life quality by promoting

295 tumor generation. Textile sewage is therefore a potential health hazard that should require a  
296 better communication on the dyes used by the industry and its method of disposal. In that  
297 respect, it appears as particularly important to efficiently treat industrial effluent containing  
298 azo dyes before they get discharged into the environment. The second issue concerns the  
299 pharmacological effect of these textile dyes. Besides 4-OHT, few compounds have anti-  
300 estrogenic activity. It is therefore of interest to note that some of the dyes that we  
301 characterized have similar anti-estrogenic activity, suggesting that they may be lead  
302 compounds for the development of new 4-OHT like anti-estrogenic compounds.

303

## 304 **References**

305

306 Alves de Lima, R.O., Bazo A.P., Salvadori, D.M., Rech, C.M., de Palma Oliveira, D., de  
307 Aragão Umbuzeiro, G., 2007. Mutagenic and carcinogenic potential of a textile azo dye  
308 processing plant effluent that impacts a drinking water source. *Mutat Res.* 626 (1-2), 53-  
309 60.

310 Baêta, B.E., Aquino, S.F., Silva, S.Q., Rabelo, C.A., 2011. Anaerobic degradation of azo dye  
311 Drimaren blue HFRL in UASB reactor in the presence of yeast extract a source of carbon  
312 and redox mediator. *Biodegradation*, DOI: 10.1007/s10532-011-9499-4.

313 Bakshi, D.K., Sharma, P., 2003. Genotoxicity of textile dyes evaluated with Ames test and  
314 rec-assay. *J Environ Pathol Toxicol Oncol.* 22 (2), 101-9.

315 Ben Mansour, H., Corroler, D., Barillier, D., Ghedira, K., Chekir, L., Mosrati, R., 2007.  
316 Evaluation of genotoxicity and pro-oxidant effect of the azo dyes: acids yellow 17, violet  
317 7 and orange 52, and of their degradation products by *Pseudomonas putida* mt-2. *Food*  
318 *Chem Toxicol.* 45 (9),1670-7.

319 Chou, P. H., Matsui, S., Matsuda, T., 2006. Detection and identification of dyes showing  
320 AhR-binding affinity in treated sewage effluents. *Water Sci Technol.* 53 (11), 35-42.

321 Chou, P. H., Matsui, S., Misaki, K., Matsuda, T., 2007. Isolation and identification of  
322 xenobiotic aryl hydrocarbon receptor ligands in dyeing wastewater. *Environ Sci Technol.*  
323 41(2), 652-7.

324 Cosmetic Ingredient Review Expert Panel., 2007. Final report on the safety assessment of  
325 disperse blue 7. *Int J. Toxicol.* 26, suppl 2, 65-77; DOI: 10.1080/10915810701351210

326 Datta, P., Lundin-Schiller, S., 2008. Estrogenicity of the synthetic food colorants tartrazine,  
327 erythrosin B, and sudan I in an estrogen-responsive human breast cancer cell line.  
328 *Journal of the Tennessee Academy of Science*, 83 (4).

329 De Aragão Umbuzeiro, G., Freeman, H. S., Warren, S. H., de Oliveira, D.P., Terao, Y.,  
330 Watanabe, T., Claxton, L. D., 2005. The contribution of azo dyes to the mutagenic  
331 activity of the Cristais River. *Chemosphere*. 60 (1), 55-64.

332 Dogan, E.E., Yesilada, E., Ozata, L., Yologlu, S., 2005. Genotoxicity testing of four textile  
333 dyes in two crosses of *Drosophila* using wing somatic mutation and recombination test.  
334 *Drug Chem Toxicol*. 28 (3), 289-301.

335 Durnev, A. D., Oreshchenko, A. V., Kulakova, A.V., Beresten, N. F., 1995. Analysis of  
336 cytogenetic activity of food dyes. *Vopr Med Khim*. 41 (5), 50-3.

337 Eremektar, G., Selcukb, H., Mericc, S., 2007. Investigation of the relation between COD  
338 fractions and the toxicity in a textile finishing industry wastewater: effect of  
339 preozonation. *Desalination* 211, 314–320.

340 García-Reyero, N., Grau, E., Castillo, M., López de Alda, M. J., Barceló, D., Piña, B., 2001.  
341 Monitoring of endocrine disruptors in surface waters by the yeast recombinant assay. *J*  
342 *Environ Toxicol Chem*. 20 (6), 1152-8.

343 Gregory, P., 2007. Toxicology of textile dyes in “environmental aspects of textile dyeing” in  
344 *Environmental aspects of textile dyeing*; Christie R .M. eds.; WP chp 3, pp 44-73.

345 Grinevicius, V. M., Geremias, R., Laus, R., Bettega, K. F., Laranjeiras, M. C., Fávère, V. T.,  
346 Wilhelm Filho, D., Pedrosa, R. C., 2009. Textile effluents induce biomarkers of acute  
347 toxicity, oxidative stress, and genotoxicity. *Arch Environ Contam Toxicol*. 57 (2), 307-  
348 14.

349 Manu B., Chaudhari S., 2003. Decolorization of indigo and azo dyes in semicontinuous  
350 reactors with long hydraulic retention time. *Process biochemistry*. 38(8), 1213-1221.

351 Mathur, N., Bhatnagar, P., 2007. Mutagenicity assessment of textile dyes from Sanganer  
352 (Rajasthan). *J Environ Biol*. 28 (1), 123-6.

353 Matur, N., Bhatnagar, P., Bakre, P., 2005. Assessing mutagenicity of textiles dyes from Pali  
354 (Rajasthan) using Ames bioassay applied ecology and environmental research 4(1): 111-  
355 118.

356 O’Neill, C., Freda, R., Hawkes, D. L., Hawkes, N. D., Lourenço, H. M., Pinheiro, W. D.,  
357 1999. Colour in textile effluents – sources, measurement, discharge consents and  
358 simulation: a review. *Journal of Chem Tech and Biotech*. 74, (11), 1009-1018.

359 Pearce, C.I., Lloyd, J.R., Guthrie, J.T., 2003. The removal of colour from textile wastewater  
360 using whole bacterial cells: a review. *Dyes and Pigments*. (58),3, Pages 179–196

361 Phugare, S. S., Kalyani, D. C., Patil, A. V., Jadhav, J. P., 2011. Textile dye degradation by  
362 bacterial consortium and subsequent toxicological analysis of dye and dye metabolites  
363 using cytotoxicity, genotoxicity and oxidative stress studies. *J Hazard Mater*. 186 (1),  
364 713-23.

365 Phugare, S. S., Kalyani, D. C., Surwase, S. N., Jadhav, J. P., 2011. Ecofriendly degradation,  
366 decolorization and detoxification of textile effluent by a developed bacterial consortium.  
367 *Ecotoxicol Environ Saf*. 74, (5), 1288-96.

368 Pillon, A., Servant, N., Vignon, F., Balaguer, P., Nicolas, J. C., 2005., In vivo  
369 bioluminescence imaging to evaluate estrogenic activities of endocrine disrupters. *Anal*  
370 *Biochem*. 340(2), 295-302.

371 Rajeswari, K., Subashkumar R., Vijayaraman K., 2011. Biodegradation of Mixed Textile  
372 Dyes by Bacterial Strains Isolated from Dyewaste Effluent. *Research Journal of*  
373 *Environmental Toxicology*, 5: 97-107.

374 Robinson, T., Chandran, B., Nigam, P., 2001. Studies on the decolourisation of an artificial  
375 textile-effluent by white-rot fungi in N-rich and N-limited media. *Appl Microbiol*  
376 *Biotechnol*. 57(5-6):810-3.

377 Robinson, T., Chandran, B., Nigam, P., 2002. Removal of dyes from a synthetic textile dye  
378 effluent by biosorption on apple pomace and wheat straw. *Water Res*. 36(11):2824-30.

379 Routledge, E.J., Parker, J., Odum, J., Ashby, J., Sumpter, J.P., (1998). Some alkyl hydroxy  
380 benzoate preservatives (parabens) are estrogenic. *Toxicol. Appl. Pharmacol*. 153, 12-19.

381 Roychoudhury, A., Giri, A. K., 1989. Effects of certain food dyes on chromosomes of *Allium*  
382 *cepa*. *Mutat Res*. 223 (3), 313-9.

383 Schneider, K., Hafner, C., Jäger, I., 2004. Mutagenicity of textile dye products. *J Appl*  
384 *Toxicol*. 24 (2), 83-91.

385 Sharma, K.P., Sharma, S., Sharma, S., Singh, P.K., Kumar, S., Grover, R., Sharma, P.K.,  
386 2007. A comparative study on characterization of textile wastewaters (untreated and  
387 treated) toxicity by chemical and biological tests. *Chemosphere*. 69, 48–54.

388 Soni, P., Sharma, S., Sharma, S., Kumar, S., Sharma, K.P., 2006. A comparative study on the  
389 toxic effects of textile dye wastewaters (untreated and treated) on mortality and RBC of a  
390 freshwater fish *Gambusia affinis* (Baird and Gerard). *J Environ Biol*. 27(4):623-8.

391 Tigini, V., Giansanti, P., Mangiavillano, A., Pannocchia, A., Varese, G. C., 2011. Evaluation  
392 of toxicity, genotoxicity and environmental risk of simulated textile and tannery  
393 wastewaters with a battery of biotests. *Ecotoxicol Environ Saf.* 74 (4), 866-73.

394 Thomas, O., Theraulaz, F., Agnel, C., Suryani, S., 1996. Advanced UV examination of  
395 wastewater. *Environmental Technology*, 17 (5), 251-261.

396 Tsuda, S., Murakami, M., Matsusaka, N., Kano, K., Taniguchi, K., Sasaki, Y. F., 2001, DNA  
397 damage induced by red food dyes orally administered to pregnant and male mice. *Toxicol*  
398 *Sci.* 61 (1), 92-9.

399 UVPro, 2000. Procédés et dispositifs pour la manipulation, l'archivage, la caractérisation,  
400 l'analyse d'un échantillon liquide. Patent 00402038-4

401 Verma, Y., 2008. Acute toxicity assessment of textile dyes and textile and dyes industrial  
402 effluents using *Daphnia magna* bioassay. *Toxicol. Ind. Health* 24, 491–500.

403 You, S.J, Teng, J.Y., 2009. Performance and dye-degrading bacteria isolation of a hybrid  
404 membrane process. *J Hazard Mater.* 172(1):172-9.

405 Yusuff, R.O., Sonibare, J.A., 2004. Characterization of textile industries effluents in Kaduna  
406 Nigeria and pollution implications, *Global Nest: the Int. J.* (6),3, 212-221.

407

408 **Figure legends**

409

410 **Figure 1:** Induction of estrogenic activities of some textile dyes. Average  $\pm$  standard  
411 deviation (n=3).

412

413 **Figure 2:** Anti-estrogenic activity of textile (A) dyes as measured by the yeast estrogen screen  
414 (B). Competitive binding essay of 3 textile dyes (0.01 g/l) against various concentrations of  
415 E2. Average  $\pm$  standard deviation (n=3).

416

417 **Figure 3:** Estrogenic (A) and anti-estrogenic (B) activity of textile effluent, (C) competitive  
418 binding assay of textile effluent (dilution factor 5-fold) against various concentrations of E2.  
419 Average  $\pm$  standard deviation (n=3).

420

421 **Figure 4:** UV/Visible deconvolution of textile effluent spectrum.

422

423

424

425

426 **Table 1:** Textile dyes analysed in this study. <sup>a</sup>Dyes and informations obtained by Everlight  
 427 Chemical SA.

<i>Compound</i>	<i>CAS #</i>	<i>Family</i>	<i>Formula</i>
Benzopurpurine 4B (Red 4 B)	992-59-6	azo	C <sub>34</sub> H <sub>26</sub> N <sub>6</sub> Na <sub>2</sub> O <sub>6</sub> S <sub>2</sub>
Direct Black VSF AZO-FREE (Direct Black 22)	6473-13-8	azo	C <sub>44</sub> H <sub>32</sub> N <sub>13</sub> Na <sub>3</sub> O <sub>11</sub> S <sub>3</sub>
Yellow 4 G (Solvent yellow 19)	10343-55-2	azo	C <sub>16</sub> H <sub>11</sub> CrN <sub>4</sub> O <sub>8</sub> S
Direct Blue 71 (Blue BRR)	4399-55-7	azo	C <sub>40</sub> H <sub>23</sub> N <sub>7</sub> Na <sub>4</sub> O <sub>13</sub> S <sub>4</sub>
Brown RL	12238-94-7	azo	C <sub>15</sub> H <sub>14</sub> O <sub>5</sub>
red Alpacide 3BL	12238-49-2	azo	C <sub>18</sub> H <sub>10</sub> N <sub>2</sub> O <sub>2</sub> Cl <sub>2</sub>
brown GV			C <sub>31</sub> H <sub>21</sub> N <sub>7</sub> O <sub>6</sub> Na <sub>2</sub> S
Orange 7GL	12222-37-6	azo	C <sub>42</sub> H <sub>28</sub> N <sub>7</sub> Na <sub>4</sub> O <sub>15</sub> S <sub>4</sub>
Direct Black PMSF		azo	C <sub>37</sub> H <sub>25</sub> N <sub>5</sub> Na <sub>2</sub> O <sub>6</sub> S <sub>2</sub>
Everzol Navy Blue FBN <sup>a</sup>	93912-64-2	azo	C <sub>37</sub> H <sub>29</sub> ClN <sub>10</sub> O <sub>22</sub> S <sub>7</sub> Na <sub>6</sub>
Blue HFRL <sup>b</sup>			
Direct Red 89 BNL 200% Blue DERF <sup>b</sup>		azo	
Everzol Yellow ED <sup>a</sup>	Confidential data		
Reactive dyes red 3BS (Red 195)	93050-79-4	azo	C <sub>31</sub> H <sub>19</sub> ClN <sub>7</sub> O <sub>19</sub> S <sub>6</sub>
Everzol Blue ED <sup>a</sup>	2580-78-1	antraquinone	C <sub>22</sub> H <sub>18</sub> N <sub>2</sub> O <sub>11</sub> S <sub>3</sub> Na <sub>2</sub>
Yellow Flavina CXL <sup>b</sup>			
Everzol Navy ED	17095-24-8	azo	C <sub>26</sub> H <sub>25</sub> N <sub>5</sub> O <sub>19</sub> S <sub>6</sub> Na <sub>4</sub>
Solvent yellow 56	2481-94-9	azo	C <sub>16</sub> H <sub>19</sub> N <sub>3</sub>
Yellow 3GF <sup>b</sup>			C <sub>30</sub> H <sub>26</sub> N <sub>4</sub> Na <sub>2</sub> O <sub>8</sub> S <sub>2</sub>
Everzol Red ED <sup>a</sup>	Confidential data		
Grey GGL <sup>b</sup>			
Blue ED 250	89157-03-9	azo	C <sub>31</sub> H <sub>24</sub> ClN <sub>7</sub> O <sub>19</sub> S <sub>6</sub> Na <sub>5</sub>

428 <sup>a</sup>Dyes and informations obtained by Everlight Chemical SA. <sup>b</sup>Reactive azo dye (copper  
 429 complex, 70% pure for Blue HFRL) kindly provided by a textile industry unfortunately there  
 430 is no available information on its structure (Baëta et al. 2011)

431

432 **Table 2:** Relative estrogen agonistic and antagonistic activities of industrial dyes. Estrogenic  
 433 activity of textile dyes (1g/l). The anti-estrogenic activity of textile dyes has been determinate  
 434 at 1g/l and expressed in 4-OHT%. The half maximal effective concentration (AC50) was  
 435 calculated based on the sigmoidal dose–effect curve of 4-OHT.

<i>Compounds</i>	<i>Test Estrogen Screen agonist activity (E2%)</i>	<i>Test Estrogen Screen antagonist activity (4-OHT %)</i>	<i>Test Estrogen Screen antagonist AC 50 (g/l)</i>
E <sub>2</sub> (10 <sup>E-9</sup> M)	100	-	-
OTH	-	100	2*10-5



Benzopurpurine 4B (Red 4 B)	0	100	0.07
Direct Black VSF AZO-FREE (Direct Black 22)	0	20	>1
Yellow 4 G (Solvent yellow 19)	0	13	>1
Direct Blue 71 (Blue BRR)	0	22	>1
*Brown RL	0	-	-
Red Alpacide 3BL	0	44	>1
*Brown GV	0	-	-
*Orange 7GL	0	-	-
*Direct Black PMSF	0	-	-
Everzol Navy Blue FBN	0	94	0.05
Blue HFRL	0	100	0.015
Direct Red 89 BNL 200%	0	100	0.025
Blue DERF	0	10	>1
*Everzol Yellow ED	0	-	-
Reactive dyes red 3BS (Red 195)	22	0	0
Everzol Blue ED	0	35	>1
Yellow Flavina CXL)	36	0	0
Everzol Navy ED	0	32	>1
Solvent yellow 56	11	0	0
Yellow 3GF	0	0	0
Everzol Red ED	0	30	>1
*Grey GGL	0	-	-
Blue ED 250	0	0	0

---

436

437





