

3-months elimination diet in childhood EoE: nutritional and immunological aspects

Diana Colson, Nicolas Kalach, Pascale Soulaines, Bénédicte Michaud,
Lucienne Chatenoud, C Dupont

► **To cite this version:**

Diana Colson, Nicolas Kalach, Pascale Soulaines, Bénédicte Michaud, Lucienne Chatenoud, et al.. 3-months elimination diet in childhood EoE: nutritional and immunological aspects. Food Allergy and Anaphylaxis Meeting (FAAM 2013), Feb 2013, Nice, France. 3 (Suppl 3), pp.P4, 2013. <inserm-00848073>

HAL Id: inserm-00848073

<http://www.hal.inserm.fr/inserm-00848073>

Submitted on 25 Jul 2013

HAL is a multi-disciplinary open access archive for the deposit and dissemination of scientific research documents, whether they are published or not. The documents may come from teaching and research institutions in France or abroad, or from public or private research centers.

L'archive ouverte pluridisciplinaire **HAL**, est destinée au dépôt et à la diffusion de documents scientifiques de niveau recherche, publiés ou non, émanant des établissements d'enseignement et de recherche français ou étrangers, des laboratoires publics ou privés.



POSTER PRESENTATION

Open Access

3-months elimination diet in childhood EoE: nutritional and immunological aspects

D Colson^{1*}, N Kalach², P Soulaines³, B Michaud⁴, L Chatenoud⁵, C Dupont³

From Food Allergy and Anaphylaxis Meeting (FAAM 2013)
Nice, France. 7-9 February 2013

Background

The use of dietary treatment in eosinophilic esophagitis (EoE) in children is regularly described in literature. The aim of the study was to evaluate anthropomorphic and biological parameters in EoE children before and after elimination diet.

Methods

A cohort of 87 consecutive patients with EoE (>15 eosinophils/hpf in esophageal biopsies) was retrospectively analyzed. All children followed the so-called modified six-food elimination diet (SFED), excluding 6 main offending foods retrieved together with those eliciting positive SPT and APT, supplemented with a nutritional support by an amino acid formula. Patients had the following treatment sequence: endoscopy/clinical/biological assessment, 3 months elimination diet and second endoscopy/clinical/ biological assessment.

Results

Analysis included 49 patients. At enrolment, BMI was significantly lower in females than in males ($p < 0.05$), and total circulating IgG and IgM levels were higher in females ($p < 0.05$). Patients having recovered at the end of the study happened to exhibit at enrolment a BMI significantly higher than those having shown only partial or no recovery ($p < 0.05$). Elimination diet led to complete recovery (no symptoms & < 5 eosinophils/hpf) in 53% cases. Following the elimination diet, BMI remained lower in females than in males. Also, blood eosinophils counts ($p < 0.0001$), IgG ($p < 0.003$) and IgM levels ($p < 0.05$) showed a significant decrease as well as specific IgE titers against CMP ($p < 0.02$) and egg ($p < 0.02$), with a tendency for decrease with fish ($p < 0.09$) and wheat ($p < 0.09$).

Conclusion

Female/male differences are observed in EoE: lower BMI, higher risk for malnutrition and more active biological parameters. The elimination diet leads to maintenance of nutritional status, decreased immune activity, identical female/male differences.

Disclosure of interest

D Colson: Grant/research support from Nutritia Nutrition Clinique, N Kalach: None declared, P Soulaines: None declared, B Michaud: None declared, L Chatenoud: None declared, C Dupont: None declared.

Author details

¹Nutritia Nutrition Clinique, Saint Ouen, France. ²Institut Catholique de Lille, Lille, France. ³Hopital Necker Enfants-Malades, Paris, France. ⁴Laboratoire d'Immunologie Biologie Inserm U 1013, Hopital Necker Enfants-Malades, Paris, France. ⁵U1013, Hopital Necker Enfants-Malades, Paris, France.

Published: 25 July 2013

doi:10.1186/2045-7022-3-S3-P4

Cite this article as: Colson et al.: 3-months elimination diet in childhood EoE: nutritional and immunological aspects. *Clinical and Translational Allergy* 2013 **3**(Suppl 3):P4.

¹Nutritia Nutrition Clinique, Saint Ouen, France
Full list of author information is available at the end of the article