

Leaky Ryanodine receptors increases Amyloid-beta load and induces memory impairments in Tg2576 mouse model of Alzheimer disease

Bénédicte Oulès, Dolores Del Prete, Barbara Greco, Xuexin Zhang, Inger Lauritzen, Mohamed Trebak, Fabio Benfenati, Frédéric Checler, Mounia Chami

► **To cite this version:**

Bénédicte Oulès, Dolores Del Prete, Barbara Greco, Xuexin Zhang, Inger Lauritzen, et al.. Leaky Ryanodine receptors increases Amyloid-beta load and induces memory impairments in Tg2576 mouse model of Alzheimer disease. From *Molecular Neurodegeneration: Basic biology and disease pathways*, Sep 2013, Cannes, France. Biomed Central, 8 (Suppl 1), pp.P54, 2013. <inserm-00869877>

HAL Id: inserm-00869877

<http://www.hal.inserm.fr/inserm-00869877>

Submitted on 4 Oct 2013

HAL is a multi-disciplinary open access archive for the deposit and dissemination of scientific research documents, whether they are published or not. The documents may come from teaching and research institutions in France or abroad, or from public or private research centers.

L'archive ouverte pluridisciplinaire **HAL**, est destinée au dépôt et à la diffusion de documents scientifiques de niveau recherche, publiés ou non, émanant des établissements d'enseignement et de recherche français ou étrangers, des laboratoires publics ou privés.

POSTER PRESENTATION

Open Access

Leaky Ryanodine receptors increases Amyloid-beta load and induces memory impairments in Tg2576 mouse model of Alzheimer disease

Bénédicte Oulès^{1*}, Dolores Del Prete^{2,3}, Barbara Greco², Xuexin Zhang⁴, Inger Lauritzen³, Mohamed Trebak⁴, Fabio Benfenati², Frédéric Checler³, Mounia Chami³

From Molecular Neurodegeneration: Basic biology and disease pathways
Cannes, France. 10-12 September 2013

Background

In Alzheimer disease (AD), the perturbation of the endoplasmic reticulum (ER) calcium (Ca²⁺) homeostasis has been linked to presenilins (PS) [1], the catalytic core in gamma-secretase complexes cleaving the amyloid precursor protein (APP) thereby generating amyloid beta peptides. Here we investigate whether APP perturbs ER Ca²⁺ homeostasis and whether ER Ca²⁺ could in turn influence amyloid beta production.

Materials and methods

We used neuroblastoma SH-SY5Y cell lines stably over-expressing human APP695 cDNA harboring or not the Swedish double mutation (K670M→N671L) (APP^{swe} and APP695 respectively) and Tg2576 mice carrying human APP^{swe} cDNA under the control of the hamster Prion promoter. We analyzed subcellular Ca²⁺ signals by using aequorin targeted Ca²⁺ probes, Fura2, AM dye and patch clamp experiments. We analyzed the expression of the Ryanodine receptor (RyR) by western blot and quantitative RT-PCR. We studied the impact of RyR blockade by dantrolene on APP metabolism and Amyloid beta peptide production using biochemistry, ELISA and immunohistochemistry approaches and investigated the underlying mechanisms through the study of APP phosphorylation, and beta- and gamma secretase expression and activities. We finally studied pre- and post-synaptic proteins expression and learning and memory paradigms in Tg2576 mice upon vehicle or dantrolene treatment.

Results

We show that overexpression of APP695, or APP^{swe} triggers increased Ryanodine receptors (RyR) expression and enhances RyR-mediated ER Ca²⁺ release in SH-SY5Y neuroblastoma cells and in APP^{swe}-expressing (Tg2576) mice. Interestingly, dantrolene-induced lowering of RyR-mediated Ca²⁺ release leads to the reduction of both intracellular and extracellular amyloid beta load in neuroblastoma cells as well as in primary cultured neurons derived from Tg2576 mice. This Aβ reduction can be accounted for by decreased Thr-668-dependent APP phosphorylation and β- and gammaoligomers and slows down learning and memory deficits in Tg2576 mice [2].

Conclusions

Overall, our data document a key role of RyR in Aβ production and learning and memory performances, and delineate RyR-mediated control of Ca²⁺ homeostasis as a physiological paradigm that could be targeted for innovative therapeutic approaches.

Acknowledgements

This work was supported by INSERM, CNRS, AFM, FRM, LABEX DISTALZ (excellence laboratory, program investment for the future), the National Institutes of Health, "L'École de l'INSERM Liliane Bettencourt" for supporting the MD-PhD curriculum of B.O., and the Italian Institute of Technology, Genova, Italy.

Authors' details

¹INSERM U 807, Paris V University, F-75015 Paris, France. ²Neuroscience and Brain Technologies - Istituto Italiano di Tecnologia, 16163 Genova, Italy.

³Institut de Pharmacologie Moléculaire et Cellulaire, UMR7275 CNRS/UNSA, 06560 Valbonne, France. ⁴Center for Cardiovascular Sciences, MC8, Albany Medical College, Albany, NY 12208; USA.

Published: 4 October 2013

¹INSERM U 807, Paris V University, F-75015 Paris, France
Full list of author information is available at the end of the article

References

1. Bezprozvanny I, Mattson MP: Neuronal calcium mishandling and the pathogenesis of Alzheimer's disease. *Trends Neurosci* 2008, **31**:454-63.
2. Oules B, Del Prete D, Greco B, Zhang X, Lauritzen I, Sevalle J, Moreno S, Paterlini-Brechot P, Trebak M, Checler F, Benfenati F, Chami M: Ryanodine receptor blockade reduces amyloid-beta load and memory impairments in Tg2576 mouse model of Alzheimer disease. *J Neurosci* 2012, **32**:11820-34.

doi:10.1186/1750-1326-8-S1-P54

Cite this article as: Oulès *et al.*: Leaky Ryanodine receptors increases Amyloid-beta load and induces memory impairments in Tg2576 mouse model of Alzheimer disease. *Molecular Neurodegeneration* 2013 **8**(Suppl 1): P54.

**Submit your next manuscript to BioMed Central
and take full advantage of:**

- Convenient online submission
- Thorough peer review
- No space constraints or color figure charges
- Immediate publication on acceptance
- Inclusion in PubMed, CAS, Scopus and Google Scholar
- Research which is freely available for redistribution

Submit your manuscript at
www.biomedcentral.com/submit

