

## **Interactions between Gastrointestinal Nematodes and Malaria in a Cohort of Children in an Amazonian Village**

Aurélia Stefani, Marie Cheuret, Duc N'guyen, Stéphane Simon, Paul Brousse, Bernard Carne, Mathieu Nacher

► **To cite this version:**

Aurélia Stefani, Marie Cheuret, Duc N'guyen, Stéphane Simon, Paul Brousse, et al.. Interactions between Gastrointestinal Nematodes and Malaria in a Cohort of Children in an Amazonian Village. *Journal of Tropical Pediatrics*, Oxford University Press (OUP), 2016, Epub ahead of print. <10.1093/tropej/fmw063>. <inserm-01410100>

**HAL Id: inserm-01410100**

**<http://www.hal.inserm.fr/inserm-01410100>**

Submitted on 6 Dec 2016

**HAL** is a multi-disciplinary open access archive for the deposit and dissemination of scientific research documents, whether they are published or not. The documents may come from teaching and research institutions in France or abroad, or from public or private research centers.

L'archive ouverte pluridisciplinaire **HAL**, est destinée au dépôt et à la diffusion de documents scientifiques de niveau recherche, publiés ou non, émanant des établissements d'enseignement et de recherche français ou étrangers, des laboratoires publics ou privés.

1 **Interactions between gastro intestinal nematodes and malaria in a cohort of children in an**  
2 **Amazonian village.**

3 Aurélie Stefani<sup>[1],[2]</sup>, Marie Cheuret<sup>[3]</sup>, Duc N'Guyen<sup>[3]</sup>, Stéphane Simon<sup>[1]</sup>, Paul Brousse<sup>[4]</sup>,  
4 Bernard Carme<sup>[1],[2],[5]</sup>, Mathieu Nacher<sup>[1],[2],[5]</sup>.

5

6 1. EPaT team, (EA 3593), UFR de Médecine – Université des Antilles et de la Guyane, Cayenne,  
7 French Guiana.

8 2. STRonGer Programme, Institut Pasteur de la Guyane, Cayenne, French Guiana.

9 3. Laboratoire de Parasitologie-Mycologie, Cayenne General Hospital, Cayenne, French Guiana.

10 4. Département des Centres de Santé, Cayenne General Hospital, Cayenne, French Guiana.

11 5. Centre d'Investigation Clinique (CIC INSERM 1424), Cayenne General Hospital, Cayenne,  
12 French Guiana INSERM.

13 Corresponding Author: Pr Mathieu Nacher, CIC INSERM 1424, Centre Hospitalier de Cayenne,  
14 Avenue des Flamboyants, 97300, Cayenne, French Guiana

15

16

17

18 **Summary**

19 **Introduction.** Most studies on nematode malaria interactions were conducted outside of the Americas.  
20 The objective of the present study was thus to study the relation between malaria and nematodes in a  
21 cohort of children in an Amazonian village.

22 **Methods.** Odds ratios for intestinal nematode infections as an explanatory variable to malaria-resistant  
23 vs. malaria-sensitive were computed.

24 **Results.** *Ascaris lumbricoides* was significantly more frequent in the “resistant” malaria group than in  
25 the “sensitive” one.

26 **Conclusions.** Despite its low statistical power, the present results find that **Ascaris** was associated  
27 with less malaria, as observed by a number of studies.

28

29 **Keywords:** *Plasmodium falciparum*, *Plasmodium vivax*, relapses, GI nematodes, *Ascaris*  
30 *lumbricoides*, French Guiana.

31

32

33

34

## 35 **Introduction**

36 Gastrointestinal nematode infections and malaria have a broadly overlapping distribution. A number  
37 of studies from different continents have shown complex interactions between different GI nematodes  
38 and malaria.

39 Observational studies in Thailand showed that *Ascaris lumbricoides* was associated with a dose-  
40 dependent association with protection from cerebral malaria and acute renal failure <sup>1, 2</sup>. Other studies  
41 have shown that *Ascaris* was associated with lower incidence or prevalence of malaria <sup>3, 4</sup>. In contrast,  
42 it was observed notably in Africa and Madagascar, that hookworm was associated with a greater  
43 incidence of malaria <sup>5-9</sup>.

44 These research questions have received relatively little attention given the omnipresence of  
45 coinfections in tropical regions. The difficulty of this question lies in the different dynamics of  
46 transmission between malaria and different helminthes with different immunomodulatory properties or  
47 hematologic consequences, which complicates the analysis of results. In addition, most results come  
48 from observational studies which are prone to biases and confounding (nutritional status,  
49 socioeconomic level, anemia, background immunity, self-treatment, etc.).

50 In French Guiana, studies performed in 2000 - 2005 ( unpublished data) have shown the persistence of  
51 a high prevalence of gastrointestinal nematode infections among communities living in the interior of  
52 French Guiana, particularly among Amerindian children living along the middle and upper Oyapock  
53 River (Camopi, Trois-Sauts). The most frequent parasite species were *Ascaris lumbricoides*,  
54 *Strongyloides stercoralis*, and *Necator americanus*. These studies have shown a prevalence exceeding  
55 20% for these three nematodes.

56 In parallel, the incidence of malaria was high among children (0-7 years) in Camopi over the 2001-  
57 2009 period with a mean of 238, 514 and 21‰ person-years for *P. falciparum*, *P. vivax* and mixed  
58 infections (microscopic diagnosis), respectively. Finally, a univariate Cox regression analysis showed  
59 that there was a link between anthelmintic treatment and malaria in these children <sup>10</sup>.

60 The trends suggest that while malaria seems less symptomatic among *Ascaris*-infected persons than  
61 those without *Ascaris*, it seems that incidence is higher among patients with hookworm than those  
62 without hookworm. This may result from immune modulation in *Ascaris*-infected patients and  
63 hematologic factors in hookworm-infected patients. Although discernible trends seem to emerge, there  
64 are conflicting results, which are often due to methodological differences. It is impossible to  
65 demonstrate causal relations in observational studies. However, converging elements may point to  
66 causal relations between two variables. To this end repeating the study in different contexts is  
67 important to determine if a finding is robust. Most studies on GI-nematode malaria interactions were  
68 conducted outside of the Americas except two studies in Colombia and Brazil <sup>4,7</sup>. The objective of the  
69 present study was thus to determine whether there were any relation between malaria and GI  
70 nematodes in the context of a cohort of children under seven years of age living in a small Amazonian  
71 village in French Guiana.

## 72 **Methods**

73 The village of Camopi, located in the Oyapock malaria endemic area, consists of a central village and  
74 28 hamlets localized within 15 km<sup>2</sup> along the Oyapock and the Camopi Rivers. The village is isolated  
75 from the coast and separated from Brazil by the Oyapock River, which represents the border. The  
76 1200 inhabitants of Camopi are mainly Wayampi and Teko Amerindians from the linguistic family of  
77 the tupi-guarani. They respectively live on the banks of the Oyapock and the Camopi Rivers. The  
78 population is young with an average age of 18 years.

79 The patients from the Camopi cohort described elsewhere<sup>10, 11</sup> were reviewed and classified in two  
80 groups, one resistant and one sensitive according to the number of malaria episodes. *Plasmodium*  
81 *vivax* relapses were defined as infections occurring within 90 days of a first *P. vivax* episode <sup>12</sup>.

82 Two groups of children were identified regarding their past history of malaria infection in the period  
83 2001-2009: one "sensitive" group had  $\geq 7$  malaria episodes; and one "resistant" group only had  $\leq 1$   
84 within 3 years or  $\leq 2$  within 6 years.

85 Subsequently to this group definition, three field missions were conducted to collect stool samples in  
86 order to test the hypothesis that malaria-resistant patients had a significantly different prevalence of GI  
87 nematodes than the "malaria-sensitive" group.

88 Overall 91 stool samples were collected. All past malaria history, nematodes infection and  
89 anthelmintic treatments were recorded for each child included in the study.

90 Laboratory diagnosis of three helminth infections was performed by multiplex real-time PCR (*Ascaris*  
91 *lumbricoides*, *Necator americanus* or hookworm, and *Strongyloides stercoralis* or threadworm) <sup>13</sup>.

92 Odds ratios for intestinal nematode infections as an explanatory variable to malaria-resistant vs.  
93 malaria-sensitive were computed with Stata 10® software (College Station, Texas).

94 Odds ratios for intestinal nematode infections as an explanatory variable to « relapses » vs. « no  
95 relapses» were also computed. Given the small number of observations multivariate analyses were  
96 not used.

97

## 98 **Results**

99 Overall 86.8% of the children were infected by one nematode: 68.1% (n=62) were infected by  
100 hookworm, 52.7% (n=48) by threadworm and 34.1% (n=31) by *Ascaris*. A total of 65% of the  
101 children had a co-infection.

102 Coinfections between *N. americanus* and *S. stercoralis* were more frequent than coinfections with  
103 *Ascaris*(see Figure 2). This is probably because hookworm and threadworm are present in the same  
104 environmental types due to their similar modes of transmission.

105 Among the 91 children who participated in the study, 84 were included. A total of 41 children were  
106 “malaria sensitive” and 43 of them were “malaria resistant” regarding the chosen group definition.

107 Table 1 shows that *Ascaris* was significantly more frequent in the “resistant” malaria group than in the  
108 “sensitive” one (p=0.003). No association was found between the two other nematodes and malaria.

109 Among the 57 children that had at least one *P. vivax* episode, 41 had one or more relapses whereas 16  
110 children had no relapses. *Ascaris* was more frequent in children that had no relapses but this failed to  
111 reach statistical significance. Table 1 shows that relapses seemed less frequent in the *Ascaris* group.  
112 The difference was not significant but the power to detect such a difference given the sample size was  
113 only 30%.

## 114 **Discussion**

115 Despite its very low sample size, the present study, as a number of other studies<sup>1, 3, 4, 7, 14</sup> reviewed<sup>15</sup>,  
116 has found a negative association between *Ascaris* and malaria in this Amazonian setting. To explain  
117 the proximal explanation of these observations between *Ascaris* and malaria, complementary  
118 immunological hypotheses have been put forward{Nacher, 2002 #40}. This reinforces the suggestion  
119 that *Ascaris* has a protective impact on both the severity and patency of clinical infections. *Ascaris*  
120 *lumbricoides* has often been singled out as the most significant worm presumably because it also  
121 represents the one with the largest biomass of antigenic immunomodulatory material. In addition,  
122 *Ascaris* antigens have been reported to have a particular ability to induce a strong IgE response.  
123 Hypothetical ultimate causes lie in the mutual benefits for the worm and malaria parasites to protect  
124 their host in order to reproduce more effectively. Patients co-infected with malaria and a nematode  
125 seem to have more gametocytes, fewer symptoms associated with malaria and infection of longer  
126 duration. The anemic host has an increased attractiveness of the host for the vectors presumably  
127 leading to an increase of the number of mosquito bites. This phenomenon could affect incidence and  
128 transmission.

129 Although the results were not statistically significant at the 5% level, there was a trend for decreased  
130 relapses in *Ascaris*-infected patients. Surely, statistical power was much too low (28%). The  
131 hypothesis is again that *Ascaris*-mediated immunomodulation may interfere with the activation of  
132 latent hypnozoites leading to the *vivax* relapses. This had never been shown before. Given the novelty  
133 and implications of such an observation, larger studies should test this hypothesis.

134

135 The weakness of the present study was its sample size. However, despite its low statistical power the  
136 present results also find that *Ascaris* was associated with less malaria, as observed by a number of  
137 studies. An interesting corollary finding was that there also seemed to be fewer *P.vivax* relapses in the  
138 *Ascaris* group which would be a novel observation with intriguing implications on the activation of  
139 latent hypnozoites. However, this observation needs to be replicated with larger sample sizes. This  
140 data from South America brings additional data to further improve the understanding of nematodes-  
141 *Plasmodium* coinfections.

142

143

144

145 **Conflict of interest statement:** The authors declare that there is no conflict of interest.

146

147

## 148 **References**

149

- 150 1. Nacher M, Gay F, Singhasivanon P, et al. *Ascaris lumbricoides* infection is associated with  
151 protection from cerebral malaria. *Parasite immunology* 2000; **22**(3): 107-13.
- 152 2. Nacher M, Singhasivanon P, Silachamroon U, et al. Helminth infections are associated with  
153 protection from malaria-related acute renal failure and jaundice in Thailand. *The American journal of*  
154 *tropical medicine and hygiene* 2001; **65**(6): 834-6.
- 155 3. Brutus L, Watier L, Hanitrasoamampionona V, Razanatsoarilala H, Cot M. Confirmation of the  
156 protective effect of *Ascaris lumbricoides* on *Plasmodium falciparum* infection: results of a  
157 randomized trial in Madagascar. *The American journal of tropical medicine and hygiene* 2007; **77**(6):  
158 1091-5.
- 159 4. Melo GC, Reyes-Lecca RC, Vitor-Silva S, et al. Concurrent helminthic infection protects  
160 schoolchildren with *Plasmodium vivax* from anemia. *PloS one* 2010; **5**(6): e11206.
- 161 5. Boel M, Carrara VI, Rijken M, et al. Complex Interactions between soil-transmitted helminths  
162 and malaria in pregnant women on the Thai-Burmese border. *PLoS neglected tropical diseases* 2010;  
163 **4**(11): e887.
- 164 6. Degarege A, Animut A, Legesse M, Erko B. Malaria severity status in patients with soil-  
165 transmitted helminth infections. *Acta tropica* 2009; **112**(1): 8-11.



- 166 7. Fernandez-Nino JA, Idrovo AJ, Cucunuba ZM, et al. Paradoxical associations between soil-  
167 transmitted helminths and Plasmodium falciparum infection. *Transactions of the Royal Society of*  
168 *Tropical Medicine and Hygiene* 2012; **106**(11): 701-8.
- 169 8. Hillier SD, Booth M, Muhangi L, et al. Plasmodium falciparum and helminth coinfection in a  
170 semi urban population of pregnant women in Uganda. *The Journal of infectious diseases* 2008;  
171 **198**(6): 920-7.
- 172 9. Pullan RL, Kabatereine NB, Bukirwa H, Staedke SG, Brooker S. Heterogeneities and  
173 consequences of Plasmodium species and hookworm coinfection: a population based study in  
174 Uganda. *The Journal of infectious diseases* 2011; **203**(3): 406-17.
- 175 10. Stefani A, Hanf M, Nacher M, Girod R, Carme B. Environmental, entomological,  
176 socioeconomic and behavioural risk factors for malaria attacks in Amerindian children of Camopi,  
177 French Guiana. *Malaria journal* 2011; **10**: 246.
- 178 11. Hustache S, Nacher M, Djossou F, Carme B. Malaria risk factors in Amerindian children in  
179 French Guiana. *The American journal of tropical medicine and hygiene* 2007; **76**(4): 619-25.
- 180 12. Hanf M, Stephani A, Basurko C, Nacher M, Carme B. Determination of the Plasmodium vivax  
181 relapse pattern in Camopi, French Guiana. *Malaria journal* 2009; **8**: 278.
- 182 13. Basuni M, Muhi J, Othman N, et al. A pentaplex real-time polymerase chain reaction assay for  
183 detection of four species of soil-transmitted helminths. *The American journal of tropical medicine*  
184 *and hygiene* 2011; **84**(2): 338-43.
- 185 14. Murray MJ, Murray AB, Murray MB, Murray CJ. Parotid enlargement, forehead edema, and  
186 suppression of malaria as nutritional consequences of ascariasis. *The American journal of clinical*  
187 *nutrition* 1977; **30**(12): 2117-21.
- 188 15. Nacher M. Interactions between worms and malaria: good worms or bad worms? *Malaria*  
189 *journal* 2011; **10**: 259.

190