



**HAL**  
open science

## Nanopores: maltoporin channel as a sensor for maltodextrin and lambda-phage.

Emir Berkane, Frank Orlik, Alain Charbit, Christophe Danelon, Didier Fournier, Roland Benz, Mathias Winterhalter

### ► To cite this version:

Emir Berkane, Frank Orlik, Alain Charbit, Christophe Danelon, Didier Fournier, et al.. Nanopores: maltoporin channel as a sensor for maltodextrin and lambda-phage.. *Journal of Nanobiotechnology*, BioMed Central, 2005, 3, pp.3. 10.1186/1477-3155-3-3 . inserm-00092492

**HAL Id: inserm-00092492**

**<https://www.hal.inserm.fr/inserm-00092492>**

Submitted on 12 Sep 2006

**HAL** is a multi-disciplinary open access archive for the deposit and dissemination of scientific research documents, whether they are published or not. The documents may come from teaching and research institutions in France or abroad, or from public or private research centers.

L'archive ouverte pluridisciplinaire **HAL**, est destinée au dépôt et à la diffusion de documents scientifiques de niveau recherche, publiés ou non, émanant des établissements d'enseignement et de recherche français ou étrangers, des laboratoires publics ou privés.

Research

Open Access

## Nanopores: maltoporin channel as a sensor for maltodextrin and lambda-phage

E Berkane<sup>1,2</sup>, F Orlik<sup>2</sup>, A Charbit<sup>3</sup>, C Danelon<sup>1</sup>, D Fournier<sup>1</sup>, R Benz<sup>2</sup> and M Winterhalter\*<sup>1,4</sup>

Address: <sup>1</sup>Institut Pharmacologie & Biologie Structurale-CNRS UMR5089, 205, rte de Narbonne, F-31077 Toulouse, France, <sup>2</sup>Lehrstuhl für Biotechnologie, Biozentrum, Am Hubland, D-97074 Würzburg, Germany, <sup>3</sup>Inserm U-570, CHU Necker-Enfants Malades, 156, rue de Vaugirard, F- 75730 Paris Cedex 15, France and <sup>4</sup>International University Bremen, School of Engineering and Science, D-28727 Bremen, Germany

Email: E Berkane - emirrt@yahoo.fr; F Orlik - orlik@biozentrum.uni-wuerzburg.de; A Charbit - charbit@necker.fr; C Danelon - Christophe.Danelon@epfl.ch; D Fournier - fournier@ipbs.fr; R Benz - roland.benz@mail.uni-wuerzburg.de; M Winterhalter\* - m.winterhalter@iu-bremen.de

\* Corresponding author

Published: 02 March 2005

Received: 18 September 2004

*Journal of Nanobiotechnology* 2005, **3**:3 doi:10.1186/1477-3155-3-3

Accepted: 02 March 2005

This article is available from: <http://www.jnanobiotechnology.com/content/3/1/3>

© 2005 Berkane et al; licensee BioMed Central Ltd.

This is an Open Access article distributed under the terms of the Creative Commons Attribution License (<http://creativecommons.org/licenses/by/2.0>), which permits unrestricted use, distribution, and reproduction in any medium, provided the original work is properly cited.

### Abstract

**Background:** To harvest nutrition from the outside bacteria e.g. *E. coli* developed in the outer cell wall a number of sophisticated channels called porins. One of them, maltoporin, is a passive specific channel for the maltodextrin uptake. This channel was also named LamB as the bacterial virus phage Lambda mis-uses this channel to recognise the bacteria. The first step is a reversible binding followed after a lag phase by DNA injection. To date little is known about the binding capacity and less on the DNA injection mechanism. To elucidate the mechanism and to show the sensitivity of our method we reconstituted maltoporin in planar lipid membranes. Application of an external transmembrane electric field causes an ion current across the channel. Maltoporin channel diameter is around a few Angstrom. At this size the ion current is extremely sensitive to any modification of the channels surface. Protein conformational changes, substrate binding etc will cause fluctuations reflecting the molecular interactions with the channel wall. The recent improvement in ion current fluctuation analysis allows now studying the interaction of solutes with the channel on a single molecular level.

**Results:** We could demonstrate the asymmetry of the bacterial phage Lambda binding to its natural receptor maltoporin.

**Conclusion:** We suggest that this type of measurement can be used as a new type of biosensors.

Nature created and optimized proteins for specific tasks which makes them often interesting in material science. For example, membrane transporters could control the permeability of artificial nanometer sized container. A

typical application could be to control the enzymatic activity in a liposome [1]. Another possible application is to reconstitute channels into planar lipid bilayer and use time dependent conductance as a signal [2,3]. Application

of an external electric field drives the ions through the nano (and subnano) meter sized channel. Any larger molecule that diffuses into and temporarily sticks to the channel interior will cause typical fluctuations of the ion current which allow to conclude on its mode of translocation. Such studies were used to follow sugar translocation through maltoporin [4]. Similar types of measurements were done to investigate the translocation of antibiotics like ampicillin [5]. Subtle changes in the channel size or small conformational changes can be recorded and this technique could be developed towards an instrument to probe very soft forces.

Porins are attractive candidates for applications because they are very stable. Moreover, recombinant technology permits production of porins in *E. coli* with high yields [6]. A third advantage is the availability of the high resolution 3-D crystal structure showing details of substrate binding sites which facilitates enormously a rational engineering of modified proteins.

The outer cell wall of Gram-negative bacteria from *E. coli* is fairly permeable to smaller solutes below a molecular weight of about 400 Da [6]. Such substances can freely permeate under a concentration gradient through general diffusion porins in the outer cell wall. Under stress, e.g. in case of lack of nutrition, the pure diffusion process is too slow and the bacteria need to improve the efficiency of the translocation. For those cases, nature has created a series of rather specific and highly sophisticated membrane channels. The most extensively studied examples of specific porins are the maltooligosaccharide-specific channel Maltoporin of *E. coli* [4,7,8]. Maltoporin forms ion-conducting channels when reconstituted into lipid bilayers [9,10]. The 3D structure of Maltoporin revealed that the monomer of Maltoporin of *E. coli* consists of an 18 stranded  $\beta$ -barrel with short turns at the periplasmic side and large irregular loops at the outside of the cell [11].


The bacteriophage Lambda is a virus recognizing Maltoporin at the outer cell surface [12]. In absence of this membrane channel, phage Lambda does not recognize the bacteria. Or, even minor mutations allow the bacteria to defend themselves against virus attacks. The virus itself can, in turn mutate to restore binding ability. According to the high resolution X-ray structure the water filled channel is far too small to permit the translocation of the double strain DNA (about 20 Å) [11]. The infection mechanism thus must involve one of the following processes: Phage binding will cause a strong conformational change within the Maltoporin or, after binding the phage releases a DNA translocation machinery to bring its DNA across the hydrophobic membrane. To date none of these intermediate steps has been observed so far and the underlying process remains unclear. Recently, gpJ, a protein in the

phage terminal was identified to be involved in the Maltoporin recognition process [13].

A typical set-up for conductance measurements is shown in figure 1. The measurement cell consists of two chambers separated by a hole (less than 0.1 mm diameter) in a thin poly(tetrafluorethylene) film sandwiched between two half-cells made of Teflon (Goodfellow, Cambridge, UK). Prior to each measurement this hole has to be pre-treated to render it lipophilic by coating it with a hexadecane/hexane (1:100 v:v) droplet. After allowing for hexane evaporation, each chamber is filled with 1.5 ml buffer (for example, 1 M KCl, unbuffered, about pH 6). Black lipid bilayers were formed according to the classical Montal-Mueller technique by spreading lipids in hexane/chloroform (9:1) across the aqueous buffer [14]. For sake of stability we used diphytanoyl-phosphocholine (DphPC, Avanti Polar Lipids). After 20 min allowing for evaporation, the buffer level is lowered below the hole level and rose again. Typically after the first or second trial a stable unilamellar membrane is formed. In order to insert single porin trimers within reasonable time, but to avoid insertion of their multiples, a careful balance between the concentration of the protein solution, detergent concentration and buffer volume has to be found. One single porin trimer has to find the membrane and to insert while all others must be inactivated, e.g. by precipitation. Maltoporin from the stock (1 mg/ml in 1% OPOE) was diluted  $10^2$ - $10^5$  times in the buffer containing 1% OPOE. From our own experience in our laboratory the insertion was optimal if smallest amounts (less than 1  $\mu$ l) were injected. In a second measurement we used painted membranes as described previously [15]. Here the Teflon chamber consists of a larger hole (diameter 800  $\mu$ m and larger). Membranes were formed by painting 1  $\mu$ l of a 1% solution of DphPC in n-decane across the hole. This type of membrane facilitates multichannel insertion.

Membrane current was measured via homemade Ag/AgCl electrodes. One electrode was used as ground and the other connected to the headstage of an Axopatch 200B amplifier (Axon Instruments, USA), allowing the application of adjustable potentials (typically, 100 mV) across the membrane. A similar set-up was used in the second measurement.

We recently investigated the sugar penetration on a single molecular level [4]. We were able to reconstitute a single Maltoporin trimer into the lipid bilayer. Addition of sugar into the bulk phase resulted in a blocking of the channel in a concentration dependent manner. At low sugar concentration individual closure of the channel could be observed. Maltohexaose induces higher frequencies of closure and longer closing times than a smaller sugar like maltose. The analysis of the time-resolved conductance as




**Figure 1**

Schematic representation of a typical planar bilayer set-up for ion current recording. 1.a) Two half cells made of Delrine separated by a 25 μm Teflon foil with a hole in the center. Both parts are clamped together. 1.b) Below a microscope picture of the Teflon septum containing a hole. 1.c) Schema of a lipid bilayer with a reconstituted trimeric porin. The Cl<sup>-</sup> ions are attracted to the positive electrode and K<sup>+</sup> to the negative one. Ions are permeating the channel in the MHz range which is beyond the current time resolution. 1.d) The insertion of a single channel will give rise to a jump in conductance. Any objects diffusing in the channel may reduce the permeation time of ions and may be detected either in conductance fluctuations or an averaged reduced conductance.


a function of sugar concentration yielded the binding constant as well as the "on" and "off" rates for the sugar binding. Here we used a modified sugar through covalent binding of an ANDS (3-amino-naphtalene-2,7-disulfonic acid) molecule to the reducing end of a Maltoheptaose as schematically shown in fig. 2A (for details, see [16]). The crystal structure suggests that the maltose molecule enter the channel only with the nonreducing end from the out-

side (or reducing end from the periplasmic side). Subsequently this molecule can only enter from the cis-side in our setup. In fig. 2B we see that addition on the periplasmic side (trans side) inhibit the entry whereas addition to the outer side (cis side) caused blocking. A good control experiment in order to test the activity is to add unmodified sugar molecules to the previous experiment. In fig. 2C we clearly observe the ability to translocate unmodified



**Figure 2**

Typical recordings of ion current through a single Maltoporin trimer in presence of modified maltohexaose (see [16] for details). **(A)** Shows the unmodified maltohexaose and on the right hand side the modified sugar molecule. We designed this molecule according the crystal structure to guarantee the low penetration ability from one side. **(B)** M6-ANDS was added to *trans* (left) and then to *cis* (right). Sugar analogue modulates ion current only to the *cis*-side, the side of Maltoporin addition. The average residence time is 5.0 ms. **(C)** First, M6-ANDS was injected to the *trans*-side and no variation in the ion current occurs. As control, maltohexaose was added to the same side (left). The natural substrate is translocated demonstrating that it enters the channel from *trans* with the reducing end first. Then, M6-ANDS was added additionally to the *cis*-side (right) generating long current interruptions superimposed to maltohexaose blockade events seen in the figure of the left side. The dashed lines corresponding to zero current. Membrane bathing solution was 1 M KCl, 10 mM Tris, 1 mM CaCl<sub>2</sub>, pH 7.4, the applied voltage was + 150 mV.



**Figure 3**

Here we show the ability to recognize bacterial phage Lambda by blocking the ion conductance through the natural receptor Maltoporin. We first reconstituted about 300 Maltoporin channel in a solvent containing planar lipid bilayer. This leads to a stable conductance after about 30 min with no further protein insertion. Titration of 7 and 42 nM of the fusion protein MBP-gpJ from the bacterial virus Lambda to the compartment corresponding the intracellular side of the channel showed no effect. However, titration to the opposite side corresponding to the extracellular side caused a significant reduction of the ion conductance. Membrane bathing solution was unbuffered 1 M KCl giving a pH of about 6. The applied voltage was + 20 mV.

sugars. Addition of small amounts of unmodified sugar to the trans-side caused the expected number of events. Further addition of unmodified sugar to the opposite site enhances the sugar induced blocking. These data can be used for a fundamental analysis to probe e.g. the individual energy barrier and it seems that nature has optimized this channel to have the best turnover number. On the other hand these channels can potentially serve to discriminate sucrose from maltose.

In a second series of experiments we were interested to probe for Lambda phage binding. In principle this should be possible despite the enormous size (about 100 nm size in comparison to 4 nm sized channels). However in a preliminary step we have produced larger quantities of the phage endterminal protein gpJ fused to Maltose Binding Protein (MBP). We reconstituted a larger number of maltoporin in solvent containing membranes and titrated small quantities of the fusion construct MBP-gpJ. We know from the experiments described above that most of the channels are oriented the same direction during the reconstitution. In fig. 3 we show a first result that titration of gpJ to the opposite side of protein addition had no

effect. In contrast, addition of gpJ to the side of porin addition caused rapid blocking of the channel. This observation suggest that the porin inserts with the short turns first and that the protein part exposed to the extracellular side is naturally accessible to Lambda phages. These first results are promising and we currently work on improving the resolution. Here we have to note that this observation is in clear contrast by a report on phage lambda binding in a multichannel preparation [17]. The origin of this discrepancy might be simultaneous multiple insertion. Our observation here is in agreement with other reports showing the same orientation [4,5,18]. However, reason why porins inserts in artificial membranes differently than in natural ones remains unclear. One may speculate that the strong asymmetry of natural membranes or unknown chaperons will facilitate the entry with the long loops first.

Sensing with membrane channel is a new way in screening for solute molecules and several promising examples are already shown [2,3,16,19,20]. The actual bottleneck is the complexity in membrane channel assembly. However, the current development in automatized patch-clamping will open a wide range of possibilities [21,22]. We plan to

reduce the volume on each side of the membrane and the size of the lipid patch. We currently work with pore diameters of about 1  $\mu\text{m}$  with less background capacitance and thus a better time resolution and to simplify the channel assembly.

## References

- Colletier J-P, Chaize B, Winterhalter M, Fournier D: **Protein encapsulation in liposomes: efficiency depends on interactions between protein and phospholipid bilayer.** *BMC Biotechnology* 2002, **2**:9-17.
- Howorka S, Nam J, Bayley H, Kahne D: **Stochastic detection of monovalent and bivalent protein-ligand interactions.** *Angew Chemie Int Ed* 2004, **43**:842-846.
- Bezrukov SM: **Ion channels as molecular coulter counters to probe metabolite transport.** *J Membr Biol* 2000, **174**:1-13.
- Kullman L, Winterhalter M, Bezrukov SM: **Transport of maltodextrins through maltoporin: A single-channel study.** *Biophys J* 2002, **82**:803-812.
- Nestorovich EM, Danelon C, Winterhalter M, Bezrukov SM: **Designed to penetrate: Time-resolved interaction of single antibiotic molecules with bacterial pores.** *Proc Natl Acad Sci (USA)* 2002, **99**:9789-94.
- Van Gelder P, Dumas F, Winterhalter M: **Understanding the function of bacterial outer membrane channels by reconstitution into black lipid membranes.** *Biophys Chem* 2000, **85**:153-67.
- Szmelcman S, Hofnung M: **Maltose transport in *Escherichia coli* K12. Involvement of the bacteriophage lambda receptor.** *J Bacteriol* 1975, **124**:112-118.
- Schmid K, Ebner R, Altenbuchner J, Schmitt R, Lengeler JW: **Plasmid-mediated sucrose metabolism in *Escherichia coli* K12: mapping of the *scr* genes of pUR400.** *Mol Microbiol* 1988, **2**:1-8.
- Szmelcman S, Schwartz M, Silhavy TJ, Boos W: **Maltose transport in *Escherichia coli* K-12: a comparison of transport kinetics in wild-type and lambda-resistant mutants with the dissociation constant of the maltose-binding protein as measured by fluorescent quenching.** *Eur J Biochem* 1976, **65**:13-19.
- Benz R, Schmid A, Nakae T, Vos-Scheperkeuter GH: **Pore formation by LamB of *Escherichia coli* in lipid bilayer membranes.** *J Bacteriol* 1986, **165**:978-986.
- Dutzler R, Wang Y-F, Rizkallah PJ, Rosenbusch JP, Schirmer T: **Crystal structures of various maltooligosaccharides bound to Maltoporin reveal a specific sugar translocation pathway.** *Structure* 1996, **4**:127-134.
- Roa M, Scandella D: **Multiple steps during the interaction between coliphage lambda and its receptor protein in vitro.** *Virology* 1976, **72**:182-194.
- Wang J, Hofnung M, Charbit A: **The C-terminal portion of the tail fiber protein of bacteriophage lambda is responsible for binding to LamB, its receptor at the surface of *Escherichia coli* K-12.** *J Bact* 2000, **182**:508-512.
- Montal M, Mueller P: **Formation of bimolecular membranes from lipid monolayers and a study of their electrical properties.** *Proc Nat Acad Sci (USA)* 1972, **69**:3561-3566.
- Benz R, Janko K, Boos W, Luger P: **Formation of large, ion permeable membrane channels by matrix protein (porin) of *Escherichia coli*.** *Biochem Biophys Acta* 1978, **511**:305-319.
- Danelon C, Brando T, Winterhalter M: **Probing the orientation of reconstituted Maltoporin channels at the single-protein level.** *J Biol Chem* 2003, **278**:35542-51.
- Van Gelder P, Dumas F, Rosenbusch J, Winterhalter M: **Oriented channels reveal asymmetric energy barriers for sugar translocation through maltoporin of *Escherichia coli*.** *Eur J Biochem* 2000, **267**:79-84.
- Andersen C, Schiffler B, Charbit A, Benz R: **PH-induced collapse of the extracellular loops closes *Escherichia coli* maltoporin and allows the study of asymmetric sugar binding.** *J Biol Chem* 2002, **277**:41318-25.
- Akeson M, Branton D, Kasianowicz JJ, Brandin E, Deamer DW: **Microsecond time-scale discrimination among polycytidylic acid, polyadenylic acid, and polyuridylic acid as homopolymers or as segments within single RNA molecules.** *Biophys J* 1999, **77**:3227-33.
- Kasianowicz JJ, Burden DL, Han LC, Cheley S, Bayley H: **Genetically engineered metal ion binding sites on the outside of a channel's transmembrane beta-barrel.** *Biophys J* 1999, **76**:837-45.
- Bruggemann A, George M, Klau M, Beckler M, Steindl J, Behrends JC, Fertig N: **High quality ion channel analysis on a chip with the NPC technology.** *Assay Drug Dev Technol* 2003, **1**:665-673.
- Schmidt C, Mayer M, Vogel H: **A Chip-Based Biosensor for the Functional Analysis of Single Ion Channels.** *Angew Chem Int Ed Engl* 2000, **39**:3137-3140.

Publish with **BioMed Central** and every scientist can read your work free of charge

"BioMed Central will be the most significant development for disseminating the results of biomedical research in our lifetime."

Sir Paul Nurse, Cancer Research UK

Your research papers will be:

- available free of charge to the entire biomedical community
- peer reviewed and published immediately upon acceptance
- cited in PubMed and archived on PubMed Central
- yours — you keep the copyright

Submit your manuscript here:  
[http://www.biomedcentral.com/info/publishing\\_adv.asp](http://www.biomedcentral.com/info/publishing_adv.asp)

