

Poly(ADP-ribose) polymerase (PARP-1) is not involved in DNA double-strand break recovery.

Georges Noël, Nicole Giocanti, Marie Fernet, Frédérique Mégnin-Chanet,
Vincent Favaudon

► **To cite this version:**

Georges Noël, Nicole Giocanti, Marie Fernet, Frédérique Mégnin-Chanet, Vincent Favaudon. Poly(ADP-ribose) polymerase (PARP-1) is not involved in DNA double-strand break recovery.. BMC Cell Biology, BioMed Central, 2003, 4, pp.7. 10.1186/1471-2121-4-7 . inserm-00115574

HAL Id: inserm-00115574

<https://www.hal.inserm.fr/inserm-00115574>

Submitted on 6 Dec 2006

HAL is a multi-disciplinary open access archive for the deposit and dissemination of scientific research documents, whether they are published or not. The documents may come from teaching and research institutions in France or abroad, or from public or private research centers.

L'archive ouverte pluridisciplinaire **HAL**, est destinée au dépôt et à la diffusion de documents scientifiques de niveau recherche, publiés ou non, émanant des établissements d'enseignement et de recherche français ou étrangers, des laboratoires publics ou privés.

Research article

Open Access

Poly(ADP-ribose) polymerase (PARP-1) is not involved in DNA double-strand break recovery

Georges Noël^{1,2}, Nicole Giocanti¹, Marie Fernet^{1,3}, Frédérique Mégnin-Chanet¹ and Vincent Favaudon*¹

Address: ¹Unité 350 INSERM, Institut Curie-Recherche, Bâts. 110-112, Centre Universitaire, 91405 Orsay Cedex, France, ²Centre de Protonthérapie d'Orsay, Bât. 101, Centre Universitaire, BP 65, 91402 Orsay Cedex, France and ³Present address: DNA Repair Group, International Agency for Research on Cancer, 150 cours Albert Thomas, 69372 Lyon Cedex 08, France

Email: Georges Noël - noel@ipno.in2p3.fr; Nicole Giocanti - nicole.giocanti@curie.u-psud.fr; Marie Fernet - fernet@iarc.fr; Frédérique Mégnin-Chanet - frederique.megnin@curie.u-psud.fr; Vincent Favaudon* - vincent.favaudon@curie.u-psud.fr

* Corresponding author

Published: 16 July 2003

Received: 19 February 2003

BMC Cell Biology 2003, 4:7

Accepted: 16 July 2003

This article is available from: <http://www.biomedcentral.com/1471-2121/4/7>

© 2003 Noël et al; licensee BioMed Central Ltd. This is an Open Access article: verbatim copying and redistribution of this article are permitted in all media for any purpose, provided this notice is preserved along with the article's original URL.

Abstract

Background: The cytotoxicity and the rejoining of DNA double-strand breaks induced by γ -rays, H_2O_2 and neocarzinostatin, were investigated in normal and PARP-1 knockout mouse 3T3 fibroblasts to determine the role of poly(ADP-ribose) polymerase (PARP-1) in DNA double-strand break repair.

Results: PARP-1^{-/-} were considerably more sensitive than PARP-1^{+/+} 3T3s to induced cell kill by γ -rays and H_2O_2 . However, the two cell lines did not show any significant difference in the susceptibility to neocarzinostatin below 1.5 nM drug. Restoration of PARP-1 expression in PARP-1^{-/-} 3T3s by retroviral transfection of the full PARP-1 cDNA did not induce any change in neocarzinostatin response. Moreover the incidence and the rejoining kinetics of neocarzinostatin-induced DNA double-strand breaks were identical in PARP-1^{+/+} and PARP-1^{-/-} 3T3s. Poly(ADP-ribose) synthesis following γ -rays and H_2O_2 was observed in PARP-1-proficient cells only. In contrast neocarzinostatin, even at supra-lethal concentration, was unable to initiate PARP-1 activation yet it induced H2AX histone phosphorylation in both PARP1^{+/+} and PARP-1^{-/-} 3T3s as efficiently as γ -rays and H_2O_2 .

Conclusions: The results show that PARP-1 is not a major determinant of DNA double-strand break recovery with either strand break rejoining or cell survival as an endpoint. Even though both PARP-1 and ATM activation are major determinants of the cell response to γ -rays and H_2O_2 , data suggest that PARP-1-dependent poly(ADP-ribose) synthesis and ATM-dependent H2AX phosphorylation, are not inter-related in the repair pathway of neocarzinostatin-induced DNA double-strand breaks.

Background

Ionizing radiation induces multiple lesions in cell DNA including oxidative base damage, single-strand breaks (SSB) and double-strand breaks (DSB) in proportion to

the radiation dose. Among the enzymes that have evolved for the repair of radiation damage, poly(ADP-ribose) polymerase (PARP-1), Ataxia mutated kinase (ATM) and the heterotrimeric DNA-dependent protein kinase (DNA-

PK), play a leading role. ATM is rapidly activated by DSB to phosphorylate proteins in chromatin, most notably H2AX histone [1,2] and the catalytic subunit of the DNA-PK complex (DNA-PKcs). ATM is also essential to coupling of DNA damage detection to NF- κ B activation and proper management of the oxidative stress inherent in radiation exposure. DNA-PKcs is recruited by the Ku70-Ku86 complex at sites of DSB [3,4]. Activated DNA-PKcs phosphorylates a range of protein substrates and, along with XRCC4 and Ligase 4 is essential to V(D)J recombination and DSB repair through the non-homologous end joining (NHEJ) pathway [5].

PARP-1, an ubiquitous 113 kDa enzyme, is also required for the detection and signalization of DNA strand interruptions. It is involved in early DNA damage recognition [6], base excision repair [7,8] and genome surveillance in a variety of situations (for a review see [9]). Activated PARP-1 carries out synthesis and transfer of long linear or branched ADP-ribose polymers (pADPr) to carboxyl groups in a limited number of nuclear protein acceptors, including PARP-1 itself in a reaction initiated by PARP-1 binding to SSB [10]. In addition, several DNA damage signaling or repair proteins possess high-affinity binding motifs for pADPr, among others XRCC1, DNA ligase III, p21^{Waf1} and p53 [11,12]. Two subunits of the DNA-PK heterotrimer, namely, Ku70 and DNA-PKcs also present high affinity motifs for pADPr binding [12], and PARP-1 co-immunoprecipitates with these proteins [13–15]. *In vitro*, the DNA-PKcs subunit can be ADP-ribosylated and stimulated by PARP-1; PARP-1 can in turn be phosphorylated by DNA-PKcs [16].

Whether these protein complexes and post-translational modifications play a role in repair from radioinduced DSB, has been challenged recently [17,18]. In the prospect of unravelling this question, we reasoned that radiomimetic compounds acting to produce DSB with high selectivity and efficiency in the DNA of target cells should be used instead of ionizing radiation, since radiation generates oxidative stress response and elicits a large spectrum of lesions located at random in chromatin.

We chose neocarzinostatin (NCS) for this purpose. NCS is the prototype of the "protein antibiotic" family. It is a complex consisting of a dodecadiyne antibiotic (NCS_{Chrom}) reversibly bound to a carrier protein [19]. NCS is active in the nanomolar range, and NCS_{Chrom} cleaves DNA in the course of a suicide reaction leaving no residual active drug after a few minutes incubation. The major DNA lesions induced by NCS_{Chrom} in DNA result from radical attack [20] and consist of a blunt end break bearing a thymidine-5'-aldehyde residue on one strand [21], with an atypical abasic site at two nucleotide interval on the complementary strand [22,23]. This abasic site is

substrate for endonuclease III [24] in such a way that NCS-induced damage is rapidly converted into DSB in living cells [25,26]. *E. coli* [27,28], yeast [29] or mammalian cells [30–37] bearing a defect in DSB repair, are consistently hypersensitive to induced cell kill by NCS.

PARP-1 proficient (PARP-1^{+/+}) and PARP-1 knockout (PARP-1^{-/-}) 3T3 fibroblasts from syngenic mice, were used to investigate the role of PARP-1 in the recovery of NCS-induced DSB. PARP-1^{-/-} 3T3s complemented with the full PARP-1 cDNA, were also used in this assay. The incidence and repair kinetics of DSB were measured in parallel in PARP-1^{+/+} and PARP-1^{-/-} 3T3s, and the effect of 4-amino-1,8-naphthalimide, a potent PARP-1 inhibitor, was established in PARP-1^{+/+} 3T3s. The cytotoxic effect of γ -rays and H₂O₂ was determined for comparison.

Immunofluorescence studies were also performed to assess pADPr synthesis and H2AX histone phosphorylation following exposure to NCS, γ -rays or H₂O₂. The results show that PARP-1 is likely not to play a crucial role in the repair of DSB, at least in the absence of other types of DNA damage.

Results

Cytotoxicity of NCS, γ -rays and H₂O₂

PARP-1^{+/+} and PARP-1^{-/-} 3T3s were exposed to increasing concentrations of NCS and cytotoxicity was determined through a growth assay (see Methods). Below 1.5 nM NCS PARP-1^{+/+} and PARP-1^{-/-} 3T3s demonstrated exactly the same susceptibility to the lethal effect of NCS (Figure 1A). However, PARP-1^{+/+} were substantially more sensitive to NCS than PARP-1^{-/-} 3T3s in the high range of NCS concentration.

Pre-treatment with the PARP-1 inhibitor 4-amino-1,8-naphthalimide (ANI) did not alter the susceptibility of PARP-1^{+/+} 3T3s to NCS (Figure 2A). In contrast, PARP-1^{-/-} were considerably more sensitive than PARP-1^{+/+} 3T3s to the lethal effect of γ -rays (Figure 2B) and H₂O₂ (Figure 2C).

The PARP-1^{+/+} and PARP-1^{-/-} 3T3 clones used in these experiments originated each from independent isolates immortalized at random. There was therefore a possibility that minor differences might exist between both cell lines with regard to cell death mechanism or efficiency at a high level of DSB, and contribute to altered NCS susceptibility in addition to the PARP-1 defect. To settle this question, we used PARP-1^{-/-} 3T3s in which the full PARP-1 cDNA had been re-inserted by retroviral infection with a pBabe construct, allowing complete restoration of the PARP-1 activity [38]. The NCS response of these cells was compared to that of PARP-1^{-/-} 3T3s from the same clonal isolate transfected with the void vector. Both cell lines




Figure 1

Effect of PARP-1 knockout on the susceptibility of mouse 3T3 fibroblasts to the lethal effect of NCS. A (top). Cytotoxicity of NCS against PARP-1^{+/+} and PARP-1^{-/-} 3T3s. The growth assays were performed as described under Methods. For PARP-1^{-/-} 3T3s, the data were fitted to an exponential equation, $S = e^{-\alpha [NCS]}$ where S is the surviving fraction ($\alpha = 0.398 \pm 0.019 \text{ nM}^{-1}$). For PARP-1^{+/+} 3T3s, a linear-quadratic equation was used in order to smooth the curve and gave the same value of α . The insert shows the response of PARP-1^{+/+} and PARP-1^{-/-} 3T3s in the low range of NCS concentration. The amount of NCS (LC₃₇) required to reduce cell survival to 1/e of that in control was 1.79 and 2.48 nM for PARP-1^{+/+} and PARP-1^{-/-} 3T3s, respectively. B (bottom). Comparison of the response of PARP-1^{-/-} 3T3s from a unique isolate transduced with a pBabe-puro retroviral vector carrying the full-length PARP-1 cDNA (pBabe^{PARP-1}) or the void vector (pBabe^{void}). Both cell lines showed the same exponentially-dependent response to NCS ($\alpha = 0.566 \pm 0.044 \text{ nM}^{-1}$).




Figure 2

Differential susceptibility of PARP-1^{+/+} and PARP-1^{-/-} 3T3s to γ -rays and H₂O₂. A (top). Absence of a significant effect of ANI (30 μ M) on the cytotoxic response of PARP-1^{+/+} 3T3s to NCS. The results are expressed as the ratio of the mean lethal NCS concentrations (LC₃₇) determined from survival curves in the same way as in Figure 1. B (middle). γ -Ray survival of PARP-1^{+/+}

and PARP-1^{-/-} 3T3s. A modified form of the linear-quadratic equation, $S = a.e^{-\alpha D - \beta D^2} + b.e^{-\gamma D}$ where D is the radiation dose and (a + b) = 1, was set [6] to take into account the existence of a minor fraction (b) of the cell population experiencing cytolytic cell death at high radiation doses. The values calculated for best fit with the experimental data were: $\alpha = 0.134 \pm 0.040 \text{ Gy}^{-1}$, $\beta = 0.168 \pm 0.030 \text{ Gy}^{-2}$ for PARP-1^{-/-} 3T3; $\alpha = 0.0258 \pm 0.0323 \text{ Gy}^{-1}$, $\beta = 0.0422 \pm 0.0120 \text{ Gy}^{-2}$ for PARP-1^{+/+} 3T3s. α represents the contribution to radiation-induced cell death of lethal, non-repairable DNA damage. The quadratic parameter, β relates to unrepaired sublethal damage. Though this is still a matter of controversy [66], β is thought to represent the probability of interaction between separate breaks to exchange chromosomal aberrations [67]. The mean lethal radiation doses (D₃₇), i. e., the doses required to reduce cell survival to 1/e of that in control, are given in Table 1. C (bottom). Cytotoxicity of H₂O₂ against PARP-1^{+/+} and PARP-1^{-/-} 3T3s. The dose-response curves were fitted to a biexponential equation,

$S = a.e^{-\alpha[H_2O_2]} + b.e^{-\beta[H_2O_2]} + c$ with (a + b + c) = 1. c corresponds to the plateau of cell survival at infinite H₂O₂ concentration (0.48 and 0.046 for PARP-1^{+/+} and PARP-1^{-/-} 3T3s, respectively). The initial slope of the cells' response to H₂O₂ was calculated at 2.6 and 25.8 mM⁻¹ for PARP-1^{+/+} and PARP-1^{-/-} 3T3s, respectively.




Figure 3
Time-dependence of DSB rejoining in PARP-1^{+/+} and PARP-1^{-/-} 3T3s following exposure to 30 nM NCS. [2-¹⁴C]Thymidine-labeled cells were exposed to NCS for 10-min. After drug removal, cells were rapidly chilled in ice at the time indicated, harvested, and embedded into agarose plugs followed by lysis and PFGE (see Methods). The results are expressed as the percentage of remaining damage in Gy.eq. This percentage was determined from the fraction of activity released from the plugs; calibration was made with reference to samples irradiated in ice. The origin of the time scale starts from the introduction of NCS.

showed exactly the same susceptibility to NCS, with a purely exponential dose-dependence (Figure 1B).

Incidence of NCS-induced DSB

DSB in PARP-1^{+/+} and PARP-1^{-/-} 3T3s were measured by PFGE. Briefly, cells were exposed to increasing concentrations of NCS, up to 30 nM for 10-min at 37°C, then chilled in ice, harvested in PBS-EDTA buffer, and inserted into agarose plugs and lysed for PFGE analysis. The effect of NCS was compared to that of γ -rays (up to 60 Gy) on cells irradiated in ice.

The incidence of DSB was found to increase linearly with the γ -ray dose, in agreement with earlier reports [39]. It also grew linearly with the NCS concentration, and incubation with 1 nM NCS yielded the same amount of DSB as 1.21 Gy γ -rays (data not shown). PARP-1^{+/+} and PARP-1^{-/-} 3T3s did not show any difference in this assay. Taking into account the fact that 1 Gy γ -rays produces 41 DSB per diploid cell nucleus [40], the average incidence of DSB formed by 1 nM NCS in each 3T3 fibroblast was estimated at ca. 50 DSB. With consideration to the number of DSB induced by the treatment, the lethal efficiency of NCS was in the same range as that for γ -rays (Table 1).




Figure 4
Immunofluorescent visualization of PARP-1, pADPR synthesis and H2AX phosphorylation in PARP-1^{+/+} and PARP-1^{-/-} 3T3s exposed to various treatments including H₂O₂ (1 mM, 10-min) without or with 30 μ M ANI, γ -rays (20 Gy, 10-min) or NCS (2 or 30 nM, 10-min). PARP-1^{-/-} 3T3s complemented with the PARP-1 cDNA (pBabe^{PARP-1}) are shown for comparison. Cells were grown on coverslips, treated, fixed, incubated with antibodies and counterstained with DAPI. For each preparation the gain of the camera was adjusted relative to DAPI emission. The top part of the figure shows isolated views. The bottom panels show DAPI and immunofluorescence views in coincidence. The bar (top right view) represents 20 μ m.

Rejoining of NCS-induced DSB

To determine whether the repair of NCS-induced DSB was deficient in PARP-1^{-/-} relative to PARP-1^{+/+} 3T3s, DSB rejoining in cells treated with 30 nM NCS was measured through PFGE. The data (Figure 3) did not demonstrate any difference in the rejoining kinetics between both cell lines.

Immunofluorescence assays

PARP-1^{+/+} and PARP-1^{-/-} 3T3s were assayed through immunofluorescence for the determination of (i) PARP-1 expression, (ii) PARP-1 activity visualized by pADPr synthesis following γ -ray irradiation, NCS and H₂O₂, and (iii) ATM-dependent H2AX histone phosphorylation in response to DNA damage induced by γ -rays, H₂O₂ or NCS. The results are shown in Figure 4.

Table 1: Comparison of the mean lethal dose D_{37} (or LC_{37}), i. e., the radiation dose (or NCS concentration) leaving $1/e = 37\%$ survival, among the two cell lines investigated.

	PARP-1 ^{+/+} 3T3	PARP-1 ^{-/-} 3T3
D_{37} γ-Rays (Gy)	4.96	2.26
LC_{37} NCS (Gy.eq)	2.21	3.00

The D_{37} (or LC_{37}) values were calculated from the dose-dependent (γ-rays) or concentration-dependent survival curves (NCS), taking into account the mean incidence of NCS-induced DSB relative to γ-rays (1 nM NCS = 1.21 Gy.eq).

PARP-1 expression was detected in the nucleus of PARP-1^{+/+} 3T3s and was missing in PARP-1^{-/-} 3T3s, as expected. Also as expected, pADPr synthesis was observed selectively in the nucleus of PARP-1^{+/+} cells following γ-ray or H₂O₂ exposure, and was repressed by ANI. Stable transfection with a retroviral vector coding for the full PARP-1 gene, restored PARP-1 expression and function.

In contrast to γ-rays or H₂O₂, NCS did not induce any measurable pADPr synthesis in PARP-1^{+/+} 3T3s, even at a concentration of 30 nM representing ca. 20-fold the IC₅₀ value and yielding the same amount of DSB as 36 Gy radiation. However, consistent with induction of DSB in chromosomal DNA, NCS at relatively low concentration (2 nM) initiated rapid phosphorylation of H2AX in the same way as γ-rays or H₂O₂. In agreement with this observation, MRE11 phosphorylation and focus formation through an ATM- and NBS1-dependent mechanism, has recently been shown to occur after NCS treatment [37].

Discussion

The incidence of NCS-induced DSB was the same in PARP-1^{+/+} and PARP-1^{-/-} 3T3s, indicating that both cell lines incorporated equal amounts of NCS_{Chrom}. Moreover no difference, even minor was found between PARP-1^{+/+} and PARP-1^{-/-} in the kinetics of DSB rejoining (Figure 3).

Consistent with this, the susceptibility to NCS-induced lethality was the same in PARP-1^{+/+} and PARP-1^{-/-} 3T3s below 1.5 nM drug. PARP-1^{+/+} were comparatively more sensitive to NCS in the high range of NCS concentration. This is thought to result from minor differences in the susceptibility to DSB-induced cell death arising between different clonal isolates of 3T3 fibroblasts immortalized at random. As a matter of fact, retrovirus-mediated insertion of the PARP-1 cDNA in a single PARP-1^{-/-} 3T3 clone, did not bear any change in NCS susceptibility (Figure 1).

ANI totally repressed pADPr elongation in response to H₂O₂ (Figure 4) but did not alter the cytotoxicity of NCS in PARP-1^{+/+} 3T3s (Figure 2). These results corroborate those of other authors who showed that PARP-1 inhibitors did not affect cell kill by the topoisomerase IIα poison etoposide, another DSB inducer [41–43]. In con-

trast, and in agreement with earlier reports [6,44,45] PARP-1^{-/-} 3T3s were considerably more sensitive than PARP-1^{+/+} 3T3s to the lethal effect of radiation (Figure 2C). PARP-1^{-/-} 3T3s were also more sensitive than PARP-1^{+/+} to sub-millimolar concentrations of H₂O₂, a potent inducer of pADPr synthesis in response to SSB and oxidative base damage [46], highlighting the toxic effect of a deficiency in base excision repair in PARP-1 knockout cells [47]. However, for both cell lines, and contrary to CHO cells that exhibited an exponentially concentration-dependent response to H₂O₂ [46], the survival curves presented a plateau indicating resistance of a major (PARP-1^{+/+}) or minor fraction (PARP-1^{-/-}) of the cell population (Figure 2C). A similar effect was reported by other authors in Chinese hamster V79 cells and in normal and AT human fibroblasts exposed to H₂O₂ or organic peroxides [48,49]. Differential expression of catalase, superoxide dismutase or glutathione peroxidase was ruled out [48] and at the moment there is no explanation to this observation.

Other authors using single-cell gel electrophoresis (comet assay) have shown that the rejoining of radiation-, H₂O₂- or methyl methane sulfonate-induced SSB, is dramatically hampered by PARP-1 knockout, and significantly delayed by PARP-1 inhibitors in L1210 cells [50] and PARP-1^{+/+} 3T3s [51,52] yet these inhibitors have poor incidence on the level of H₂O₂-induced cell death [46,53]. Ectopic expression of the PARP-1 cDNA in PARP-1^{-/-} 3T3s restored a wild-type phenotype [51], demonstrating unequivocally that the repair capacity of PARP-1-deficient cells was drastically limited. We propose that this effect is relevant to the role of PARP-1 in base excision repair only, as there was no difference between PARP-1^{+/+} and PARP-1^{-/-} 3T3s with regard to the rejoining kinetics of NCS-induced DSB (Figure 3). Moreover, immunofluorescence assays showed that, even though NCS at supra-lethal concentration did not induce detectable pADPr synthesis, the ATM-dependent phosphorylation of H2AX was activated in response to relatively low levels of NCS, irrespective of whether PARP-1 was defective or not (Figure 4). Bowman *et al.* [43] also showed that the topoisomerase IIα poison etoposide did not bring about pADPr synthesis in L1210 cells, even at concentrations in excess of those causing sig-

nificant levels of apoptosis. Likewise, Dziegielewski and Beerman [54] recently showed that a defect in either ATM or DNA-PKcs may not confer cells enhanced susceptibility to the NCS homologue C-1027. On the other hand single-strand nicks in DNA are effective activators for PARP-1, not for DNA-PK [55,56]. Therefore, PARP-1-dependent pADPr elongation and ATM-dependent H2AX phosphorylation, two major post-translational modifications occurring in response to DNA single- and double-strand breaks, respectively, are likely not to be interrelated or redundant. This may elicit functional synergy between PARP-1 and ATM, consistent with the embryonic lethality observed in the double ATM/PARP-1 knockout [57].

Conclusions

Substantial differences in PARP-1 activation by ionizing radiation and alkylating agents have been reported [58] and our data suggest that PARP-1 is not a major determinant of cell survival to DNA double-strand breaks as long as the level of oxidative stress (as determined from pADPr synthesis) is limited. In consideration of studies with other DNA-damaging agents (reviewed in [59]) and of the comparatively high susceptibility of PARP-1 null 3T3s to H₂O₂ (Figure 2C), we propose that enhanced radiation sensitivity of PARP-1 null relatively to normal 3T3s, proceeds from defects in base excision repair in the former.

Methods

Antibodies and reagents

Mouse monoclonal antibodies directed against the PARP-1 C-terminal domain (clone 7-D3-6) and pADPr (clone 10 H) were from Becton-Dickinson Biosciences and Alexis Corporation, respectively. Rabbit polyclonal antibody raised against γ -H2AX, was from Trevigen. Alexa-488[®]-conjugated goat anti-rabbit and anti-mouse secondary antibodies, were purchased from Molecular Probes. HRP-conjugated goat anti-mouse and anti-rabbit IgG came from Jackson Immunoresearch.

[2-¹⁴C]Thymidine was purchased from PerkinElmer Life Sciences. 4-amino-1,8-naphthalimide (ANI) came from ACROS Organics. H₂O₂ was from Sigma-Aldrich. Other chemicals were of the highest purity available and came from VWR International. The products for cell culture were from Invitrogen.

The neocarzinostatin holoprotein, prepared and titrated as described [60], was stored as a sterile 1 mM stock solution in 2 mM sodium formate buffer, pH 4.0, at liquid nitrogen temperature.

Cells and cell cultures

Spontaneously immortalized 3T3 mouse embryo fibroblasts obtained from homozygous wild-type (PARP-1^{+/+}) and knockout (PARP-1^{-/-}) C57BL/6 mice [44], were pro-

vided by Dr. Gilbert de Murcia. It should be stressed that the knockout in PARP-1^{-/-} fibroblasts was complete, i. e., no fragment of the functional domain of PARP-1 protein was expressed in these cells. PARP-1^{-/-} 3T3s in which the full PARP-1 gene was reinserted by a retroviral pBabe-puro construct [38] were a generous gift of Dr. Moshe Oren.

3T3 fibroblasts were maintained as exponentially growing monolayers in Dulbecco modified Eagle's minimum essential medium (DMEM), with antibiotics and 10% foetal calf serum as described [6]. The mean doubling time of cells was 27-h and 52-h for PARP-1^{+/+} and PARP-1^{-/-} 3T3s, respectively.

Cytotoxicity determination

Due to a low cloning efficiency of 3T3 fibroblasts in a feeder layer technique, all cytotoxicity determinations were performed using growth assays. PARP-1^{+/+} or PARP-1^{-/-} 3T3 fibroblasts were plated in triplicate (10⁵ cells per 25 cm² vented flasks) and incubated overnight before treatment. After treatment, the flasks were rinsed twice with HBSS and returned to fresh medium for exactly 5 doubling times, with one medium change at day 7 for PARP-1^{-/-} 3T3s. Fibroblasts were harvested by 0.05% trypsin-0.02% EDTA, pelleted, resuspended in medium and counted under microscope in a Malassez cuvette.

Radiation and drug treatment

Irradiation of cells was performed at room temperature in medium equilibrated with 5% CO₂ in air, using an IBL-637 (¹³⁷Cs) γ -ray irradiator (Cis-Biointernational). The dose rate was 0.86 Gy/min or 8.0 Gy/min for survival or immunofluorescence studies, respectively.

Aliquots of NCS were thawed just before use, adjusted to the suitable dilution in ice-cold PBS buffer, pH 6.0, and immediately introduced into culture flasks (25 cm², 5 ml medium) with gentle agitation. The whole treatment was performed in dim light to avoid photo-induced degradation of the drug. The cytotoxic effect of NCS at fixed concentration was investigated as a function of the length of drug exposure. Cytotoxicity increased steeply with the length of contact and reached completion after 6-min only. More prolonged incubation did not result in increased cell death. Therefore, the length of contact with NCS was 10-min throughout.

For studies of H₂O₂ response, serial dilutions of concentrated (9.8 M) H₂O₂ were made in pure water, then in culture medium immediately prior to use. The length of contact with H₂O₂ was 10-min.

In some experiments ANI was used as a PARP-1 inhibitor [61–63]. When present, ANI (30 μ M) from a 3 mM stock solution in pure DMSO was introduced 1-h prior to irra-

diation or NCS, and removed 1-h later with two HBSS washes. The DMSO concentration in medium was 1% and was kept constant throughout experiments with ANI. Controls were made with DMSO alone at the same concentration.

DNA double-strand break determination

The determination of DSB formation and repair was carried out by pulsed field agarose gel electrophoresis (PFGE) through the clamped homogeneous electric field (CHEF) technique [64,65]. [^{14}C]Thymidine-labeled cells (2–5 10^6 cells per 25 cm^2 flask) were put in ice following treatment, harvested in ice-cold PBS supplemented with 2 mM EDTA by gentle scraping, and collected by sedimentation at 4°C. Pellets from 5 10^5 cells were resuspended in 150 μl , final volume, of PBS buffer at 37°C, and mixed under mild vortexing with an equal volume of 1.6% w/v low-melting-point agarose (BDH Laboratory) in PBS buffer at 37°C. The suspension (300 μl) was immediately pipetted into pre-chilled 4 × 10 mm moulds and allowed to form plugs on ice for 30-min. Embedded cells were subsequently lysed by immersion of the plugs in 2 ml of a solution containing 2% N-lauroyl sarkosine, 40 mM EDTA, 1 mg/ml proteinase K in PBS, pH 7.8 and incubated, firstly for 2 h in ice, secondly for 24 h at 50°C. The plugs were subsequently washed twice with PBS, and treated for 1-h with 200 $\mu\text{g}/\text{ml}$ RNase A. The plugs were finally washed twice with PBS and stored at 4°C overnight prior electrophoresis.

The plugs were inserted into the wells of an 0.8% w/v agarose gel (BioRad, chromosomal grade) made in 0.75 × TAE buffer (40 mM tris-acetate, 2 mM EDTA, pH 8), and the gels submitted to PFGE at 14°C in a CHEF-DR III apparatus (Bio-Rad) with buffer recirculation at 14°C. Migration was for 72 h at 2 V/cm with three switch times, namely, 1200-s, 1500-s and 1800-s with angles of 96°, 100° and 106°, respectively. The molecular weight markers were *S. pombe* and *S. cerevisiae* chromosomes (Bio-Rad).

After electrophoresis, the gels were stained with 0.5 $\mu\text{g}/\text{ml}$ ethidium bromide under mild agitation (30-min), followed by destaining for 1-h in fresh buffer. DNA fluorescence was visualized over an UV transilluminator with camera recording. The gels were subsequently dried *in vacuo* over a piece of Whatman paper and analyzed using a Phosphorimager[®] apparatus (Amersham Pharmacia Biotech), allowing a precise determination of the migration profile of DNA fragments and of the fractional radioactivity released from the plugs (FAR).

Immunofluorescence of PARP-1, pADPr and γ -H2AX

3T3 fibroblasts were grown for 24-h on coverslips and exposed to either γ -rays (5 Gy for γ H2AX, 50 Gy for

pADPr), H_2O_2 (1 mM, 10-min contact, 37°C), or graded concentrations of NCS (2 nM, 5 nM or 30 nM, 10-min contact, 37°C), and fixed 10-min after the beginning of treatment. For PARP-1 and pADPr immunofluorescence, fixation was by 4% formaldehyde (10-min, room temperature) in PBS followed by two PBS washes and neutralization of residual formaldehyde by 50 mM NH_4Cl (10-min, room temperature). After a further PBS wash (5-min, 4°C), cells were permeabilized by 0.5% Triton X-100 in PBS (5-min, 4°C), and finally rinsed twice with PBS. For γ H2AX determination, cells were fixed in a 1:1 v:v acetone:methanol mixe (-20°C, 10-min), dried in air and rehydrated in PBS (15-min, room temperature).

Cell preparations on coverslips were subsequently incubated with 1% bovine serum albumin in PBS (10-min, room temperature) and exposed to anti-pADPr, anti-PARP-1 or anti- γ H2AX primary antibody (1/200 dilution, 1-h, 37°C). The coverslips were rinsed thrice with PBS, and incubated with Alexa-488[®]-conjugated secondary antibody (1/500 dilution, 30-min, 37°C), rinsed thrice with PBS, counterstained with 4,6-diamidino-2-phenylindole (DAPI) and mounted. Immunofluorescence was visualized using a Zeiss Axiophot microscope equipped with a Micromax chilled camera (Princeton Applied Research).

Abbreviations

ANI, 4-amino-1,8-naphthalimide; ATM, ataxia telangiectasia mutated protein kinase; CHEF, clamped homogeneous electric field; DAPI, 4,6-diamidino-2-phenylindole; DNA-PK, DNA-dependent protein kinase; DSB, double-strand break; EDTA, ethylene dinitrilo tetraacetate; HBSS, Hank's balanced salt solution; NCS, holo-neocarzinostatin; $\text{NCS}_{\text{Chrom}}$, NCS chromophore; pADPr, poly(ADP-ribose); PARP-1, poly(ADP-ribose) polymerase; PBS, phosphate buffered saline; PFGE, pulsed-field agarose gel electrophoresis; SSB, single-strand break; TAE, Tris-acetate-EDTA buffer.

Authors' Contributions

GN and NG carried out the survival and DNA repair determinations. MF and FMC performed immunofluorescence analyses. VF conceived the study, and participated in its design and coordination. All authors have read and approved the final manuscript.

Acknowledgements

Grateful thanks are due to Drs. G. de Murcia and J. Ménessier-de Murcia (UPR 9003 CNRS, ESBS, Illkirch-Graffenstaden, France) for the generous gift of PARP-1^{-/-} 3T3 fibroblasts, and to Dr. Moshe Oren (Dept of Molecular Cell Biology, Weizmann Institute of Science, Rehovot, Israel) who provided PARP-1^{-/-} 3T3s complemented with the PARP-1 cDNA. This work was supported by grants from *Electricité de France* (RB 2001–02) and the *Institut Curie* (Protein Remodelling Program), and by financial aid from the *Institut National de la Santé et de la Recherche Médicale*. M.F. is recipient of a post-doctoral fellowship from the *Ligue Nationale Contre le Cancer*.

References

- Burma S, Chen BP, Murphy M, Kurimasa A and Chen DJ: **ATM phosphorylates histone H2AX in response to DNA double-strand breaks** *J Biol Chem* 2001, **276**:42462-42467.
- Andegeko Y, Moyal L, Mittelman L, Tsarfaty I, Shiloh Y and Rotman G: **Nuclear retention of ATM at sites of DNA double strand breaks** *J Biol Chem* 2001, **276**:38224-38230.
- Taccioli GE, Gottlieb TM, Blunt T, Priestley A, Demengeot J, Mizuta R, Lehman AR, Alt FW, Jackson SP and Jeggo PA: **Ku80: product of the XRCC5 gene and its role in DNA repair and V(D)J recombination** *Science* 1994, **265**:1442-1445.
- Wu X and Lieber MR: **Protein-protein and protein-DNA interaction regions within the DNA end-binding protein Ku70-Ku86** *Mol Cell Biol* 1996, **16**:5186-5193.
- Jeggo PA, Taccioli GE and Jackson SP: **Ménage à trois: double-strand break repair, V(D)J recombination and DNA-PK** *Bioessays* 1995, **17**:949-957.
- Fernet M, Ponette V, Dienaud-Alexandre E, Ménéssier-de Murcia J, de Murcia G, Giocanti N, Méglin-Chanet F and Favaudon V: **Poly(ADP-ribose) polymerase, a major determinant of early cell response to ionising radiation** *Int J Radiat Biol* 2000, **76**:1621-1629.
- Dantzer F, Nasheuer HP, Vonesch JL, de Murcia G and Ménéssier-de Murcia J: **Functional association of poly(ADP-ribose) polymerase with DNA polymerase α -primase complex: a link between DNA strand break detection and DNA replication** *Nucleic Acids Res* 1998, **26**:1891-1898.
- Masson M, Niedergang C, Schreiber V, Müller S, Ménéssier-de Murcia J and de Murcia G: **XRCC1 is specifically associated with poly(ADP-ribose) polymerase and negatively regulates its activity following DNA damage** *Mol Cell Biol* 1998, **18**:3563-3571.
- de Murcia G and Shall S: *From DNA damage and stress signalling to cell death. Poly ADP-ribosylation reactions* Oxford University Press, New York; 2000.
- de Murcia G and Ménéssier-de Murcia J: **Poly(ADP-ribose) polymerase: a molecular nick-sensor** *Trends Biochem Sci* 1994, **19**:172-176.
- Wesierska-Gadek J, Bugajska-Schretter A and Cerni C: **ADP-ribosylation of p53 tumor suppressor protein: mutant but not wild-type p53 is modified** *J Cell Biochem* 1996, **62**:90-101.
- Pleschke JM, Kleczkowska HE, Strohm M and Althaus FR: **Poly(ADP-ribose) binds to specific domains in DNA damage checkpoint proteins** *J Biol Chem* 2000, **275**:40974-40980.
- Morrison C, Smith GCM, Stingl L, Jackson SP, Wagner EF and Wang ZQ: **Genetic interaction between PARP and DNA-PK in V(D)J recombination and tumorigenesis** *Nat Genet* 1997, **17**:479-482.
- Ruscetti T, Lehnert BE, Halbrook J, Le Trong H, Hoekstra MF, Chen DJ and Peterson SR: **Stimulation of the DNA-dependent protein kinase by poly(ADP-ribose) polymerase** *J Biol Chem* 1998, **273**:14461-14467.
- Galante S and Kohwi-Shigematsu T: **Poly(ADP-ribose) polymerase and Ku autoantigen form a complex and synergistically bind to matrix attachment sequences** *J Biol Chem* 1999, **274**:20521-20528.
- Ariumi Y, Masutani M, Copeland TD, Mimori T, Sugimura T, Shimotohno K, Ueda K, Hatanaka M and Noda M: **Suppression of the poly(ADP-ribose) polymerase activity by DNA-dependent protein kinase in vitro** *Oncogene* 1999, **18**:4616-4625.
- Ariumi Y, Ueda K, Masutani M, Copeland TD, Noda M, Hatanaka M and Shimotohno K: **In vivo phosphorylation of poly(ADP-ribose) polymerase is independent of its activation** *FEBS Lett* 1998, **436**:288-292.
- Brown ML, Franco D, Burkle A and Chang Y: **Role of poly(ADP-ribosylation) in DNA-PKcs-independent V(D)J recombination** *Proc Natl Acad Sci USA* 2002, **99**:4532-4537.
- Favaudon V: **On the mechanism of reductive activation in the mode of action of some anticancer drugs** *Biochimie (Paris)* 1982, **64**:457-475.
- Dedon PC and Goldberg IH: **Free-radical mechanisms involved in the formation of sequence-dependent bistranded DNA lesions by the antitumor antibiotics bleomycin, neocarzinostatin, and calicheamicin** *Chem Res Toxicol* 1992, **5**:311-332.
- Kappen LS and Goldberg IH: **Deoxyribonucleic acid damage by neocarzinostatin chromophore: strand breaks generated by selective oxidation of C-5' of deoxyribose** *Biochemistry* 1983, **22**:4872-4878.
- Favaudon V, Charnas RL and Goldberg IH: **Poly(deoxyadenylic-deoxythymidylic acid) damage by radiolytically activated neocarzinostatin** *Biochemistry* 1985, **24**:250-259.
- Povirk LF, Houlgrave CW and Han YH: **Neocarzinostatin-induced DNA base release accompanied by staggered oxidative cleavage of the complementary strand** *J Biol Chem* 1988, **263**:19263-19266.
- Hashimoto M, Greenberg MM, Kow YW, Hwang JT and Cunningham RP: **The 2-deoxyribonolactone lesion produced in DNA by neocarzinostatin and other damaging agents forms cross-links with the base-excision repair enzyme endonuclease III** *J Am Chem Soc* 2001, **123**:3161-3162.
- Ohtsuki K and Ishida N: **Neocarzinostatin-induced breakdown of deoxyribonucleic acid in HeLa-S3 cells** *J Antibiot (Tokyo)* 1975, **28**:143-148.
- Beerman TA and Goldberg IH: **The relationship between DNA strand-scission and DNA synthesis inhibition in HeLa cells treated with neocarzinostatin** *Biochim Biophys Acta* 1977, **475**:281-293.
- Tatsumi K and Nishioka H: **Effect of DNA repair systems on antibacterial and mutagenic activity of an antitumor protein, neocarzinostatin** *Mutat Res* 1977, **48**:195-203.
- Denklauf D, Stahl R and Kohnlein W: **Effect of neocarzinostatin on E. coli mutants deficient in DNA repair** *Z Naturforsch [C]* 1989, **44**:791-796.
- Moustacchi E and Favaudon V: **Cytotoxic and mutagenic effects of neocarzinostatin in wild-type and repair-deficient yeasts** *Mutat Res* 1982, **104**:87-94.
- Shiloh Y, Tabor E and Becker Y: **Cellular hypersensitivity to neocarzinostatin in ataxia-telangiectasia skin fibroblasts** *Cancer Res* 1982, **42**:2247-2249.
- Babilon RW, Soprano KJ and Henderson EE: **Hypersensitivity and reduced inhibition of DNA synthesis in ataxia telangiectasia lymphoblasts treated with low levels of neocarzinostatin** *Mutat Res* 1985, **146**:79-87.
- Helbig R, Zdzienicka MZ and Speit G: **The effect of defective DNA double-strand break repair on mutations and chromosome aberrations in the Chinese hamster cell mutant XR-V15B** *Radiat Res* 1995, **143**:151-157.
- Povirk LF: **DNA damage and mutagenesis by radiomimetic DNA-cleaving agents: bleomycin, neocarzinostatin and other enediynes** *Mutat Res* 1996, **355**:71-89.
- Muller C and Salles B: **Regulation of DNA-dependent protein kinase activity in leukemic cells** *Oncogene* 1997, **15**:2343-2348.
- Muller C, Calsou P and Salles B: **The activity of the DNA-dependent protein kinase (DNA-PK) complex is determinant in the cellular response to nitrogen mustards** *Biochimie (Paris)* 2000, **82**:25-28.
- van Duijn-Goedhart A, Zdzienicka MZ, Sankaranarayanan K and van Buul PP: **Differential responses of Chinese hamster mutagen sensitive cell lines to low and high concentrations of calicheamicin and neocarzinostatin** *Mutat Res* 2000, **471**:95-105.
- Yuan SSF, L. CH, Hou MF, Chan TF, Kao YH, Wu YC and Su JH: **Neocarzinostatin induces Mre11 phosphorylation and focus formation through an ATM- and NBS1-dependent mechanism** *Toxicol* 2002, **177**:123-130.
- Wang X, Michael D, de Murcia G and Oren M: **p53 activation by nitric oxide involves down-regulation of Mdm2** *J Biol Chem* 2002, **277**:15697-15702.
- Frankenberg-Schwager M: **Review of repair kinetics for DNA damage induced in eukaryotic cells in vitro by ionizing radiation** *Radiother Oncol* 1989, **14**:307-320.
- Dahm-Daphi J and Dikomey E: **Separation of DNA fragments induced by ionizing irradiation using a graded-field gel electrophoresis** *Int J Radiat Biol* 1995, **67**:161-168.
- Chatterjee S, Cheng MF, Berger RB, Berger SJ and Berger NA: **Effect of inhibitors of poly(ADP-ribose) polymerase on the induction of GRP78 and subsequent development of resistance to etoposide** *Cancer Res* 1995, **55**:868-873.
- Sebestyen A, Mihalik R, Petak I and Kopper L: **Modulation of apoptosis signaling in etoposide-treated lymphoma cells** *Anticancer Res* 1997, **17**:2609-2614.
- Bowman KJ, Newell DR, Calvert AH and Curtin NJ: **Differential effects of the poly(ADP-ribose) polymerase (PARP) inhibitor**

- NU1025 on topoisomerase I and II inhibitor cytotoxicity in L1210 cells in vitro** *Br J Cancer* 2001, **84**:106-112.
44. Ménissier-de Murcia J, Niedergang C, Trucco C, Ricoul M, Dutrillaux B, Mark M, Oliver FJ, Masson M, Dierich A, LeMeur M, Walztinger C, Chambon P and de Murcia G: **Requirement of poly(ADP-ribose) polymerase in recovery from DNA damage in mice and in cells** *Proc Natl Acad Sci USA* 1997, **94**:7303-7307.
 45. Trucco C, Rolli V, Oliver FJ, Flatter E, Masson M, Dantzer F, Niedergang C, Dutrillaux B, Ménissier-de Murcia J and de Murcia G: **A dual approach in the study of poly(ADP-ribose) polymerase: in vitro random mutagenesis and generation of deficient mice** *Mol Cell Biochem* 1999, **193**:53-60.
 46. Cantoni O, Murray D and Meyn RE: **Effect of 3-aminobenzamide on DNA strand-break rejoining and cytotoxicity in CHO cells treated with hydrogen peroxide** *Biochim Biophys Acta* 1986, **867**:135-143.
 47. Dantzer F, de La Rubia G, Ménissier-de Murcia J, Hostomsky Z, de Murcia G and Schreiber V: **Base excision repair is impaired in mammalian cells lacking poly(ADP-ribose) polymerase-I** *Biochemistry* 2000, **39**:7559-7569.
 48. Kaneko M, Kodama M and Inoue F: **Bimodal pattern of killing of Chinese hamster V79 variant cells by hydrogen peroxide** *Free Radic Res* 1994, **20**:229-239.
 49. Shackelford RE, Innes CL, Sieber SO, Heinloth AN, Leadon SA and Paules RS: **The Ataxia telangiectasia gene product is required for oxidative stress-induced G(1) and G(2) checkpoint function in human fibroblasts** *J Biol Chem* 2001, **276**:21951-21959.
 50. Bowman KJ, White A, Golding BT, Griffin RJ and Curtin NJ: **Potentiation of anti-cancer agent cytotoxicity by the potent poly(ADP-ribose) polymerase inhibitors NU1025 and NU1064** *Br J Cancer* 1998, **78**:1269-1277.
 51. Trucco C, Oliver FJ, de Murcia G and Ménissier-de Murcia J: **DNA repair defect in poly(ADP-ribose) polymerase-deficient cell lines** *Nucleic Acids Res* 1998, **26**:2644-2649.
 52. Atorino L, Di Meglio S, Farina B, Jones R and Quesada P: **Rat germinal cells require PARP for repair of DNA damage induced by gamma-irradiation and H₂O₂ treatment** *Eur J Cell Biol* 2001, **80**:222-229.
 53. Cantoni O, Cattabeni F, Stocchi V, Meyn RE, Cerutti P and Murray D: **Hydrogen peroxide insult in cultured mammalian cells: relationships between DNA single-strand breakage, poly(ADP-ribose) metabolism and cell killing** *Biochim Biophys Acta* 1989, **1014**:1-7.
 54. Dziegielewska J and Beerman TA: **Cellular responses to the DNA strand-scission enediyne C-1027 can be independent of ATM, ATR, and DNA-PK kinases** *J Biol Chem* 2002, **277**:20549-20554.
 55. Weinfeld M, Chaudhry MA, D'Amours D, Pelletier JD, Poirier GG, Povirk LF and Lees-Miller SP: **Interaction of DNA-dependent protein kinase and poly(ADP-ribose) polymerase with radiation-induced DNA strand breaks** *Radiat Res* 1997, **148**:22-28.
 56. D'Silva I, Pelletier JD, Lagueux J, D'Amours D, Chaudhry MA, Weinfeld M, Lees-Miller SP and Poirier GG: **Relative affinities of poly(ADP-ribose) polymerase and DNA-dependent protein kinase for DNA strand interruptions** *Biochim Biophys Acta* 1999, **1430**:119-126.
 57. Ménissier-de Murcia J, Mark M, Wendling O, Wynshaw-Boris A and de Murcia G: **Early embryonic lethality in PARP-1 Atm double-mutant mice suggests a functional synergy in cell proliferation during development** *Mol Cell Biol* 2001, **21**:1828-1832.
 58. Le Rhun Y, Kirkland JB and Shah GM: **Cellular responses to DNA damage in the absence of poly(ADP-ribose) polymerase** *Biochem Biophys Res Commun* 1998, **245**:1-10.
 59. Griffin RJ, Curtin NJ, Newell DR, Golding BT, Durkacz BW and Calvert AH: **The role of inhibitors of poly(ADP-ribose) polymerase as resistance-modifying agents in cancer therapy** *Biochimie* 1995, **77**:408-422.
 60. Favaudon V: **Gamma-radiolysis study of the reductive activation of neocarzinostatin by the carboxyl radical** *Biochimie (Paris)* 1983, **65**:593-560.
 61. Banasik M, Komura H, Shimoyama M and Ueda K: **Specific inhibitors of poly(ADP-ribose) synthetase and mono(ADP-ribosyl) transferase** *J Biol Chem* 1992, **267**:1569-1575.
 62. Berges F and Zeller WJ: **Combination effects of poly(ADP-ribose) polymerase inhibitors and DNA-damaging agents in ovarian tumor cell lines, with special reference to cisplatin** *J Cancer Res Clin Oncol* 1996, **122**:665-670.
 63. Schlicker A, Peschke P, Bürkle A, Hahn EW and Kim JH: **4-amino-1,8-naphthalimide: a novel inhibitor of poly(ADP-ribose) polymerase and radiation sensitizer** *Int J Radiat Biol* 1999, **75**:91-100.
 64. Blöcher D, Einspenner M and Zajackowski J: **CHEF electrophoresis, a sensitive technique for the determination of DNA double-strand breaks** *Int J Radiat Biol* 1989, **56**:437-448.
 65. Woudstra EC, Roesink JM, Rosemann M, Brunsting JF, Driessen C, Orta T, Konings AW, Peacock JH and Kampinga HH: **Chromatin structure and cellular radiosensitivity: a comparison of two human tumour cell lines** *Int J Radiat Biol* 1996, **70**:693-703.
 66. Cornforth MN: **Analyzing radiation-induced complex chromosome rearrangements by combinatorial painting** *Radiat Res* 2001, **155**:643-659.
 67. Cornforth MN and Bedford JS: **A quantitative comparison of potentially lethal damage repair and the rejoining of interphase chromosome breaks in low passage normal human fibroblasts** *Radiat Res* 1987, **111**:385-405.

Publish with **BioMed Central** and every scientist can read your work free of charge

"BioMed Central will be the most significant development for disseminating the results of biomedical research in our lifetime."

Sir Paul Nurse, Cancer Research UK

Your research papers will be:

- available free of charge to the entire biomedical community
- peer reviewed and published immediately upon acceptance
- cited in PubMed and archived on PubMed Central
- yours — you keep the copyright

Submit your manuscript here:
http://www.biomedcentral.com/info/publishing_adv.asp

