

**Reciprocal regulation of brain and muscle Arnt-like protein 1 and peroxisome proliferator-activated receptor alpha defines a novel positive feedback loop in the rodent liver circadian clock.**

Laurence Canaple, Juliette Rambaud, Ouria Dkhissi-Benyahya, Béatrice Rayet, Nguan Soon Tan, Liliane Michalik, Franck Delaunay, Walter Wahli, Vincent Laudet

► **To cite this version:**

Laurence Canaple, Juliette Rambaud, Ouria Dkhissi-Benyahya, Béatrice Rayet, Nguan Soon Tan, et al.. Reciprocal regulation of brain and muscle Arnt-like protein 1 and peroxisome proliferator-activated receptor alpha defines a novel positive feedback loop in the rodent liver circadian clock.. Mol Endocrinol, 2006, 20 (8), pp.1715-27. 10.1210/me.2006-0052 . inserm-00135943

**HAL Id: inserm-00135943**

**<https://www.hal.inserm.fr/inserm-00135943>**

Submitted on 19 Mar 2007

**HAL** is a multi-disciplinary open access archive for the deposit and dissemination of scientific research documents, whether they are published or not. The documents may come from teaching and research institutions in France or abroad, or from public or private research centers.

L'archive ouverte pluridisciplinaire **HAL**, est destinée au dépôt et à la diffusion de documents scientifiques de niveau recherche, publiés ou non, émanant des établissements d'enseignement et de recherche français ou étrangers, des laboratoires publics ou privés.

**Reciprocal regulation of BMAL1 and PPAR $\alpha$  defines a novel positive feedback loop in the rodent liver circadian clock.**

Laurence Canaple<sup>\*¶</sup>, Juliette Rambaud\*, Ouria Dkhissi-Benyahya<sup>°</sup>, Béatrice Rayet<sup>§</sup>, Nguan Soon Tan<sup>#</sup>, Liliane Michalik<sup>#</sup>, Franck Delaunay<sup>§</sup>, Walter Wahli<sup>#</sup>, Vincent Laudet<sup>\*1</sup>

\* Structure and Evolution of Nuclear Receptors, CNRS UMR 5161, IFR 128 BioSciences Lyon-Gerland, Laboratoire de Biologie Moléculaire de la Cellule, Ecole Normale Supérieure, 46 allée d'Italie, 69364 Lyon cedex 07, France.

° INSERM U-371, Laboratoire Cerveau et Vision, IFR 19-UCBL, 18 avenue du Doyen Lépine, 69675 Bron, France.

§ Université de Nice Sophia Antipolis CNRS FRE 2721 Bâtiment de Sciences Naturelles 28, Avenue Valrose 06108 Nice cedex 2 France.

# Center for Integrative Genomics, University of Lausanne, Genopode, CH-1015 Lausanne, Switzerland.

¶ Present address: ANIMAGE, CNRS UMR 5515, 59 Boulevard Pinel, 69677 Bron cedex, France.

<sup>1</sup> Corresponding author: CNRS UMR 5161, IFR 128 BioSciences Lyon-Gerland, Laboratoire de Biologie Moléculaire de la Cellule, Ecole Normale Supérieure, 46 allée d'Italie, 69364 Lyon cedex 07, France. Tel : 33 4 72 72 81 90 ; Fax : 33 4 72 72 80 80 ; e-mail : vincent.laudet@ens-lyon.fr

The authors have nothing to declare.

Running Title: PPAR $\alpha$  and liver circadian clock

Key words: PPAR $\alpha$ , BMAL1, Peripheral clock, Fibrates

**Abstract**

Recent evidence has emerged that PPAR $\alpha$ , which is largely involved in lipid metabolism, can play an important role in connecting circadian biology and metabolism. In the present study, we investigated the mechanisms by which PPAR $\alpha$  influences the pacemakers acting in the central clock located in the suprachiasmatic nucleus and in the peripheral oscillator of the liver. We demonstrate that PPAR $\alpha$  plays a specific role in the peripheral circadian control as it is required to maintain the circadian rhythm of the master clock gene *bmal1* *in vivo*. This regulation occurs *via* a direct binding of PPAR $\alpha$  on a PPRE located in the *bmal1* promoter. Reversely, BMAL1 is an upstream regulator of *PPAR\alpha* gene expression. We further demonstrate that fenofibrate induces circadian rhythm of clock gene expression in cell culture and up-regulates hepatic *bmal1* *in vivo*. Altogether, these results provide evidence for an additional regulatory feedback loop involving BMAL1 and PPAR $\alpha$  in peripheral clocks.

## Introduction

Circadian rhythms enable numerous organisms to adapt to daily environmental changes such as light, temperature and social communication and serve to synchronize multiple molecular, biochemical, physiological and behavioral processes. Circadian rhythms persist with an approximate 24 h periodicity even in temporally isolated subjects, indicating the presence of an autonomous time keeping system called circadian clock. In mammals, circadian rhythms are generated by the main pacemaker located in the suprachiasmatic nucleus (SCN) of the hypothalamus (1). To ensure that internal time coincides with environmental time, the clock must be adjusted, a process known as entrainment. In mammals, light received by the eyes synchronizes the oscillator through the retino-hypothalamic tract and hence synchronizes the behavior of the organism with the daily 24 h light/dark cycle (*For review : (2-6)*).

In addition to the SCN, other peripheral tissues such as liver, heart, kidney (7, 8) as well as isolated cells (9) express clock genes giving rise to circadian rhythms with a different phase from that observed in the SCN. Interestingly, these peripheral clocks can be reset by alternative routes independently of the SCN, for example by forced change of feeding time (10, 11). Several lines of evidence suggest that the peripheral circadian clocks are not SCN-independent but require inputs from the SCN to drive the rhythmicity and ensure an ordered response of the organism to environmental changes (12, 13). Thus, the SCN is believed to coordinate rhythms in the brain and body *via* a combination of neural and humoral diffusible and synaptic signals (7, 14, 15)

Genetic analyses have identified master clock genes such as *clock*, *bmal1*, *period* (*per1*, 2) and *cryptochrome* genes (*cry1*, 2), as well as the orphan nuclear receptor genes, *rev-erba* and *rora* (16, 17). Other transcription factors functioning in the circadian regulation of gene expression, including DBP-related factors (18), Rev-erb $\beta$ , ROR $\beta$  and  $\gamma$  (19) have also

been identified. The clock mechanism mainly involves an integrated network of interacting self-sustained transcriptional-translational feedback loops, composed of positive and negative regulators, which drive their own rhythmic expression and the one of clock-controlled genes to perform a fine tuning of circadian gene expression (20).

Recent reports have highlighted the interplay between circadian oscillators, metabolism and physiology. Whereas genes involved in the glucose and lipid metabolism are known to exhibit circadian variations (21-23), molecular studies have revealed a critical role for *bmal1* and *clock* genes in regulating glucose homeostasis (24) and lipid metabolism (25-28). Moreover, the cross-talk between ROR $\alpha$  and Rev-erb $\alpha$  was shown to be physiologically important for the control of cholesterol and triglyceride metabolism (29-31). In turn, the peripheral clocks can be coordinately regulated by multiple circulating factors, which are affected by the metabolic status of the organism. Indeed, glucose, one of the major food metabolites that exhibits a plasma diurnal rhythm, is a direct resetting signal in cultured cells by down-regulating *per1* and *per2* RNA levels (32). The levels of glucose-regulated hormones such as insulin or glucagon immediately up-regulate *per1* and *per2* expression (13). In addition, other studies have revealed an important role of glucocorticoids and retinoids in the resetting of peripheral clocks (33, 34). Though the elucidation of the mechanisms that govern the connection between metabolism and circadian clock has just begun, it appears that several members of the nuclear receptor family are involved in this pathway.

Evidence has emerged that peroxisome proliferator-activated receptor alpha (PPAR $\alpha$ ), a member of the nuclear receptor superfamily that regulates the expression of numerous genes involved in lipid metabolism and energy homeostasis, can play a role in the normal circadian regulation. First, *PPAR\alpha* has been identified as a circadian clock-controlled gene with a diurnal rhythm at the mRNA and protein levels in rats and mice in many peripheral organs such as liver, heart, kidney and to a lesser extent in the SCN, where the central pacemaker is

located (7, 35, 36). This circadian expression of *PPAR $\alpha$*  may be in part controlled by hormonal factors, since insulin and glucocorticoids regulate its mRNA expression (35, 37-39). A recent study has also shown that the circadian expression of *PPAR $\alpha$*  mRNA is regulated by the peripheral oscillators in a CLOCK-dependent manner (27). Second, since daily variations in lipogenic and cholesterogenic gene expression are attenuated or abolished in mice in which the *PPAR $\alpha$*  gene has been disrupted, PPAR $\alpha$  may be an important mediator for the circadian regulation of lipid metabolism (40, 41). It is now believed that PPAR $\alpha$  has a wider general role in transducing hormone messages involved in dietary status (42). These observations thus suggest that PPAR $\alpha$  may be required in the control of circadian food-dependent fluctuations in gene expression. Third, PPAR $\alpha$  is connected to the regulation of other nuclear hormone receptors such as Rev-erb $\alpha$ , as fenofibrate, a PPAR $\alpha$  agonist, induces human and rat *rev-erb $\alpha$*  expression in liver through the direct binding of PPAR $\alpha$  on an atypical DR2 element located in the *rev-erb $\alpha$*  promoter (30, 43). It was recently shown that CLOCK plays an important role in lipid homeostasis by regulating the circadian transactivation of PPARE-controlled target genes (26) and of *PPAR $\alpha$*  gene itself *via* an E-box rich region *in vivo* and *in vitro* (27). Fourth, the partner of PPAR $\alpha$ , RXR $\alpha$ , interacts with CLOCK protein in a ligand-dependent manner and inhibits CLOCK/BMAL1-dependent activation *via* an E-box element (34). Other results have also suggested that *PPAR $\alpha$* -deficiency disturbs the normal circadian regulation of certain SREBP sensitive genes in the liver (40, 44).

Despite much evidence supporting a role of PPAR $\alpha$  in metabolic control and energy homeostasis (45) and the accumulation of data connecting metabolism and circadian biology (3, 46, 47), little is known concerning the influence of PPAR $\alpha$  on the circadian clock. In the present study, we investigated the mechanisms by which PPAR $\alpha$  can influence the pacemakers acting in the SCN and in the liver. We report that *PPAR $\alpha$*  deficient (*PPAR $\alpha$ <sup>-/-</sup>*)

mice present similar locomotor activity with wild type (WT) mice without any molecular alteration of clock gene expression in the SCN. Interestingly, we show for the first time that PPAR $\alpha$  is a direct regulator of *bmal1* expression in liver *via* its direct binding on a PPRE located on the *bmal1* promoter. This regulation is required to maintain the normal circadian oscillation of *bmal1* *in vivo*. Stressing the importance of the regulatory pathway that exists between PPAR $\alpha$  and the peripheral clock genes, we show that fenofibrate up-regulates *bmal1* gene expression in murine liver and induces circadian rhythm of clock gene expression in cell culture in a PPAR $\alpha$  dependent manner. Reversely, we also observe that BMAL1 is an upstream regulator of the *PPAR\alpha* gene expression. Taken together our data implicate PPAR $\alpha$  in a new regulatory loop that controls peripheral circadian clocks.

## Results

### *PPAR $\alpha$ <sup>-/-</sup> mice display normal circadian locomotor activity and clock gene expression in the SCN.*

In order to evaluate the influence of PPAR $\alpha$  on the function of the central circadian oscillator, behavioral analysis of the circadian rhythm was carried out using *PPAR $\alpha$ <sup>-/-</sup>* and WT littermate mice. Animals were first synchronized for 2 weeks to a 12L:12D cycle. Under these light conditions, WT and *PPAR $\alpha$ <sup>-/-</sup>* mice entrained normally and consolidated their locomotor activity to the dark period of the LD cycle (figure 1A). No difference in the total amount of daily activity was observed between WT and *PPAR $\alpha$ <sup>-/-</sup>* mice (figure 1B). When placed in constant darkness, the *PPAR $\alpha$* -deficient mice do not display an arrhythmic behaviour with endogenous period similar between both genotypes (24,11 $\pm$ 0,11 hrs and 24,12  $\pm$ 0,08 hrs respectively in *PPAR $\alpha$* -deficient and WT mice).

To determine if the inactivation of *PPAR $\alpha$*  can alter the master oscillator in the SCN at the molecular level, we further compared the circadian expression profiles of clock genes in the SCN of WT and *PPAR $\alpha$*  mutant mice (figure 1C). There is no significant difference in the amplitude and in the phase of the circadian expression of the tested clock genes (*bmal1*, *per2*, *per3*, *cry2* and *rev-erb $\alpha$* ). This clearly suggests that there is no major molecular effect of the *PPAR $\alpha$*  deletion at the central clock level and that the expression of *PPAR $\alpha$*  in the SCN is not essential for the basal maintenance of the central circadian timing system.



***PPAR $\alpha$  is required to maintain the amplitude of the circadian expression of *bmal1* in the murine liver.***

To evaluate the role of PPAR $\alpha$  on the circadian system of the liver (a peripheral clock), where it is mainly expressed, we analyzed the circadian expression of several clock genes in liver isolated from WT and PPAR $\alpha$  mutant mice. As already reported, PPAR $\alpha$  expression effectively follows a circadian rhythm in peripheral tissues such as liver, kidney and muscle (data not shown). Figure 2 shows that all the clock genes tested (*bmal1*, *per1*, *per2*, *per3*, *cry2* and *rev-erba*) are expressed in a circadian manner with no modification in the phase of their rhythm between both genotypes. By contrast, the amplitudes of *bmal1* and *per3* expression are drastically affected in PPAR $\alpha$ -deficient mice by comparison to the WT, with a significant decrease at circadian times CT1 and CT21 for *bmal1* (where CT0 is subjective day beginning at 7 am and CT12 is subjective night beginning at 7 pm) and an increase at CT8 for *per3*. These data suggest that PPAR $\alpha$  does not influence the phase synchronisation properties of the liver clock but affect the amplitude of two major clock genes *bmal1* and *per3*.

***Food-induced phase resetting entrains circadian PPAR $\alpha$  expression and is globally maintained in liver of PPAR $\alpha$ <sup>-/-</sup> mice.***

Although the day-night cycle is the most obvious time cue, animals can also respond to other synchronizing signals and feeding time appears to be a potent temporal cue, or *Zeitgeber*, for the liver clock (10, 11). According to PPAR $\alpha$  role in the hepatic lipid metabolism during starvation, we hypothesized its potential role in food phase resetting of the liver clock. To examine whether the daily feeding time can affect the phase of the PPAR $\alpha$  circadian expression in the liver, WT mice were fed for two weeks exclusively during the day or during the night. As expected, mice fed exclusively during the day displayed an inversed phase in

circadian hepatic expression of *bmall*, *per1*, *per3* and *rev-erb $\alpha$*  gene by comparison to mice fed only during the night. Similarly, feeding during the day entirely inversed the phase of liver *PPAR $\alpha$*  expression (figure 3A), result that is in agreement with the demonstration that *PPAR $\alpha$*  is a clock-controlled gene in the liver (7, 36). Control mice fed only during the night displayed a similar phase of hepatic clock gene expression and *PPAR $\alpha$*  than mice fed *ad libitum*.

Next, we studied whether feeding time can also reset the phase of *bmall*, *per1*, *per3* and *rev-erb $\alpha$*  expression in the absence of *PPAR $\alpha$*  (figure 3B). Except the *bmall* expression which was refractory to resetting, *per1*, *per3* and *rev-erb $\alpha$*  genes showed an inversed rhythm of their expression in the liver after daytime-restricted feeding compared to nighttime feeding in the *PPAR $\alpha$*  knock-out context. The present results show that feeding time can reset the expression of *per1*, *per3* and *rev-erb $\alpha$*  in the liver of *PPAR $\alpha$* <sup>-/-</sup> mice and suggest that *bmall* expression might be controlled by *PPAR $\alpha$* .

***Fenofibrate induces expression of clock genes in Rat-1 fibroblasts and up-regulates *bmall* gene expression in liver.***

Since *PPAR $\alpha$*  presents a circadian expression in liver, we first studied whether a shock with a serum-rich medium is able to induce an oscillation of *PPAR $\alpha$*  in the well established *in vitro* model Rat-1 fibroblasts (9, 13, 48-50). As shown in figure 4A, *PPAR $\alpha$*  gene expression is induced by a serum shock in Rat-1 fibroblasts with a maximal level reached 12 hours after the beginning of the treatment by comparison with the control. The induced-oscillating expressions of *rev-erb $\alpha$*  and *cry1* after serum shock were in accordance with previously reported data (9, 51). In addition, *rev-erb $\alpha$*  expression is delayed (peak at 16 hours after the serum shock treatment) compared to *PPAR $\alpha$*  expression which suggests that in Rat-1

fibroblasts *PPAR $\alpha$*  induces *rev-erba* expression. Similarly to these observations, it seems likely that serum largely participates in the synchronization of the circadian oscillation of *PPAR $\alpha$*  mRNA in fibroblast cultures.

Fibrates are well-known activators of *PPAR $\alpha$*  expression and are classical drugs used in the treatment of dyslipidemias. Using the same *in vitro* model, we then evaluated if fenofibrate can induce expression of *PPAR $\alpha$*  gene. The addition of fenofibrate into the serum-free medium triggered a rhythmic expression of *PPAR $\alpha$*  peaking 12-14 hours as observed after a serum treatment (figure 4B). Fenofibrate induced a *PPAR $\alpha$*  expression 6-fold higher than serum (figures 4A and 4B).

Fenofibrate can also induce a rhythmic expression of clock genes. The temporal induction by fenofibrate of *cry1*, *rev-erba* and *bmal1* expression into the culture medium was almost similar to that observed after a serum-shock: *cry1* mRNA level peaked at Zeitgeber times ZT8-10 (where ZT0 is time when the light switched on at 7 am and ZT12 is time when the light switched off at 7 pm), *rev-erba* at ZT16-20 and *bmal1* at ZT4 (figure 4C). Thus fenofibrate can act as a *Zeitgeber* in cell culture and trigger a rhythm of clock gene expression.

To determine if fenofibrate can also reset the liver clock *in vivo*, the response of *bmal1* and *rev-erba* genes to fenofibrate treatment was compared in liver of WT and *PPAR $\alpha$ <sup>-/-</sup>* mice (figure 4D). In accordance with our previous results, fenofibrate was also able to markedly induce *bmal1* mRNA levels in control mice but not in *PPAR $\alpha$*  deficient mice, suggesting that *bmal1* induction by fenofibrate is effectively mediated by *PPAR $\alpha$* . Similarly as it was previously reported in rat liver and human hepatocytes (30, 43), administration of fenofibrate in WT mice significantly increased the *rev-erba* mRNA levels. As expected, *rev-erba* expression in *PPAR $\alpha$*  knock-out mice was not induced by fenofibrate, confirming that murine

*rev-erb $\alpha$*  induction by fenofibrate is mediated by PPAR $\alpha$ . Taken together these data strongly suggest that fenofibrate can alter the endogenous rhythm of the liver peripheral clock and *in vivo*

***PPAR $\alpha$  is a direct regulator of *bmall* expression.***

To specify the regulation of *bmall* gene expression by PPAR $\alpha$  suggested by our analysis of *bmall* expression in the PPAR $\alpha$ <sup>-/-</sup> mice, we analysed the interaction of PPAR $\alpha$  with the *bmall* promoter. First we performed a bioinformatic research of potential PPAR $\alpha$  binding sites (PPREs) on the *bmall* promoter region using the Nubiscan software. Two major PPREs were predicted at the positions -1519 (+, TGGACATGGGTCA) and -4943 (-, AGGGCTGAGGACA), the start site corresponding to the one identified in mouse testis (52). In order to evaluate whether PPAR $\alpha$  binds to the *bmall* gene promoter *in vivo*, the occupancy of the potential PPRE binding sites by PPAR $\alpha$  was analysed using ChIP assays performed on hepatocyte DNA using an anti-PPAR $\alpha$  antibody (figure 5A). As previously shown *in vitro*, the DNA encompassing the *rev-erb $\alpha$*  Rev-DR2 site (position -45) was precipitated *in vivo* by the anti-PPAR $\alpha$  antibody after fibrate treatment (figure 5A, lower panel, lanes 7 and 9), in accordance with a PPAR $\alpha$ -Rev-erb $\alpha$  crosstalk through competition for binding to the same Rev-DR2 site (43). Moreover an amplification product was observed when the same DNA samples were PCR-amplified using primers covering the PPRE located at the position -1519 in the *bmall* promoter (figure 5A, upper panel, lanes 7 and 9). No amplification product was obtained using primers flanking the site at the position -4943 (middle panel). These data further demonstrate that in mice *in vivo*, PPAR $\alpha$  directly binds to the PPRE site located at the position -1519 of the *bmall* promoter in peripheral oscillators and therefore that *bmall* is a direct PPAR $\alpha$  target gene.

***PPAR $\alpha$  mRNA expression is severely down-regulated in the liver of *bmal1*<sup>-/-</sup> mice.***

To test the hypothesis that the regulation of circadian *PPAR $\alpha$*  expression involves *bmal1* in the liver, we analyzed the daily accumulation of *PPAR $\alpha$*  mRNA in the liver of *bmal1*<sup>-/-</sup> mice by quantitative PCR. *PPAR $\alpha$*  mRNA expression is rhythmic in *bmal1*<sup>-/-</sup> mice with a maximum around ZT12 as observed in WT mice whereas the amplitude of the peak of expression is significantly dampened (figure 5B). This result suggests that BMAL1 is an upstream regulator of *PPAR $\alpha$*  gene and is consistent with the recent observation that *PPAR $\alpha$*  expression is also CLOCK-dependent (27).

## Discussion

### *A new regulatory feedback loop involved BMAL1 and PPAR $\alpha$ in peripheral clocks.*

Molecular dissection of the mechanisms by which the clock oscillating system is controlled remains one of the most important challenges to assess the importance of the circadian regulation in diverse physiological and metabolic processes in mammals. In the present study we have integrated the *PPAR $\alpha$*  gene and its protein into a new positive regulatory feedback loop in the liver. A model summarizing our main results is shown in figure 6. We show that *PPAR $\alpha$*  play an important role in the endogenous rhythmic property of peripheral clocks *in vivo*, whereas in the central clock *PPAR $\alpha$*  deficiency does not alter the circadian expression of clock genes. These clock gene expressions are only affected in the liver of *PPAR $\alpha$* <sup>-/-</sup> deficient mice. First *PPAR $\alpha$*  is not essential to drive the central circadian system since *PPAR $\alpha$* <sup>-/-</sup> mice entrained normally without arrhythmic behaviour in constant darkness and displayed no alteration in the amplitude and the phase of circadian expression of the clock genes (*bmal1*, *per2*, *per3*, *cry2* and *rev-erba*) compared to WT mice. The lack of *PPAR $\alpha$* -dependent clock regulation *in vivo* in the SCN can be due either to the absence of a circadian *PPAR $\alpha$*  function in the central clock or to the presence of another isotype of *PPAR* ( $\beta$ ,  $\gamma$ ) or other nuclear receptors that are able to bind to the *PPRE* site and to exert a compensatory effect. Second, the expression of *bmal1*, an essential gene of the molecular oscillator, is drastically reduced in the liver of *PPAR $\alpha$* <sup>-/-</sup> deficient mice. This regulation of *bmal1* transcription by *PPAR $\alpha$*  is likely through a direct binding of *PPAR $\alpha$*  to the *PPRE* element located at the position -1519 in the *bmal1* promoter. In addition to the established transactivation of the circadian promoter *rev-erba* by *PPAR $\alpha$*  (43), our data largely support the view that *PPAR $\alpha$*  also associates with the circadian *bmal1* promoter *in vivo*. Interestingly, *PPAR $\alpha$*  expression is strongly decreased

in the liver of *bmal1*<sup>-/-</sup> deficient mice compared to WT mice. As no significant change in hepatic CLOCK level is found in *bmal1*<sup>-/-</sup> deficient mice compared with WT mice (53), this suggests that BMAL1 itself is in return involved in the circadian transactivation of *PPAR $\alpha$*  gene at the level of peripheral oscillators in mice. Other observations have also revealed that CLOCK is involved in the circadian transactivation of *PPAR $\alpha$*  (27) and interacts with its partner RXR (34). We propose that *bmal1* is directly and positively regulated by PPAR $\alpha$  and that BMAL1 imposes in return a circadian regulation on *PPAR $\alpha$*  transcription.

***PPAR $\alpha$  expression is resetted by feeding in peripheral clocks.***

Liver is known to be the organ reacting most rapidly to the temporal feeding regimen as it plays a dominant role in the metabolism and processing of food components, such as proteins, lipids and carbohydrates. It was clearly established that the circadian gene expression in peripheral hepatic cells is intimately connected to feeding (10, 11) and that PPAR $\alpha$  is involved in food processing and energy homeostasis (45). A connection between circadian gene regulation, metabolism and energy homeostasis was also established. For example, glucose metabolism (e.g., 6-phosphofructokinase-2, aldolase and glucose phosphate isomerase) is under the control of the circadian time keeping system (23, 24). Rhythmic expression of numerous enzymes and transcription factors involved in protein and amino acid metabolism (e.g., serine dehydratase, DBP and 3-hydroxy-3-methylglutaryl coenzyme A reductase, (7, 54), in fat metabolism (e.g., cholesterol 7  $\alpha$ -hydroxylase, PPAR $\alpha$ , HMGCoA lyase and reductase, (7, 35, 55) or in detoxification process (e.g., steroid 15  $\alpha$ -hydroxylase (Cyp2a4) and coumarin 7-hydroxylase (Cyp2a5), Cyp2e1, Cyp17 and glutathione-S-transferase theta 2, (23, 56) was also observed in liver. Herein we show that the circadian expression of *PPAR $\alpha$*  is resetted by feeding time and that in the absence of PPAR $\alpha$ ,

clock gene expression in liver is entrained by reversed feeding time, suggesting that PPAR $\alpha$  does not play a major role in food resetting. We underline that after an extended duration of daytime feeding, the food imposed reversed phase of circadian gene expression in peripheral liver is similar between WT and PPAR $\alpha$ <sup>-/-</sup> deficient mice. This is probably due to PPAR $\alpha$ -independent signalling pathway that plays a major role in the phase resetting of circadian gene expression by feeding time.

***Fibrates as a resetting signal in cell cultures and peripheral clocks.***

Previous studies have shown that PPAR $\alpha$  expression is positively controlled by glucocorticoids and fibrates and negatively by insulin (37-39). Fibrates are also known to up-regulate the expression of *rev-erba* in the rat liver and in both rat and human primary hepatocyte cultures (30, 43). *In vitro*, PPAR $\alpha$  mRNA was induced in rat fibroblasts culture after both a serum shock and more interestingly, a fenofibrate treatment. Moreover, fenofibrate efficiently stimulates the rhythmic expression of several clock genes such as *cry1*, *bmal1* and *rev-erba*. This suggests that fenofibrate is able to entrain rhythmic PPAR $\alpha$  and clock gene expressions in Rat-1 fibroblasts. *In vivo*, *rev-erba* mRNA level is up-regulated by fenofibrate in the mouse liver of WT animals. As this up-regulation of *rev-erba* gene by fibrates is not observed in the PPAR $\alpha$  deficient mice, we confirm *in vivo* that the induction of *rev-erba* gene expression by fibrates is mediated by PPAR $\alpha$  at the transcriptional level. Interestingly, the increased accumulation of PPAR $\alpha$  after fibrate treatment leads in turn to a higher level of expression of *bmal1* in WT mice. Other convincing evidence that the fibrate effect on the *bmal1* expression is exerted at the transcriptional level *via* the PPAR $\alpha$  protein is that fibrates have no effect on the *bmal1* expression in the PPAR $\alpha$ <sup>-/-</sup> mice. Taken together,



these results suggest that PPAR $\alpha$  could play a role in integrating chemical signals inside the liver.

***How circadian rhythm might influence a fibrate therapy?***

To date, fibrates are clinically used as hypolipidemic drugs that lower plasma cholesterol and triglycerides. They exert their effect by regulating the expression of several key genes implicated in lipid metabolism *via* PPAR $\alpha$  activation. Interestingly, fibrate therapy represents a cost-effective approach in the clinical management and the prevention of cardiovascular diseases in a growing population suffering from lifestyle-induced metabolic dysfunctions such as obesity, insulin-resistance and diabetes (57). Although important differences in lipid metabolism exist between mice and human, including the function of PPAR $\alpha$  (58, 59) our *in vitro* and *in vivo* studies suggest that the regulation by fibrates of the circadian expression of clock genes may influence the success of a treatment as it would suggest a potential induction and subsequently a dysfunction of their expression after fibrate supply in patients. The validity of this hypothesis remains to be demonstrated in human. Therefore, one additional question to address is how fibrate administration at a selected time of the day can impact the efficacy and the success of the treatment. It will be of interest to decipher the molecular mechanisms involved in the circadian expression of clock genes and PPAR $\alpha$ -regulated genes in presence or absence of fibrates to provide new insight in the downstream circadian physiological and cellular processes governed by PPAR $\alpha$  itself. Similarly, such understanding should lead to new strategies for pharmacological manipulation of the human clock to improve the treatment of dyslipidaemias.

In summary, our data indicate that PPAR $\alpha$  is a specific element of the liver oscillatory clock in mammals and plays an important role in integrating signals into the clock machinery. We clearly demonstrate that *in vivo*, PPAR $\alpha$  is required to maintain normal circadian oscillation of the master clock gene *bmal1* in liver. This regulation occurs *via* a direct binding of PPAR $\alpha$  on a PPRE located in the *bmal1* promoter. In addition, BMAL1 is an upstream regulator of

the *PPAR $\alpha$*  expression. This finding provides a new regulatory pathway for the circadian system and suggests that some transcription factors may have a specific role in the peripheral clocks. Further studies are now required to determine the exact impact of circadian rhythms on the metabolic processes governed by PPAR $\alpha$  in peripheral organs. This can be addressed by investigating the circadian regulation of PPAR $\alpha$  target genes and some of the downstream targets in the peripheral clock in WT, *PPAR $\alpha$* <sup>-/-</sup> and other clock mutant mice. The complete elucidation of the signalling elements involved in the interactions between central and peripheral clocks and the mechanisms that govern the interplay between metabolism and circadian oscillators will also have a major impact on the circadian field in the next future.

## Material and methods

### *Cell culture, serum and fibrate shock*

Rat-1 fibroblasts were grown in DMEM supplemented with 5% fetal calf serum and a mixture of penicillin/streptomycin/glutamine (PSG). The serum shock was done as described elsewhere using 50% horse serum (9). For the fibrate shock, the medium was exchanged with DMEM-PSG supplemented with 50 $\mu$ M fenofibrate (Sigma) after the cells reach confluence (time 0). This medium was replaced with fenofibrate-free DMEM-PSG medium after 2 hours. At 0, 4, 8, 12, 16, 20, 24 hours after shock, cells were lysed and kept at  $-70^{\circ}\text{C}$  until RNA extraction. Whole-cell RNAs were extracted using GenElute Mammalian Total RNA extraction Kit from Sigma.

### *Animal experiments*

Purebred WT and homozygous  $PPAR\alpha^{-/-}$  mice on an SV129 background were used. All experiments were done with male mice between 6 to 8 weeks of age. Animals were kept under a 12 hours light:dark cycle (12L:12D) and food and drinking water were available *ad libitum*, except when indicated otherwise. The experimental protocols of the current research were approved by the rules and regulations of french veterinary services.

- ***Locomotor activity recording:*** Adult male mice (n=8 for both WT and  $PPAR\alpha^{-/-}$ ) were exposed to 12L:12D cycle for at least two weeks. For monitoring locomotor activity, mice were housed individually in cages equipped with infrared motion captors placed over the cages and a computerized data acquisition system (Circadian Activity Monitoring System, INSERM, France). Activity records were analyzed with the Clocklab software package (Actimetrics, Evanston, IL). For each animal, the total duration of activity was determined every 2 or 12 hours during the LD cycle and then averaged for WT and knockout mice.

Animals were then allowed to free run in constant darkness for at least 15 days. The endogenous period in DD was subsequently determined using the Clocklab software.

- **Circadian expression of clock genes in wild type versus PPAR $\alpha$ <sup>-/-</sup> mice:** 12 WT and 12 PPAR $\alpha$ <sup>-/-</sup> mice were maintained for a period of two weeks in a 12L:12D cycle and transferred in constant darkness (DD) the day before the sacrifice. Livers and SCN were removed at circadian times CT 1, 8, 14, 21, stored at -70°C until RNA extraction and analysed by quantitative RT-PCR. The experiment was done twice.

- **Restricted feeding:** Mice (n=24 both for WT and PPAR $\alpha$ <sup>-/-</sup>) fed during the day received food when light was on (7 am to 7 pm) whereas mice (n=24 both for WT and PPAR $\alpha$ <sup>-/-</sup>) fed during the night received food from 7 pm to 7 am for two weeks. Water was freely available over the experimental period. As controls, WT and PPAR $\alpha$ <sup>-/-</sup> mice (n=24 for both) were fed *ad libitum*. Mice were transferred in DD the day before sacrifice. Livers were dissected at indicated circadian times (CT 1, 8, 14, 21), stored at -70°C until RNA extraction and analysed by quantitative-RT-PCR.

- **Fenofibrate response:** In this experiment, mice (n=32 both for WT and PPAR $\alpha$ <sup>-/-</sup>) were treated for two weeks with fenofibric acid (vehicle DMSO) mixed in the drinking water at the final concentration of 7 mM. Control animals (n=32 both for WT and PPAR $\alpha$ <sup>-/-</sup>) were treated with vehicle in the drinking water. Livers were removed at circadian times CT 1, 8, 14, 21, stored at -70°C until RNA extraction and analysed by quantitative RT-PCR.

- **Analysis of Bmal 1<sup>-/-</sup> mice:** WT and homozygous *bmal1*<sup>-/-</sup> mice on B6 background (both females and males aged from 8 to 14 weeks provided by C. Bradfield, Wisconsin, USA) were kept under 12L:12D cycles and fed *ad libitum*. Livers were removed at indicated *Zeitgeber* time. Reverse-transcribed total RNAs from 3 to 5 animals per time point were analysed by quantitative -RT-PCR as described below.

***RNA extraction and Quantitative RT-PCR***

Total RNAs were prepared from cells or from organs according to the manufacturer's instructions (Sigma) and reverse transcribed using random primers and MMLV Reverse Transcriptase (Invitrogen). cDNA were then used as template for a quantitative real-time PCR assay using the QuantiTect SYBR Green PCR reagents (Qiagen) and the DNA Engine Opticon system (MJ Research). Each couple of oligonucleotides used were designed to hybridize on different exons. The sequences of forward and reverse primers were as follows:

bmal1 forward 5'-CCAAGAAAGTATGGACACAGACAAA-3'  
 bmal1 reverse 5'-GCATTCTTGATCCTTCCTTGGT-3'  
 cry1 forward 5'-CTGGCGTGGAAGTCATCGT-3'  
 cry1 reverse 5'-CTGTCCGCCATTGAGTTCTATG-3'  
 cry2 forward 5'-TGTCCCTTCCTGTGTGGAAGA-3'  
 cry2 reverse 5'-GCTCCCAGCTTGGCTTGAA-3'  
 per1 forward 5'-GGAGACCACTGAGAGCAGCAAG-3'  
 per1 reverse 5'-CGCACTCAGGAGGCTGTAGGC-3'  
 per2 forward 5'-ATGCTCGCCATCCACAAGA-3'  
 per2 reverse 5'-GCGGAATCGAATGGGAGAAT-3'  
 per3 forward 5'-GGCGTTCTACGCGCACACTGC-3'  
 per3 reverse 5'-CGCTGGTGCACATTCATACTGCG-3'  
 ppar $\alpha$  forward 5'-CGCTATGAAGTTCAATGCCTT-3'  
 ppar $\alpha$  reverse 5'-TGCAACTTCTCAATGTAGCC-3'  
 rev-erb $\alpha$  forward 5'-CATGGTGCTACTGTGTAAGGTGTGT-3'  
 rev-erb $\alpha$  reverse 5'-CACAGGCGTGCACTCCATAG-3'  
 36B4 forward 5'-ACCTCCTTCTTCCAGGCTTT-3'  
 36B4 reverse 5'-CCCACCTTGTCTCCAGTCTTT-3'

The efficiency (>95%) and the specificity of the amplification were controlled by generating standard curves and carrying out melting curves and agarose gels of the amplicons respectively. The relative levels of each RNA were calculated by  $2^{-CT}$  (CT standing as the cycle number in which SYBR Green fluorescence exceeds a constant threshold value) and normalized to the corresponding non cyclic 36B4 RNA levels. The presented values are means  $\pm$  SEM of duplicates of the same reaction for at least 3 different mice or 3 experimental points.

The significance of differences was assessed by distribution-free two-way ANOVA. Paired student's tests were used to compare WT and *PPAR $\alpha$* <sup>-/-</sup> or *bmal1*<sup>-/-</sup> mice. Results are presented as mean  $\pm$  SEM. Differences were considered significant when  $p < 0.05$ .

### ***Chromatin Immunoprecipitation (ChIP) assays***

ChIP experiments were performed as already described in IJpenberg *et al.* (60). Briefly, WT and *PPAR $\alpha$* <sup>-/-</sup> mice (n=3) were fed for 5 days with either Wy14,643 (50mg/kg/day) or vehicle. Immunoprecipitation of liver extracts was done using a PPAR $\alpha$  antibody and the immunoprecipitated DNA was PCR amplified using primers flanking either the *rev-erba* Rev-DR2 (GTGTCACTGGGGC) or potential PPAR $\alpha$  response element (PPRE, usually AGGTCANAGGTCA) on the *bmal1* promoter predicted using the computer program NUBISCAN available at the following website [www.nubiscan.unibas.ch](http://www.nubiscan.unibas.ch) (61). An equal volume of non-precipitated genomic DNA (Input) was amplified as positive control. One fifth of PCR products were separated on an ethidium bromide-stained 2% agarose gel.

### **Acknowledgment**

This work was supported by CNRS, ENS de Lyon, Région Rhône-Alpes, FRAF, MENRT and FP5-OldClock QLK6-CT-2002-02258. We thank J.Samarut and M. Janier for their financial support, B. Desvergne and G. Triqueneaux for their helpful discussions. We are also grateful to F.J. Gonzalez for the *PPAR $\alpha$* <sup>-/-</sup> mice.

## Figure legends

### Figure 1: Entrainment and free running locomotor activity of WT and *PPAR* $\alpha$ <sup>-/-</sup> mice

**A.** Representative actograms of locomotor activity for WT and *PPAR* $\alpha$ <sup>-/-</sup> mice under 12L:12D cycle and DD conditions. After 2 weeks of entrainment under a 12L:12D cycle, mice were placed in DD. Horizontal bar at the top of each actogram depicts the lighting conditions of LD cycles. Time spans in darkness are marked by grey shadowing.

**B.** 24-h profiles of spontaneous locomotor activity of WT (open boxes) and *PPAR* $\alpha$ -deficient mice (solid boxes). The distribution of activity was determined every 12 hours during LD cycle. Shaded areas indicate the dark period. Results are expressed as the means  $\pm$  SEM of values from 8 animals per group.

**C.** Circadian expression of *bmal1*, *per2*, *per3*, *cry2* and *rev-erba* mRNAs in SCN of WT (—◆—) and *PPAR* $\alpha$ <sup>-/-</sup> mice (--■--). Real-time PCR was used to determine transcript levels at 4 circadian times (CT1, 8, 13, 21). Transcript levels are displayed as relative quantity (RQ) after normalization to the non cyclic 36B4 expression levels in the same sample. Results are expressed as the means  $\pm$  SEM of values of two independent experiments, each realized with 3 animals for both genotypes at each time point. Statistically significant differences between WT and *PPAR* $\alpha$ -deficient mice are indicated by \* ( $p < 0,05$ ).

**Figure 2:** Circadian expression of *bmal1*, *per1*, *per2*, *per3*, *cry2* and *rev-erba* mRNAs in liver of WT (—◆—) and *PPAR* $\alpha$ <sup>-/-</sup> mice (--■--). Transcript level values are expressed as relative quantity (RQ) after normalization to the corresponding non cyclic 36B4 expression levels. Results are shown as the mean  $\pm$  SEM of values of two independent experiments, each with 3 animals for both genotypes at each time point. There were significant variations between the two genotypes at CT1 and CT21 for *bmal1* and CT8 for *per3* as indicated by \*  $p < 0,01$ .

**Figure 3:** Circadian gene expression in liver after restricted feeding in WT and *PPAR* $\alpha$ <sup>-/-</sup> mice.

**A.** *PPAR* $\alpha$  expression in the liver of food-entrained WT mice (daytime (—◆—) or nighttime feeding (--■--)).

**B.** Circadian accumulation of *bmal1*, *per1*, *per3* and *rev-erba* mRNAs in liver of WT and *PPAR* $\alpha$ <sup>-/-</sup> mice as a function of daytime (—◆—) or nighttime feeding (--■--). Transcript levels are displayed as relative quantity (RQ) after normalization to the corresponding non cyclic 36B4 expression levels. The presented values are expressed as means  $\pm$  SEM of duplicates of the same reaction for 6 different mice per genotypes.

**Figure 4:** Effect of serum shock or fenofibrate on clock gene expression.

**A.** Accumulation of *PPAR* $\alpha$  (—◆—), *cry1* (—▲—) and *rev-erba* (—■—) mRNAs in Rat-1 fibroblasts shocked with 50% of horse serum. *PPAR* $\alpha$  accumulation in absence of horse serum (--■-- is reported as a control of non-induction. The presented values are means  $\pm$  SEM of duplicates of the same reaction for 3 different experimental points.

**B.** Accumulation of *PPAR* $\alpha$  mRNAs in Rat-1 fibroblasts shocked with 50  $\mu$ M fenofibrate (—■—) or with vehicle (--◆-- as controls. The presented values are means  $\pm$  SEM of duplicates of the same reaction for 3 independent experimental points.

**C.** Accumulation of *bmal1*, *cry1* and *rev-erba* mRNAs in Rat-1 fibroblasts shocked with fenofibrate (—■—) or with vehicle (--◆-- as controls. The presented values are means  $\pm$  SEM of duplicates of 3 independent experimental points.



**D.** Circadian accumulation of *bmal1* and *rev-erb $\alpha$*  mRNAs in liver of WT and *PPAR  $\alpha$* <sup>-/-</sup> mice treated (black boxes) or not (white boxes) with fenofibrate. In this experiment, mice were treated for two weeks with fenofibric acid (vehicle DMSO) mixed in the drinking water at the final concentration of 7 mM. Control animals were treated with the vehicle in the drinking water. Real-time PCR was used to determine transcript levels. Transcript level values are displayed as relative quantity (RQ) after normalization to the non cyclic 36B4 expression levels in the same sample. Results are expressed as the means  $\pm$  SEM of values from 8 animals for both genotypes at each time point.

**Figure 5:** Reciprocal regulation of BMAL1 and PPAR $\alpha$

**A.** ChIP of the potential *bmal1* and *rev-erb $\alpha$*  PPRE elements with PPAR $\alpha$  antibodies. WT and *PPAR $\alpha$* <sup>-/-</sup> mice (n=3 for both genotypes) were fed for 5 days with either Wy14,643 or vehicle (V). ChIP of liver extracts was performed with a PPAR $\alpha$  antibody (PPAR-Ab) and analysed by PCR for enrichment of the PPRE element of the *bmal1* promoter (top panels) and of the *rev-erb $\alpha$*  promoter Rev -DR2 (bottom panel). p.i.: preimmune serum; Input: non-precipitated genomic DNA.

**B.** Daily expression of *PPAR $\alpha$*  mRNAs in WT (—●—) and *bmal1* mutant mice (··■··) using real-time PCR. Transcript levels were normalized against the non cyclic 36B4 transcript level in the same sample. Results are expressed as the means  $\pm$  SEM of values from 3-5 animals at each time point. Statistically significant differences between WT and deficient mice are indicated by \* p<0,0001.

**Figure 6:** Model of cross-talk between PPAR $\alpha$  and circadian pathways depicting the control of circadian regulation by PPAR $\alpha$  in peripheral clocks.

In mammals, circadian rhythms are generated by the main pacemaker located in the suprachiasmatic nucleus (SCN) of the hypothalamus, which synchronizes the peripheral oscillators and ensure an ordered response of the organism in terms of physiology, metabolism and behavior to environmental changes. These peripheral clocks can be resetted by alternative routes such as feeding time. Herein, we show that PPAR $\alpha$  is entirely resetted by feeding time. We propose that *bmal1* transcription is directly positively regulated by PPAR $\alpha$  and that BMAL1 imposes circadian regulation on *PPAR $\alpha$*  transcription (bold arrows). The amplified accumulation of PPAR $\alpha$  under fibrate treatment leads in turn to a higher level of expression of *bmal1* gene (dashed arrows).

## References

1. Reppert SM, Weaver DR 2002 Coordination of circadian timing in mammals. *Nature* 418:935-41.
2. Challet E, Caldelas I, Graff C, Pevet P 2003 Synchronization of the molecular clockwork by light- and food-related cues in mammals. *Biol Chem* 384:711-9.
3. Albrecht U, Eichele G 2003 The mammalian circadian clock. *Curr Opin Genet Dev* 13:271-7.
4. Morse D, Sassone-Corsi P 2002 Time after time: inputs to and outputs from the mammalian circadian oscillators. *Trends Neurosci* 25:632-7.
5. Dunlap JC 1999 Molecular bases for circadian clocks. *Cell* 96:271-90.
6. Sassone-Corsi P 1998 Molecular clocks: mastering time by gene regulation. *Nature* 392:871-4.
7. Panda S, Antoch MP, Miller BH, Su AI, Schook AB, Straume M, Schultz PG, Kay SA, Takahashi JS, Hogenesch JB 2002 Coordinated transcription of key pathways in the mouse by the circadian clock. *Cell* 109:307-20.
8. Kita Y, Shiozawa M, Jin W, Majewski RR, Besharse JC, Greene AS, Jacob HJ 2002 Implications of circadian gene expression in kidney, liver and the effects of fasting on pharmacogenomic studies. *Pharmacogenetics* 12:55-65.
9. Balsalobre A, Damiola F, Schibler U 1998 A serum shock induces circadian gene expression in mammalian tissue culture cells. *Cell* 93:929-37.
10. Stokkan KA, Yamazaki S, Tei H, Sakaki Y, Menaker M 2001 Entrainment of the circadian clock in the liver by feeding. *Science* 291:490-3.
11. Damiola F, Le Minh N, Preitner N, Kornmann B, Fleury-Olela F, Schibler U 2000 Restricted feeding uncouples circadian oscillators in peripheral tissues from the central pacemaker in the suprachiasmatic nucleus. *Genes Dev* 14:2950-61.
12. Guo H, Brewer JM, Champhekar A, Harris RB, Bittman EL 2005 Differential control of peripheral circadian rhythms by suprachiasmatic-dependent neural signals. *Proc Natl Acad Sci U S A* 102:3111-6
13. Balsalobre A, Marcacci L, Schibler U 2000 Multiple signaling pathways elicit circadian gene expression in cultured Rat-1 fibroblasts. *Curr Biol* 10:1291-4.
14. Brown SA, Schibler U 1999 The ins and outs of circadian timekeeping. *Curr Opin Genet Dev* 9:588-94.
15. Hastings MH, Reddy AB, Maywood ES 2003 A clockwork web: circadian timing in brain and periphery, in health and disease. *Nat Rev Neurosci* 4:649-61.
16. Reppert SM, Weaver DR 2001 Molecular analysis of mammalian circadian rhythms. *Annu Rev Physiol* 63:647-76.
17. Lowrey PL, Takahashi JS 2000 Genetics of the mammalian circadian system: Photic entrainment, circadian pacemaker mechanisms, and posttranslational regulation. *Annu Rev Genet* 34:533-562.
18. Ripperger JA, Shearman LP, Reppert SM, Schibler U 2000 CLOCK, an essential pacemaker component, controls expression of the circadian transcription factor DBP. *Genes Dev* 14:679-89.
19. Ueda HR, Hayashi S, Chen W, Sano M, Machida M, Shigeyoshi Y, Iino M, Hashimoto S 2005 System-level identification of transcriptional circuits underlying mammalian circadian clocks. *Nat Genet* 37:187-92.
20. Shearman LP, Sriram S, Weaver DR, Maywood ES, Chaves I, Zheng B, Kume K, Lee CC, van der Horst GT, Hastings MH, Reppert SM 2000 Interacting molecular loops in the mammalian circadian clock. *Science* 288:1013-9.

21. Marquez S, Crespo P, Carlini V, Garbarino-Pico E, Baler R, Caputto BL, Guido ME 2004 The metabolism of phospholipids oscillates rhythmically in cultures of fibroblasts and is regulated by the clock protein PERIOD 1. *Faseb J* 18:519-21.
22. Oishi K, Miyazaki K, Kadota K, Kikuno R, Nagase T, Atsumi G, Ohkura N, Azama T, Mesaki M, Yukimasa S, Kobayashi H, Iitaka C, Umehara T, Horikoshi M, Kudo T, Shimizu Y, Yano M, Monden M, Machida K, Matsuda J, Horie S, Todo T, Ishida N 2003 Genome-wide expression analysis of mouse liver reveals CLOCK-regulated circadian output genes. *J Biol Chem* 278:41519-27.
23. Akhtar RA, Reddy AB, Maywood ES, Clayton JD, King VM, Smith AG, Gant TW, Hastings MH, Kyriacou CP 2002 Circadian cycling of the mouse liver transcriptome, as revealed by cDNA microarray, is driven by the suprachiasmatic nucleus. *Curr Biol* 12:540-50.
24. Rudic RD, McNamara P, Curtis AM, Boston RC, Panda S, Hogenesch JB, Fitzgerald GA 2004 BMAL1 and CLOCK, two essential components of the circadian clock, are involved in glucose homeostasis. *PLoS Biol* 2:e377.
25. Turek FW, Joshu C, Kohsaka A, Lin E, Ivanova G, McDearmon E, Laposky A, Losee-Olson S, Easton A, Jensen DR, Eckel RH, Takahashi JS, Bass J 2005 Obesity and metabolic syndrome in circadian Clock mutant mice. *Science* 308:1043-5
26. Inoue I, Shinoda Y, Ikeda M, Hayashi K, Kanazawa K, Nomura M, Matsunaga T, Xu H, Kawai S, Awata T, Komoda T, Katayama S 2005 CLOCK/BMAL1 is involved in lipid metabolism via transactivation of the peroxisome proliferator-activated receptor (PPAR) response element. *J Atheroscler Thromb* 12:169-74
27. Oishi K, Shirai H, Ishida N 2005 CLOCK is involved in the circadian transactivation of peroxisome-proliferator-activated receptor alpha (PPARalpha) in mice. *Biochem J* 386:575-81.
28. Shimba S, Ishii N, Ohta Y, Ohno T, Watabe Y, Hayashi M, Wada T, Aoyagi T, Tezuka M 2005 Brain and muscle Arnt-like protein-1 (BMAL1), a component of the molecular clock, regulates adipogenesis. *Proc Natl Acad Sci U S A* 102:12071-6
29. Raspe E, Duez H, Gervois P, Fievet C, Fruchart JC, Besnard S, Mariani J, Tedgui A, Staels B 2001 Transcriptional regulation of apolipoprotein C-III gene expression by the orphan nuclear receptor RORalpha. *J Biol Chem* 276:2865-71.
30. Vu-Dac N, Chopin-Delannoy S, Gervois P, Bonnelye E, Martin G, Fruchart JC, Laudet V, Staels B 1998 The nuclear receptors peroxisome proliferator-activated receptor alpha and Rev-erbalpha mediate the species-specific regulation of apolipoprotein A-I expression by fibrates. *J Biol Chem* 273:25713-20.
31. Vu-Dac N, Gervois P, Grotzinger T, De Vos P, Schoonjans K, Fruchart JC, Auwerx J, Mariani J, Tedgui A, Staels B 1997 Transcriptional regulation of apolipoprotein A-I gene expression by the nuclear receptor RORalpha. *J Biol Chem* 272:22401-4.
32. Hirota T, Okano T, Kokame K, Shirotani-Ikejima H, Miyata T, Fukada Y 2002 Glucose down-regulates Per1 and Per2 mRNA levels and induces circadian gene expression in cultured Rat-1 fibroblasts. *J Biol Chem* 277:44244-51.
33. Balsalobre A, Brown SA, Marcacci L, Tronche F, Kellendonk C, Reichardt HM, Schutz G, Schibler U 2000 Resetting of circadian time in peripheral tissues by glucocorticoid signaling. *Science* 289:2344-7.
34. McNamara P, Seo SP, Rudic RD, Sehgal A, Chakravarti D, FitzGerald GA 2001 Regulation of CLOCK and MOP4 by nuclear hormone receptors in the vasculature: a humoral mechanism to reset a peripheral clock. *Cell* 105:877-89.
35. Lemberger T, Saladin R, Vazquez M, Assimacopoulos F, Staels B, Desvergne B, Wahli W, Auwerx J 1996 Expression of the peroxisome proliferator-activated receptor

- alpha gene is stimulated by stress and follows a diurnal rhythm. *J Biol Chem* 271:1764-9.
36. Ueda HR, Chen W, Adachi A, Wakamatsu H, Hayashi S, Takasugi T, Nagano M, Nakahama K, Suzuki Y, Sugano S, Iino M, Shigeyoshi Y, Hashimoto S 2002 A transcription factor response element for gene expression during circadian night. *Nature* 418:534-9
  37. Lemberger T, Staels B, Saladin R, Desvergne B, Auwerx J, Wahli W 1994 Regulation of the peroxisome proliferator-activated receptor alpha gene by glucocorticoids. *J Biol Chem* 269:24527-30
  38. Steineger HH, Sorensen HN, Tugwood JD, Skrede S, Spydevold O, Gautvik KM 1994 Dexamethasone and insulin demonstrate marked and opposite regulation of the steady-state mRNA level of the peroxisomal proliferator-activated receptor (PPAR) in hepatic cells. Hormonal modulation of fatty-acid-induced transcription. *Eur J Biochem* 225:967-74.
  39. Shaw D, Goldman BD 1995 Gender differences in influence of prenatal photoperiods on postnatal pineal melatonin rhythms and serum prolactin and follicle-stimulating hormone in the Siberian hamster (*Phodopus sungorus*). *Endocrinology* 136:4237-46.
  40. Patel DD, Knight BL, Wiggins D, Humphreys SM, Gibbons GF 2001 Disturbances in the normal regulation of SREBP-sensitive genes in PPAR alpha-deficient mice. *J Lipid Res* 42:328-37.
  41. Gibbons GF, Patel D, Wiggins D, Knight BL 2002 The functional efficiency of lipogenic and cholesterogenic gene expression in normal mice and in mice lacking the peroxisomal proliferator-activated receptor-alpha (PPAR-alpha). *Adv Enzyme Regul* 42:227-47.
  42. Kersten S, Seydoux J, Peters JM, Gonzalez FJ, Desvergne B, Wahli W 1999 Peroxisome proliferator-activated receptor alpha mediates the adaptive response to fasting. *J Clin Invest* 103:1489-98.
  43. Gervois P, Chopin-Delannoy S, Fadel A, Dubois G, Kosykh V, Fruchart JC, Najib J, Laudet V, Staels B 1999 Fibrates increase human REV-ERBalpha expression in liver via a novel peroxisome proliferator-activated receptor response element. *Mol Endocrinol* 13:400-9.
  44. Knight BL, Hebbachi A, Hauton D, Brown AM, Wiggins D, Patel DD, Gibbons GF 2005 A role for PPARalpha in the control of SREBP activity and lipid synthesis in the liver. *Biochem J* 389:413-21.
  45. Desvergne B, Michalik L, Wahli W 2004 Be fit or be sick: peroxisome proliferator-activated receptors are down the road. *Mol Endocrinol* 18:1321-32.
  46. Gachon F, Nagoshi E, Brown SA, Ripperger J, Schibler U 2004 The mammalian circadian timing system: from gene expression to physiology. *Chromosoma* 113:103-12
  47. Canaple L, Kakizawa T, Laudet V 2003 The days and nights of cancer cells. *Cancer Res* 63:7545-52
  48. Rosbash M 1998 Why the rat-1 fibroblast should replace the SCN as the in vitro model of choice. *Cell* 93:917-9
  49. Yagita K, Tamanini F, van Der Horst GT, Okamura H 2001 Molecular mechanisms of the biological clock in cultured fibroblasts. *Science* 292:278-81
  50. Hastings MH 2005 Circadian biology: fibroblast clocks keep ticking. *Curr Biol* 15:R16-8

51. Triqueneaux G, Thenot S, Kakizawa T, Antoch MP, Safi R, Takahashi JS, Delaunay F, Laudet V 2004 The orphan receptor Rev-erb $\alpha$  gene is a target of the circadian clock pacemaker. *J Mol Endocrinol* 33:585-608.
52. Yu Y, Correll PH, Vanden Heuvel JP 2002 Conjugated linoleic acid decreases production of pro-inflammatory products in macrophages: evidence for a PPAR gamma-dependent mechanism. *Biochim Biophys Acta* 1581:89-99
53. Kondratov RV, Chernov MV, Kondratova AA, Gorbacheva VY, Gudkov AV, Antoch MP 2003 BMAL1-dependent circadian oscillation of nuclear CLOCK: posttranslational events induced by dimerization of transcriptional activators of the mammalian clock system. *Genes Dev* 17:1921-32
54. Ogawa H, Ansai Y 1995 Diurnal rhythms of rat liver serine dehydratase, D-site binding protein, and 3-hydroxy-3-methylglutaryl coenzyme A reductase mRNA levels are altered by destruction of the suprachiasmatic nucleus of the hypothalamus. *Arch Biochem Biophys* 321:115-22
55. Lavery DJ, Schibler U 1993 Circadian transcription of the cholesterol 7  $\alpha$  hydroxylase gene may involve the liver-enriched bZIP protein DBP. *Genes Dev* 7:1871-84
56. Lavery DJ, Lopez-Molina L, Margueron R, Fleury-Olela F, Conquet F, Schibler U, Bonfils C 1999 Circadian expression of the steroid 15  $\alpha$ -hydroxylase (Cyp2a4) and coumarin 7-hydroxylase (Cyp2a5) genes in mouse liver is regulated by the PAR leucine zipper transcription factor DBP. *Mol Cell Biol* 19:6488-99
57. Vosper H, Khoudoli GA, Graham TL, Palmer CN 2002 Peroxisome proliferator-activated receptor agonists, hyperlipidaemia, and atherosclerosis. *Pharmacol Ther* 95:47-62.
58. Palmer CN, Hsu MH, Griffin KJ, Raucy JL, Johnson EF 1998 Peroxisome proliferator activated receptor-alpha expression in human liver. *Mol Pharmacol* 53:14-22
59. Macdonald N, Holden PR, Roberts RA 1999 Addition of peroxisome proliferator-activated receptor alpha to guinea pig hepatocytes confers increased responsiveness to peroxisome proliferators. *Cancer Res* 59:4776-80
60. IJpenberg A, Tan NS, Gelman L, Kersten S, Seydoux J, Xu J, Metzger D, Canaple L, Chambon P, Wahli W, Desvergne B 2004 In vivo activation of PPAR target genes by RXR homodimers. *Embo J* 23:2083-91.
61. Podvinec M, Kaufmann MR, Handschin C, Meyer UA 2002 NUBIScan, an in silico approach for prediction of nuclear receptor response elements. *Mol Endocrinol* 16:1269-79.




Figure 1






Figure 2




Figure 3







Figure 4

A.



B.




Figure 5




Figure 6