

Low doses of GM-CSF (molgramostim) and G-CSF (filgrastim) after cyclophosphamide (4 g/m²) enhance the peripheral blood progenitor cell harvest: results of two randomized studies including 120 patients.

Philippe Quittet, Patrice Ceballos, Ernesto Lopez, Zhao-Yang Lu, Pascal Latry, Catherine Becht, Eric Legouffe, Nathalie Fegueux, Carole Exbrayat, Damien Pouessel, et al.

► **To cite this version:**

Philippe Quittet, Patrice Ceballos, Ernesto Lopez, Zhao-Yang Lu, Pascal Latry, et al.. Low doses of GM-CSF (molgramostim) and G-CSF (filgrastim) after cyclophosphamide (4 g/m²) enhance the peripheral blood progenitor cell harvest: results of two randomized studies including 120 patients.. Bone Marrow Transplantation, Nature Publishing Group, 2006, 38 (4), pp.275-84. 10.1038/sj.bmt.1705441 . inserm-00136076

HAL Id: inserm-00136076

<https://www.hal.inserm.fr/inserm-00136076>

Submitted on 13 Mar 2007

HAL is a multi-disciplinary open access archive for the deposit and dissemination of scientific research documents, whether they are published or not. The documents may come from teaching and research institutions in France or abroad, or from public or private research centers.

L'archive ouverte pluridisciplinaire **HAL**, est destinée au dépôt et à la diffusion de documents scientifiques de niveau recherche, publiés ou non, émanant des établissements d'enseignement et de recherche français ou étrangers, des laboratoires publics ou privés.

Low Doses of GM-CSF (Molgramostim) and G-CSF (Filgrastim) after Cyclophosphamide (4g/m²) Enhances the Peripheral Blood Progenitor Cell Harvest : Results of two Randomized Studies including 120 patients.

Quittet P¹, Ceballos P¹, Lopez E¹, Lu ZY², Latry P¹, Becht C¹, Legouffe E¹, Fegueux N¹, C Exbrayat¹, Pouessel D¹, V Rouillé¹, Daures JP³, Klein B², Rossi JF^{1*}.

1, Service Hématologie et Oncologie médicale, Hopital Lapeyronie, 34295 Montpellier, France.

2, Unité de Thérapie Cellulaire, Hopital St-Eloi, 34295 Montpellier, France.

3, Unité de Biostatistiques, IURC Montpellier.

- to whom correspondence should be sent :

E-mail address: jf-rossi@chu-montpellier.fr

Running Title : G-CSF versus G-CSF plus GM-CSF for mobilization/collection of PBPC

ABSTRACT

The use of a combination of G-CSF and GM-CSF to G-CSF alone, after cyclophosphamide (4g/m^2) was compared in 2 randomized phase III studies, including 120 patients. In study A, 60 patients received $5 \times 2 \mu\text{g/kg/day}$ of G-CSF and GM-CSF compared to $5 \mu\text{g/kg/day}$ of G-CSF. In study B, 60 patients received $2.5 \times 2 \mu\text{g/kg/day}$ G-CSF and GM-CSF compared to G-CSF alone ($5 \mu\text{g/kg/day}$). With the aim to collect at least $5 \times 10^6/\text{kg}$ CD34 cells in a maximum of 3 large volume leukapheresis (LK), 123 LK were performed in study A, showing significant higher number of patients reaching $10 \times 10^6/\text{kg}$ CD34 cells (21/29 in G+GM-CSF arm vs 11/27 in G-CSF arm, $P = .00006$). In study B, 109 LK were performed, with similar results (10/27 vs 15/26, $P = .003$). In both the study, the total harvest of CD34 cells/kg was 2-fold higher in G-CSF plus GM-CSF group (18.3×10^6 in study A and 15.85×10^6 in study B) than in G-CSF group (9×10^6 in study A and 8.1×10^6 in study B), a difference particularly seen in multiple myeloma, with no significant difference in terms of mobilized myeloma cells between G-CSF and GM-CSF groups.

Key words: G-CSF, GM-CSF, hematopoietic stem cells.

INTRODUCTION

Autologous peripheral blood progenitor cells (PBPC) provide a rapid and sustained hematopoietic recovery after the administration of high dose therapy in patients with haematological malignancies and certain solid tumors (1-5). The mobilization of PBPC is usually achieved with the use of haematopoietic growth factors (HGFs) alone or in combination with chemotherapy, in which case a higher yield of CD34+ cells can be reached and collected (6). It is generally recognized that granulocyte-colony-stimulating factor (G-CSF) as a single agent mobilizes more CD34+ cells than does granulocyte-macrophage-colony stimulating factor (GM-CSF) (7,8 and for review, 9). Other HGFs, such as Flt3 ligand (10), interleukin-3 (11), stem cell factor (12-13) and erythropoietin (14) or antagonists of SDF-1-CXCR4 (15) have been tested generally in combination for mobilizing PBPC. The recommended dose of G-CSF after chemotherapy is 5µg/kg/d (16). Increasing the dose of G-CSF superior to 5µg/kg/d (17, 18) or changing the administration schedule (19) has not been proven to improve PBPC mobilization. More recently, the use of a single dose of pegfilgrastin was demonstrated as equivalent to a daily administration of filgrastim for mobilizing PBPC (20). It has been shown the therapeutic relevance of high dose of CD34 cells in autologous PBPC transplantation. The infusion of 5×10^6 CD34/kg minimum results in rapid engraftment, reduces the transfusion need and may have a clinical influence (21-23). The synergistic effect of co-administration of HGFs like G-CSF and GM-CSF on PBPC mobilization has been suggested in phase I/II study (24), in order to improve the harvest of CD34 cells (25). Several non-randomized or randomized clinical trials have been performed, showing a little or no benefit for sequential administration of standard doses (5-10 µg/kg/d) of GM-CSF and G-CSF (26-37). Nevertheless, the use of these HGFs was not well explored and particularly the minimal efficient dose for their concomitant administration, following high dose cyclophosphamide (CY). We report here two randomized studies in order to

determine first, whether the concomitant administration of GM-CSF and G-CSF could improve the PBPC mobilization and collection to achieve a target yield of 5×10^6 CD34 cells/kg, and secondly what is the minimal dose for this combination of HGFs.

MATERIALS AND METHODS

Study design.

As shown on Fig.1, this was a randomized, open-label, unicenter study, including two parts (study A and study B). The primary objective of this study was to achieve 10×10^6 CD34 cells/kg in a minimal number of leukapheresis. Calculation of the number of subjects was based on the fact that the percentage of the subjects reaching more than 10×10^6 CD34 cells/kg in the combination arm will be twice than that observed in the single-cytokine arm. With a power of 90% and a 5% risk, the number of patients is 25 per arm. For that reason, we decide to include 30 patients per arm due to the possibility of unevaluable patients. There was no stratification. The secondary objectives of this study include the tolerance, the factors influencing the mobilization of progenitor cells, the percentage of myeloma cells mobilized in PB, and the difference of mobilization at the level of 5×10^6 CD34 cells/kg between the 2 arms. In study A, 60 patients were randomized to receive after CY (4 g/m^2) the co-administration of $5 \text{ } \mu\text{g/kg/day}$ (d) of rHu-GM-CSF (Molgramostim, Schering-Plough, Kenilworth, NJ, USA) and $5 \text{ } \mu\text{g/kg/d}$ of rHu-G-CSF (Filgrastim, Amgen, Thousand Oaks, CA) (G + GM-CSF, total HGF dose = $10 \text{ } \mu\text{g/kg/d}$) compared to $5 \text{ } \mu\text{g/kg/d}$ of G-CSF (Filgrastim). In study B, 60 subsequent patients have received the same design of treatment with $2.5 \text{ } \mu\text{g/kg/d}$ of GM-CSF and $2.5 \text{ } \mu\text{g/kg/d}$ of G-CSF (G + GM-CSF, total dose of HGF = $5 \text{ } \mu\text{g/kg/d}$) compared to $5 \text{ } \mu\text{g/kg/d}$ of G-CSF. The 4 g/m^2 CY was administered by intravenous (IV) route to all the patients followed 24 hours later by the HGF administration subcutaneously until the last day of leukapheresis (LK). For patients receiving the 2 HGFs, they were injected in separate sites.

Patient Eligibility

Written informed consent was obtained from all patients enrolled in the study. Patients with histologically confirmed history of cancer and requiring a high dose myeloablative

chemotherapy with PBPC rescue were eligible. Disease status and demographic characteristics are detailed in Table I. Inclusion criteria were: age between 18 and 65 years, performance status ≤ 2 , half-life expectancy of at least 6 months, normal organ function as defined by serum creatinine, transaminase < 2 times normal range, cardiac ejection fraction within the institution's normal range, and normal blood count (ANC $> 1.5 \times 10^9/L$, platelets $> 100 \times 10^9/L$) at the day of mobilization chemotherapy. Exclusion criteria were: patients who had received HGF within 2 weeks before study entry, patients with other malignancies within 5 past years, HIV1 or 2, HTLV 1 or 2, hepatitis C sero-positivity, hepatitis B positive virological status, patients with psychiatric, addictive or any disorder which compromised their ability to give truly informed consent.

Stem cell collection.

White blood count (WBC) were assessed daily after the beginning of chemotherapy. When it reaches $0.5 \times 10^9/L$, the percentage of circulating CD34 cells was monitored daily according to the ISHAGE protocol, by using a phycoerythrin-conjugated (PE) antibody to CD34 (Q-Bend 10, Immunotech, Marseille, France) (38). For each labelled sample, 30,000 events were recorded on forward-versus-side scatter dot-plot using a FACScan (Becton-Dickinson, San-Jose, CA, USA). The estimated number of CD34 cells/kg that can be collected in one LK was calculated by the central laboratory according to the following formula: $(\%CD34 \times WBC \times 8/kg \text{ body weight})$ (8 being a coefficient corresponding approximately to 2 blood volumes in L). Then, a LK was started if the estimation of harvestable CD34 number was $\geq 10^6$ CD34/kg. If the estimation of harvestable CD34 cell number was below 10^6 CD34/kg the collection was delayed. A maximum of three LK was attempted to reach a minimum target of 5×10^6 CD34/kg. For MM patients, at this time, we performed a CD34 selection in order to decrease contaminating tumoral cells. For that reason, due to the loss of cells during this procedure, we attempt to collect 10×10^6 CD34/kg. Large volume LK was performed through a dual

peripheral venous puncture using a Cobe Spectra separator version 4 (CS 3000; Baxter Healthcare Corp Lakewood, Co). Median time of processing was 5 hours and the mean volume of processed blood ranged between 12 to 16 L.

Analysis of product of leukapheresis.

The determination of CD34 cell content in the LK was carried out according to the following procedure. Incubation of 10^6 cells with 30% AB serum (100 μ L) for 10 mn, two washings with PBS, suspension in a final volume of 100 μ L, incubation with 20 μ L of monoclonal antibody CD34 conjugated with PE, or 20 μ L of murine IgG recognizing no human antigen and conjugated with PE as a negative control, or 5 μ L anti-CD 45 conjugated with PE as a positive control for 30 mn at 4°C, cell lysis followed by two washings and resuspension in 200 μ L PBS. Cells are analysed with a FACSCAN cytometer. At least 30,000 events were acquired for each sample.

In patients with Multiple Myeloma (MM), myeloma cells were enumerated by FACS analysis using the FITC-conjugated MI15 (anti-CD138 mAb (39)). 10^6 cells were incubated with 1 μ g MI15^{FITC} (10 μ L) or to 1 μ g IgG^{FITC} negative control (20 μ L) and fluorescence was analysed with FACSCAN cytofluorometer. No assesment of CD34 cells was performed on the LK product.

Autologous stem cell transplantation (ASCT).

After a resting time of a maximum of 8 weeks, patients received high dose chemotherapy (HDC) followed by autologous PBPC transplant. MM patients were scheduled to receive two subsequent HDC and PBPC transplants. The first HDC was 140 mg/m² Melphalan. At least three months later, patients received the second HDC consisting of 200 mg/m² melphalan or 140mg/m² melphalan and total body irradiation delivering 12 Gy followed by 5 μ g/kg/d G-CSF support administrated from day 2 after HDC and until ANC > 1.5 x 10⁹/L on two consecutive counts. Patients having malignant lymphoma received a BEAM regimen

consisting on 300 mg/m² BCNU, Etoposide 200 mg/m²/d for four days, Cytarabine 200 mg/m²/d for four days and Melphalan 140 mg/m² without post-transplant G-CSF support. Breast cancer, testis and ovarian cancer patients were scheduled to receive one or two HDC and PBPC transplant. The following parameters of haematologic recovery were analyzed: day of ANC $\geq 0.5 \times 10^9/L$, day of platelet count $\geq 50 \times 10^9/L$, cost estimation of the whole transfusion procedure and duration of hospitalization.

Statistical analysis.

An intent-to-treat analysis was done taking into account all patients who received CY and HGFs. A Wilcoxon non-parametric test was carried out to assess the comparability of group at randomization. The comparison of number of LK per group were evaluated by an exact Fischer's test. The median values of CD34 cells harvested in the different groups were compared by a Mann-Whitney non-parametric test. The comparison of distribution of patients with 1 to 3 LK to reach the pre-defined threshold was done by the Chi 2 test. If none LK was performed it is considered as mobilization failure and not evaluated. Data were analysed using SAS (Statistical Analysis System) software version 6.08.

RESULTS

Patient population (Table I)

One hundred twenty consecutive patients were enrolled and randomized in the department. All patients had a disease requiring high-dose chemotherapy and PBPC transplantation as salvage therapy. They were: 58 (48%) MM patients (stage II or III), 32 (27%) non-Hodgkin Lymphoma (NHL) (large cell type with IPI \geq 3 or sensitive relapse), 8 (7%) Hodgkin disease (HD) (sensitive relapse or refractory), and 4 (3%) chronic lymphocytic leukemia (CLL) (Binet stage B and C), and 18 (15%) solid tumors (ST) including 13 metastatic (or high-risk) breast cancer, 4 metastatic testis cancer, 1 metastatic ovarian cancer. All patients were newly diagnosed and received a range of 3 to 16 courses of chemotherapy before PBSC mobilization lasting less than one year.

Sixty patients were included in study A (5 μ g/kg/d G-CSF plus 5 μ g/kg/d GM-CSF versus 5 μ g/kg/d G-CSF): 26 MM, 12 NHL, 6 HD, 2 CLL, 14 ST.

Sixty subsequent patients were included in study B (2.5 μ g/kg/d G-CSF plus 2.5 μ g/kg/d GM-CSF versus 5 μ g/kg/d G-CSF): 32 MM, 20 NHL, 2 HD, 2 CLL, 4 ST.

4 were not evaluable for the following events at time of chemotherapy: infection at time of chemotherapy (1 case), disease progression or relapses (2 cases), death before mobilization (1 case). Patients characteristics were comparable, particularly age, sex ratio, body weight and blood cell count before mobilization.

Kinetics of mobilization of circulating CD34 PBPC

In study A, the circulating CD34 cells appear in the peripheral blood (PB) respectively between day 9 to 12 in the G-CSF group and day 9 to 13 in G-CSF plus GM-CSF group. In study B, the circulating CD34 cells appear between day 9 to 11 in the G-CSF group and day 9 to 13 in G-CSF plus GM-CSF group (data not shown).

Collection of PB CD34 cell

There was a good correlation between the estimated number of CD34 cells and the collected number of CD34 cells measured (Fig. 2). Parameters of LK procedures are shown in Table II. A total of 232 cytaphereses were performed: 123 in study A, and 109 in study B.

One to three large volume LK were performed to harvest the required number of CD34 cells. The following parameters of LK were analysed: number of LK processed per patient (LK/patient), the day of first LK after CY, WBC at the first LK, and the median number of blood volumes treated per LK (BV/LK). None of these parameters were statistically different between the randomized groups.

Number of patients achieving a harvest of 5×10^6 and 10×10^6 CD34 cells/kg

In study A, 123 LK were performed (61 in the G-CSF group, 62 in the G + GM-CSF group). The distribution of patients undergoing one, two or three LK were identical in both groups. Among the patients who have undergone LK, the proportion of patients reaching a minimum target of 5×10^6 CD34 cells/kg was not statistically different between the G-CSF arm (21/27, 77.8%) and the G-CSF plus GM-CSF arm (25/29, 86.2%). When the objective of CD34 collection was 10×10^6 cells/kg, the proportion of patients reaching this target was significantly higher in the G-CSF plus GM-CSF arm (21/29, 72.4%) compared to the G-CSF arm (11/27, 40.7%; $P = .00006$).

In study B, a total of 109 LK were performed (56 in G-CSF arm and 53 in G-CSF plus GM-CSF arm). The distribution of patient undergoing one, two or three LK are identical in both groups. Among the patients who have undergone LK, the proportion of patients reaching a minimum target of 5×10^6 CD34 cells/kg was not statistically different between the G-CSF arm (20/27, 74.1%) and the G-CSF plus GM-CSF arm (20/26, 76.9%). When the objective of CD34 collection was 10×10^6 cells/kg, the proportion of patients reaching this target was significantly higher in the G-CSF plus GM-CSF (arm 15/26, 57.7%) compared to the G-CSF arm (10/27, 37%; $P = .003$) (Fig 3).

CD34 cell harvest (Table III).

In study A, the median number of CD34 cells x 10⁶/kg collected in LK1, LK2 or LK3 in the G-CSF and G + GM-CSF groups is respectively 3.1 (range 0.9-44.7), 3.7 (1.2-13.1), 2.7 (1.6-7.6) versus 7.3 (1-30.9), 6.28 (0.8-59), 5.72 (0.8-12.5). The total number of CD34 cells x10⁶/kg harvested is 2 fold higher in the G + GM-CSF group 18.3 (3-90) compared to 9 (3-45) in the G-CSF group (P= .09).

In study B, the median number of CD34 cells x 10⁶/kg collected at LK1, LK2, LK3 in the G-CSF and G + GM-CSF groups is respectively 3.7 (range 0.6-30.1), 3.0 (0.6-14.3), 3.3 (2.9-7) versus 7.07 (0.9-24.5), 8.02 (0.7-29.8), 6.71 (0.6-9.3). The total number of CD34 x 10⁶/kg harvested is about 2 fold higher in the G + GM-CSF group 15.85 (2.5-34.9) compared to 8.1 (1.6-33.9) in the G-CSF group (P= .09).

To avoid a bias due to the variable number of LK, and due to the variable blood volume treated during one LK, we have evaluated the efficacy of CD34 PBPC collection by calculating the number of CD34 cells x 10⁶/kg collected during the first LK according to the number of blood volume processed (CD34 LK₁/BV). In study A, this parameter was again increased about 2 fold in the G-CSF plus GM-CSF group 2.4 (0.3-13) compared to 1.1 (0.3-12) in the G-CSF group (P= .02). In study B, the same observation could be done in the G + GM-CSF group: 2.52 (0.3-10.2) versus 1.37 (0.2-11.6) in the G-CSF group (P= 0.04).

Patients with MM

In the study A : 14 MM patients received G-CSF and 12 received G-CSF plus GM-CSF (11 are evaluable). The total CD34 cells x 10⁶/kg harvested is 2.2 fold higher in the G-CSF plus GM-CSF group compared to the G-CSF group respectively 23.8 (11.1-89.9) versus 11.02 (3.2-29.2) (P= .005). The CD34 LK₁/BV is 2.4 fold higher in the G-CSF plus GM-CSF group 2.81 (0.9-13.3) versus 1.17 (0.3-8.1) in the G-CSF group (p=0.02). In the MM patients, we recommended to collect a minimum target of 10 x 10⁶ CD34 cells/kg before the CD34

positive selection. In these conditions, 11/11 (100%) of patients receiving the G-CSF plus GM-CSF combination reach this target compared to only 8/14 (57%) in the G-CSF group ($P = .02$) (Fig 4). The same observation were done in patients of study B, 17 MM patients received G-CSF (15 are evaluable) and 15 received G-CSF plus GM-CSF (14 are evaluable). The total CD34 cells $\times 10^6$ /kg harvested is about 2.5 fold higher in the G-CSF plus GM-CSF group (24.6 , range: 2.5-34.9) compared to the G-CSF group (9.26, range: 2.8-33.9) ($P = .03$). The CD34 LK₁/BV is 2.2 fold higher in the G-CSF plus GM-CSF group 2.95×10^6 /kg (0.4-10.2) compared to 1.35×10^6 /kg (0.1-11.6) ($P = .014$). 12/14 (86%) of patients receiving the G-CSF plus GM-CSF combination reach the defined target of 10×10^6 CD34+ cells/kg compared to only 7/15 (44%) in the G-CSF group ($P = .03$) (Fig. 4).

Patients with other lymphoid malignancies (NHL, HD and CLL)

Mobilization by CY and G-CSF plus GM-CSF did not significantly improve the amount of collected CD34 cells compared to the CY and G-CSF mobilization especially the CD34 LK₁/BV was not significantly increased (Table III). The comparison between myeloma and lymphoma patients is detailed in Table IV. MM patients received less chemotherapy courses (median 3, range: 2-6) while lymphoma patients received a median of 6 courses (median 2-16) chemotherapy courses ($P = .02$). As shown on Fig 5, no differences were seen between the two arms (G-CSF and G+GM-CSF) in the group of patients with more than one year of chemotherapy before mobilization.

In study A, among patients receiving G-CSF, the CD34 LK₁/BV is not significantly higher in MM 1.2×10^6 (0.3-8) than in Lymphoma 0.9×10^6 (0.4-12) ($P = .12$) while among patients receiving G-CSF plus GM-CSF, the CD34 LK₁/BV is 1.5 higher in MM (2.8×10^6 , range :1-13) than in Lymphoma (1.9×10^6 , range: 0.3-7) ($P = .047$). In study B, same observations was made, the CD34 LK₁/BV is not significantly higher in MM than in lymphoma ($P = .8$) while among patients receiving G-CSF plus GM-CSF, the CD34 LK₁/BV

is 2.1 higher in MM (2.95×10^6 , range: 0.4-10) than in Lymphoma (1.4×10^6 , range: 0.3-5) ($P = .01$).

Patients with solid tumors

Due to of low number of patients with solids tumors and their heterogenicity, none comparison was made between the two groups of treatment.

Mobilization failure

The percentage of patient that failed to mobilize sufficient PBSC was similar in study A (4/56, 7%) and in study B (3/53, 6%). None difference has been observed according to the HGF treatment group (Table V).

Adverse events

No serious adverse events were associated with the administration of cyclophosphamide at 4 g/m^2 . Grade 1 and 2 nausea and vomitis were easily controlled by systematic administration of andosetron. In study A, three patients with full dose of G + GM-CSF experienced grade 2 symptoms (chill, bone and muscle pains, tachycardia) within the first two days. These patients stopped themself the GM-CSF. No patient experienced grade 1 and 2 toxicities during HGF treatment in the study B.

Effects of HGF on residual plasma cells (PC) mobilization in MM patients

In study A, an estimation of residual PC in product of LK was possible in 19 patients. The number of PC was $4.6 \times 10^6/\text{kg}$ in the G-CSF group and $5.39 \times 10^6/\text{kg}$ in the G-CSF plus GM-CSF group ($P = \text{NS}$).

In study B, the numeration of residual PC in LK was done in 21 patients with MM. The number of PC was $2.58 \times 10^6/\text{kg}$ in the G-CSF group and $6.19 \times 10^6/\text{kg}$ in the G + GM-CSF group ($P = \text{NS}$).

When comparing the total PC collected to the total CD34 cells collected in each patient, the PC/CD34 ratio is in study A 1 for 2.4 in G-CSF group and 1 for 4.42 in the G-CSF plus

GM-CSF group (P = NS). In study B, the PC/CD34 ratio is 1 for 3.6 in G-CSF group and 1 for 3.97 in the G-CSF plus GM-CSF group (P = NS) showing no enhancement of PC mobilization by G-CSF plus GM-CSF regimen.

Hematologic recovery after myeloablative treatment

97 patients were transplanted: 49 patients in Study A, 48 in study B. All MM patients were scheduled to receive a transplantation with a graft purged by CD34 positive selection. In study A, the CD34 positive selection was done in only 9/14 (64.3%) MM patients receiving G-CSF and 11/11 (100%) in patients receiving G-CSF plus GM-CSF (P= .03). In study B, this regimen was done in only 8/15 (53.3%) patients receiving G-CSF and 13/14 (92.8%) patients receiving G-CSF plus GM-CSF (P= .02).

Because of purification and double HDC in MM, the median number of CD34 cells infused was similar in all group of treatment and ranged between 3.78 to 4.32 x 10⁶/kg. As expected, all patients had an engraftment, the median times to reach an ANC of $\geq 0.5 \times 10^9/L$ and platelet $\geq 50 \times 10^9/L$ were respectively 11 days (range 9 to 17) and 15 days (8-34). The transfusion cost is estimated at 1,619 \$ (265-8,752). The median duration of hospitalisation was 24 days (16-94).

DISCUSSION

These two randomized studies demonstrate two major points:

-the combination of GM-CSF plus G-CSF following CY results in a 2 fold enhancement of the number of PBPC, in a safe and efficient manner for an homogenous group of newly diagnosed patients;

-the concomitant administration of these two HGFs, had additive effects, as observed by the superiority of the combination at equivalent doses ($5\mu\text{g}/\text{kg}$ vs $2.5+2.5\mu\text{g}/\text{kg}$).

Most of the patients included in these studies, achieved a minimal target of 5×10^6 CD34/kg, but more patients receiving G-CSF plus GM-CSF achieved a target of 10×10^6 CD34/kg. This result was obtained with a median of 2 LK. No adverse effects related to LK procedures have been reported. Although CD34 cell dose is currently considered to be the preferred indicator of satisfactory engraftment, the optimal CD34 cell dose for autologous transplantation remains to be defined. A collection of a threshold value of at least 3 to 5×10^6 CD34/kg is recommended to insure rapid and successful engraftment and long term hematological recovery. A dose-response relationship is evident between the number of CD34 cells per kilogram infused and neutrophil and platelet engraftment kinetics, decrease hospital stay and cost (40-42). Infusion of very high dose superior to 15×10^6 CD34/kg was shown to be associated with a shorter time for platelet recovery compared to that observed for patients receiving between 2.5 and 15×10^6 CD34/kg (43). However, Dercksen et al found no difference for neutrophil recovery when the number of CD34+ cells were superior to 6×10^6 cells/kg (44). Recently, different groups observed a linkage between the number of CD34 harvested and reinfused, and survival or duration of the response in different diseases including breast cancer and multiple myeloma (45, 46). The mechanism of such effect remains uncertain, but may be linked to a high number of immunocompetent cells collected in addition to the stem cells, as suggested by Porrata et col. who observed a correlation between

progression-free survival and the number of lymphocytes readministered (46). In addition, GM-CSF had several immune activities, including differentiation of dendritic cells from monocytes and it has been associated to different strategies of immune therapy including anti-tumoral vaccination (46, 47). These observations may represent a rationale for re-discovering the role of GM-CSF for mobilizing both haematopoietic stem cells and immune cell effectors (48,49). The response we made through this study appears important to define the minimal efficient dose for a combination of G-CSF and GM-CSF, prior to add other compounds for mobilization such as selective CXCR4 antagonists (15).

We observed that the efficacy of dual HGFs was more evident in patients having MM, where 2.5 fold higher PBSC are collected with the combination of HGFs. However, no difference have been observed in the group of lymphoma's patients. The impressive results observed with MM compared to the lymphoma group, are probably due to prior chemotherapy compared to the lymphoma group. Some investigations have highlighted the importance of prior exposure to the specific stem cell toxicity of chemotherapy containing alkylating agent and/or anthracycline and their impact on mobilization efficacy (50). Haas R and col. suggest the same findings showing a loss of 0.2×10^6 CD34+ cells/kg per chemotherapy course including alkylating agent (51). In that way, it seems that 6 courses of chemotherapy appears as a significant critical threshold. This feature is probably not corrected by the HGF treatment, including the co-administration of G-CSF and GM-CSF. The dexamethasone administered preferentially to MM patients should also stimulate the hematopoiesis in synergy to HGFs. The role of glucocorticoids, especially dexamethasone, on proliferation of haematopoietic progenitors has been suggested *in vitro* (52, 53). This particular benefit observed in MM patients and the fact that we use a concomitant administration, may represent the differences observed between the literature and our results.

The timing of harvest is one of the major parameter of successful yield of CD34+ PBSC. The criteria used to define the optimal harvesting day was the peripheral WBC or mononuclear cells. But, these criterias are associated with a weak correlation to the CD34 cells amount in the product of leukapheresis (54). At the present time, the absolute number of circulating CD34 cells is correlated closely with the CD34 cell yield of the corresponding leukapheresis product. The usual critical threshold of circulating CD34 cells as indicator of onset of cytophoresis is 40 CD34 cells per μL (55). But, this method doesn't provide an estimation of CD34 cells collection. We use an estimation of the CD34 PBSC yield determined just before each cytophoresis to give directly an estimation of the final collect of CD34+ PBSCs.

In addition to PBSC mobilization, the mobilization of tumoral cells remains likely. The role of these residual tumoral cells in the graft for the relapse after transplantation is still controversial (56, 57). In this study, we evaluate the tumors cells co-mobilization in MM patients. We have shown no trend to mobilize more tumoral cells with G-CSF plus GM-CSF regimen than that observed with G-CSF alone. All patients recovered after myeloablative treatment.

As expected, no clear benefit appears in terms of engraftment, transfusion cost according to HGF regimen. The median number of CD34 cells infused to the patients in all treatment group is comparable (3.8 to 4.32×10^6 CD34+/kg). The majority of the patients having MM or breast cancer were scheduled to receive a double intensive chemotherapy followed by a CD34 immunoselected graft rescue. This schedule was possible more frequently in the G-CSF plus GM-CSF mobilization regimen (93 to 100% of patients) than in G-CSF group (53 to 64% of patients).

Future insight of graft manipulation as high level tumor cell purging, genetic manipulation, monocytes mobilization in view of anti-tumoral vaccination by dendritic cells, *ex-vivo*

expansion will invariably result in loss of a substantial fraction of hematopoietic progenitor cells harvested, a situation that needs to collect higher number of hematopoietic stem cells. The combination of cytokines, including GM-CSF may represent a particular interesting combination, in addition to the effects of GM-CSF on the immune system, as demonstrated in vaccination programs.

REFERENCES

1. Attal M , Harousseau JL, Stoppa AM et al. A prospective, randomized trial of autologous bone marrow transplantation and chemotherapy in multiple myeloma. *N Engl J Med* 1996; 335: 91-97.
2. Harousseau JL. Stem cell transplantation in multiple myeloma (0, 1, or 2). *Curr Opin Oncol* 2005 ; 17: 93-98.
3. Gorin NC. Autologous stem cell transplantation in hematological malignancies. *Springer Semin Immunopathol* 2004 ; 26: 3-30.
4. Philip T, Guglielmi C, Hagenbeek A et al. Autologous bone marrow transplantation as compared with salvage chemotherapy in relapses of chemotherapy-sensitive non-Hodgkin's lymphoma. *N Engl J Med*. 1995; 333: 1540-1545.
5. Gratwohl A, Schmid O, Baldomero H, et al. Accreditation Committee of the European Group for Blood and Marrow Transplantation. Haematopoietic stem cell transplantation (HSCT) in Europe 2002. Changes in indication and impact of team density. A report of the EBMT activity survey. *Bone Marrow Transplant* 34: 855-875, 2004.
6. Demirer T, Buckner CD, Bensinger WI. Optimization of stem cell mobilization. *Stem Cells* 1996; 14 : 106-116.
7. Lane TA, Law P, Maruyama et al. Harvesting and enrichment of HPC mobilized into the peripheral blood of normal donors by GM-CSF or G-CSF: potential role in allogeneic transplantation. *Blood* 1995; 85: 275-282.
8. Peters WP, Rosner G, Ross M et al. Comparative effect of GM-CSF and G-CSF on priming of peripheral blood progenitor cells for use with autologous bone marrow after high dose chemotherapy. *Blood* 1993, 81: 1709-1719.
9. Gazitt Y. Comparison between granulocyte colony-stimulating factor and granulocyte-macrophage colony-stimulating factor in the mobilization of peripheral blood stem cells. *Curr Opin Hematol* 2002, 9 : 190-198.
10. Lyman SD. Biologic effects and potential clinical applications of Flt3 ligand. *Curr Opin Hematol* 1998; 5: 192-196.
11. Haas R, Ehrhardt R, Witt B et al. Autografting with peripheral blood stem cells mobilized by sequential interleukin-3/granulocyte-macrophage colony-stimulating factor following high-dose chemotherapy in non-Hodgkin's lymphoma. *Bone Marrow Transplant* 1993; 12: 643-9.
12. Stiff P, Gingrich R, Luger S, A et al. Randomized phase 2 study of PBPC mobilization by stem cell factor and filgrastim in heavily pretreated patients with Hodgkin's disease or non-Hodgkin's lymphoma. *Bone Marrow Transplant* 2000; 26: 471-481.

13. Prosper F, Sola C, Hornedo J et al. Mobilization of peripheral blood progenitor cells with a combination of cyclophosphamide, r-metHuSCF and filgrastim in patients with breast cancer previously treated with chemotherapy. *Leukemia* 2003; 17: 437-441.
14. Waller CF, von Lintig F, Daskalakis A, et al. Mobilization of peripheral blood progenitor cells in patients with breast cancer: a prospective randomized trial comparing rhG-CSF with the combination of rhG-CSF plus rhEpo after VIP-E chemotherapy. *Bone Marrow Transplant.* 1999; 24: 19-24.
15. Liles WC, Rodger E, Broxmeyer HE, et al. Augmented mobilization and collection of CD34+ hematopoietic cells from normal human volunteers stimulated with granulocyte-colony-stimulating factor by single-dose administration of AMD3100, a CXCR4 antagonist. *Transfusion.* 2005; 45: 295-300.
16. Ozer H, Armitage JO, Bennett CL, et al. 2000 update of recommendations for the use of hematopoietic colony-stimulating factors: evidence-based, clinical practice guidelines. *J Clin Oncol* 2000; 18: 3558-3585.
17. Andre M, Baudoux E, Bron D, et al. Phase III randomized study comparing 5 or 10 microg per kg per day of filgrastim for mobilization of peripheral blood progenitor cells with chemotherapy, followed by intensification and autologous transplantation in patients with nonmyeloid malignancies. *Transfusion.* 2003; 43: 50-57.
18. Jones HM, Jones SA, Watts, MJ, et al. The effects of variation of the dose and schedule of G-CSF administration after low dose cyclophosphamide (1.5 g/m²) on both neutrophil and peripheral blood haemopoietic progenitor cell levels. *Blood* 1993; 82 (suppl 1) 233a.
19. Carrion R, Serrano D, Gomez-Pineda A, Diez-Martin JL. A randomised study of 10 microg/kg/day (single dose) vs 2 x 5 microg/kg/day (split dose) G-CSF as stem cell mobilisation regimen in high-risk breast cancer patients. *Bone Marrow Transplant* 2003; 32: 563-567.
20. Green MD, Koelbl H, Baselga J, et al. A randomized double-blind multicenter phase III study of fixed-dose single-administration pegfilgrastim versus daily filgrastim in patients receiving myelosuppressive chemotherapy. *Ann Oncol.* 2003; 14: 29-35.
21. Siena S, Schiavo R, Pedrazzoli P, Carlo-Stella C: Therapeutic relevance of CD34+ cell dose in blood cell transplantation for cancer therapy. *J Clin Oncol* 2000; 18:1360-1377.
22. Weaver CH, Hazelton B, Birch R, et al. An analysis of engraftment kinetics as a function of the CD34 content of peripheral blood progenitor cell collections in 692 patients after the administration of myeloablative chemotherapy. *Blood* 1995; 86: 3961-3969.

23. Schulman KA, Birch R, Zhen B, et al. Effect of CD34+ cell dose on resource utilization in patients after high-dose chemotherapy with peripheral-blood stem-cell support. *J Clin Oncol* 1999; 17: 1227.
24. Winter JN, Lazarus HM, Rademaker A, et al. Phase I/II study of combined granulocyte colony-stimulating factor and granulocyte-macrophage colony-stimulating factor administration for the mobilization of hematopoietic progenitor cells. *J Clin Oncol* 1996, 14: 277-286.
25. Charrier S, Chollet P, Bay JO, et al. Hematological recovery and peripheral blood progenitor cell mobilization after induction chemotherapy and GM-CSF plus G-CSF in breast cancer. *Bone Marrow Transplantation* 2000; 252: 705-710.
26. Bolwell B, Goormastic M, Yanssens T, et al. Comparison of G-CSF with GM-CSF for mobilizing peripheral blood progenitor cells and for enhancing marrow recovery after autologous bone marrow transplant. *Bone Marrow Transplant* 1994; 14: 913-918.
27. Spitzer G, Adkins D, Mathews M, et al. Randomized comparison of G-CSF + GM-CSF vs G-CSF alone for mobilization of peripheral blood stem cells: effects on hematopoietic recovery after high-dose chemotherapy. *Bone Marrow Transplant* 1997; 20: 921-930.
28. Weisdorf D, Miller J, Verfaillie C, et al. Cytokine-primed bone marrow stem cells vs peripheral blood stem cells for autologous transplantation : a randomized comparison of GM-CSF vs G-CSF. *Biol Blood Marrow Transplant* 1997, 3: 217-223.
29. Hohaus S, Martin H, Wassmann B, et al. Recombinant human granulocyte and granulocyte-macrophage colony-stimulating factor (G-CSF and GM-CSF) administered following cytotoxic chemotherapy have a similar ability to mobilize peripheral blood stem cells. *Bone Marrow Transplant* 1998; 22: 625-630.
30. Comenzo RL, Sanchowala V, Fisher C, et al. Intermediate-dose intravenous melphalan and blood stem cells mobilized with sequential GM+G-CSF or G-CSF alone to treat AL (amyloid light chain) amyloidosis. *Br J Haematol* 1999; 104 : 553-539.
31. Siena S, Bregni M, Gianni AM. Mobilization of peripheral blood progenitor cells for autografting: chemotherapy and G-CSF or GM-CSF. *Baillieres Best Pract Res Clin Haematol* 1999; 12: 27-39.
32. Madero L, Gonzalez-Vicent M, Molina J, et al. Use of concurrent G-CSF + GM-CSF vs G-CSF alone for mobilization of peripheral blood stem cells in children with malignant disease. *Bone Marrow Transplant* 2000; 26: 365-369.
33. Weaver CH, Schulman KA, Wilson-Relyea B, et al. Randomized trial of filgrastim, sargramostim, or sequential sargramostim and filgrastim after myelosuppressive

- chemotherapy for the harvesting of peripheral-blood stem cells. *J Clin Oncol* 2000; 18: 43-53.
34. Koc ON, Gerson SL, Cooper BW, et al. Randomized cross-over trial of progenitor-cell mobilization: high-dose cyclophosphamide plus granulocyte colony-stimulating factor (G-CSF) versus granulocyte-macrophage colony-stimulating factor plus G-CSF. *J Clin Oncol* 2000; 18: 1824-1830.
 35. Gazitt Y, Callander N, Freytes CO, et al. Peripheral blood stem cell mobilization with cyclophosphamide in combination with G-CSF, GM-CSF, or sequential GM-CSF/G-CSF in non-Hodgkin's lymphoma patients: a randomized prospective study. *J Hematother Stem Cell Res* 2000; 9: 737-748.
 36. Gazitt Y, Shaughnessy P, Liu Q. Differential mobilization of CD34+ cells and lymphoma cells in non-Hodgkin's lymphoma patients mobilized with different growth factors. *J Hematother Stem Cell Res* 2001; 10: 167-176.
 37. Armitage JO. Emerging applications of recombinant human granulocyte-macrophage colony-stimulating factor. *Blood* 1998; 92: 4491-4508.
 38. Sutherland DR, Anderson L, Keeney M, et al. The ISHAGE guidelines for CD34+ cell determination by flow cytometry. *International Society of Hematotherapy and Graft Engineering. J Hematother* 1996; 5: 213-226.
 39. Costes V, Magen V, Legouffe E, et al. The Mi15 monoclonal antibody (anti-syndecan-1) is a reliable marker for quantifying plasma cells in paraffin-embedded bone marrow biopsy specimens. *Hum Pathol* 1999; 30: 1405-1411.
 40. Duggan PR, Guo D, Luidier J, et al. Predictive factors for long-term engraftment of autologous blood stem cells. *Bone Marrow Transplant.* 2000; 26: 1299-1304.
 41. Uyl-de Groot CA, Huijgens PC, Rutten FF: Colony-stimulating factors and peripheral blood progenitor cell transplantation, benefits and costs. *Pharmaco Economics* 1996; 10: 23-35.
 42. Bensinger W, Appelbaum F, Rowley S, et al. Factors that influence collection and engraftment of autologous peripheral-blood stem cells. *J Clin Oncol* 1995; 13: 2547-2555.
 43. Ketterer N, Salles G, Raba M, et al. High CD34 (+) cell counts decrease hematologic toxicity of autologous peripheral blood progenitor cell transplantation. *Blood* 1998; 91: 3148-3155.
 44. Dercksen MW, Rodenhuis S, Dirkson MK, Schaasberg WP, Baars JW, van der Waal E, Slaper-Cortenbach IC, Pinedo HM, Von dem Borne AE, van der Schoot CE, Gerritsen WR: Subsets of CD34+ cells and rapid hematopoietic recovery after peripheral-blood stem-cell transplantation. *J Clin Oncol* 1995; 13: 1922-1932.

45. Wahlin A, Eriksson M, Hultdin M. Relation between harvest success and outcome after autologous peripheral blood stem cell transplantation in multiple myeloma. *Eur J Haematol.* 2004; 73: 263-268.
46. Porrata LF, Gertz MA, Geyer SM, et al. The dose of infused lymphocytes in the autograft directly correlates with clinical outcome after autologous peripheral blood hematopoietic stem cell transplantation in multiple myeloma. *Leukemia.* 2004; 18: 1085-1092.
47. O'Neil DW, Adams S, Bhardwaj N. Manipulating dendritic cell biology for the active immunotherapy cancer. *Blood* 2004; 104: 2235-2246.
48. Dranoff G. GM-CSF-based cancer vaccines. *Immunol Rev.* 2002; 188: 147-154.
49. Gazitt Y. Immunologic profiles of effector cells and peripheral blood stem cells mobilized with different hematopoietic growth factors. *Stem Cells.* 2000; 18: 390-398.
50. Koumakis G, Vassilomanolakis M, Hatzichristou H, et al. Predictive factors affecting mobilization and peripheral blood stem cell (PBSC) collection using single apheresis (SA) for rescuing patients after high-dose chemotherapy (HD.CHE) in various malignancies. *Bone Marrow Transplant* 1996; 18: 1065-1072.
51. Haas R, Möhle R, Frühauf S, Goldschmidt H, et al. Patient characteristics associated with successful mobilizing and autografting of peripheral blood progenitor cells in malignant lymphoma. *Blood* 1994; 83: 3787-3794.
52. von Lindern M, Zauner W, Mellitzer G, et al. The glucocorticoid receptor cooperates with the erythropoietinreceptor and c-Kit to enhance and sustain proliferation of erythroid progenitors in vitro. *Blood* 1999; 94: 550-559.
53. Dror Y, Ward AC, Touw IP, Freedman MH. Combined corticosteroid/granulocyte colony-stimulating factor (G-CSF) therapy in the treatment of severe congenital neutropenia unresponsive to G-CSF: Activated glucocorticoid receptors synergize with G-CSF signals. *Exp Hematol.* 2000; 28: 1381-1389.
54. Armitage S, Hargreaves R, SamsonD, et al. CD34 counts to predict the adequate collection of peripheral blood progenitor cells. *Bone Marrow Transplant* 1997; 20: 587-591.
55. Möhle R, Murea S, Pförsich M, et al. Estimation of the progenitor cell yield in a leukapheresis product by previous measurement of CD34+ cells in the peripheral blood. *Vox Sang* 1996; 71: 90-96.
56. Rill DR, Moen RC, Buschle M, et al. An approach for the analysis of relapse and marrow reconstitution after autologous marrow transplantation using retrovirus-mediated gene transfer. *Blood* 1992; 79: 2694-2700.
57. Vora AJ, Toh CH, Peel J, Greaves M. Use of granulocyte colony-stimulating factor (G-CSF) for mobilizing peripheral blood stem cells:risk of mobilizing clonal

myeloma cells in patients with bone marrow infiltration. *Br J Haematol* 1994; 86: 180-182.

Figure and Table legends.

Table I. Demographic and disease status of patients at diagnosis.

Table II. Parameters of leukapheresis (LK) procedures

Table III. Study A and B: Collection of CD34+ PBSC ($\times 10^6/\text{kg}$) for harvested patients (n) in each group.

Table IV: Differences between MM and Lymphoma

Table V. Mobilization failure in patients receiving the hematopoietic growth factor

Figure 1. Design of the study: mobilization and collection of CD34 cells.

Figure 2. Correlation between the estimated number of CD34 cells ($\%CD34 \times WBC \times 8$)/kg body weight (8 being a coefficient corresponding approximately to 2 blood volumes in L) and the number of collected CD34 cells.

Figure 3. Percentage of patients reaching 5 or 10×10^6 CD34 cells/kg in study B.

Figure 4. Percentage of patients with multiple myeloma and reaching 10×10^6 CD34 cells/kg.

Figure 5. Kinetic of circulating CD34 cells in patients with less (A) and more (B) than one year of prior chemotherapy.

Table I.

| HGF | Study A | | Study B | |
|-----------------------|-------------|--------------|-----------|--------------|
| | G-CSF | G+GM-CSF | G-CSF | G+GM-CSF |
| <i>dose (µg/kg/d)</i> | 5 | 5 + 5 | 5 | 2.5 + 2.5 |
| N | 29 | 31 | 30 | 30 |
| Age (Yr) | | | | |
| Median | 54 | 51* | 53 | 57* |
| (<i>range</i>) | (22-65) | (32-70) | (23-69) | (35-74) |
| Weight (kg) | | | | |
| Median | 62.5 | 68.5* | 65 | 67.5* |
| (<i>range</i>) | (44-114) | (47-110) | (43-104) | (40-100) |
| Sex | | | | |
| Male | 16 (55 %) | 17 (55 %)* | 16 (53 %) | 14 (47 %)* |
| Female | 13 (45 %) | 14 (47 %)* | 14 (47 %) | 15 (53 %)* |
| MM | 14 | 12 | 17 | 15 |
| Stage II | 3 | - | 3 | 3 |
| III | 11 | 12 | 14 | 12 |
| NHL | 5 | 7 | 8 | 12 |
| Fol. SR | 2 | 1 | 1 | 3 |
| Dif. | 1 | 4 | 6 | 9 |
| MZ | 1 | 2 | 1 | - |
| Cut. | 1 | - | - | - |
| HD | 4 | 2 | 2 | - |
| SR | 3 | 1 | 1 | - |
| Ref. | 1 | 1 | 1 | - |
| CLL (St. B) | 1 | 1 | 1 | 1 |
| Solid Tumors | 5 | 9 | 2 | 2 |
| Breast HR | 1 | 5 | - | 1 |
| M1 | 1 | 3 | 1 | 1 |
| Testis M1 | 2 | 1 | 1 | - |
| Ovarian M1 | 1 | - | - | - |

MM= Multiple Myeloma, stage according to Salmon-Durie classification. NHL= Non-Hodgkin's Lymphoma, Fol.= follicular, Dif.= diffuse, MZ= mantle zone, Cut.= cutaneous, SR= sensitive relapse, Ref.= refractory. HDK = Hodgkin disease, (all NHL and HDK are stage IV according to Ann Harbor classification). CLL= Chronic Lymphocytic Leukemia, stage according to Binet Classification. Breast, HR= high risk according to Scarff-Bloom/Richardson score ≥ 6 and/or more than 6 loco-regional metastatic nodes, M1 presence of long-distance metastase according to UICC-TNM classification. Ovarian= Ovarian cancer.

* not significantly different from data of the G-CSF group using a Wilcoxon non parametric test.

Table II.

| | Study A | | Study B | |
|----------------------------------|--------------------------|---------------------------|--------------------------|--------------------------|
| HGF | G-CSF | G+GM-CSF | G-CSF | G+GM-CSF |
| <i>dose (µg/kg/d)</i> | 5 | 5 + 5 | 5 | 2.5 + 2.5 |
| Median values and (range) | | | | |
| Leukapheresis N=232 | 61 | 62* | 56 | 53* |
| LK/patient # | 2 (0-3) | 2* (0-3) | 2 (0-3) | 2* (0-3) |
| Day of the 1st LK | 11 (9-13) | 10* (8-17) | 11 (8-16) | 10* (7-12) |
| WBC at 1st LK | 6 (0.7-39) | 8.8* (0.8-46.2) | 7.2 (0.9-51.2) | 4* (1-25.9) |
| BV/LK ° | 3.01 (1.5-4.3) | 2.55* (1.8-4) | 2.9 (1.8-5.1) | 2.8* (1.7-3.5) |

LK/patient[#] = Leukapheresis undergone by patient. WBC= White Blood Count x 10⁹/L. BV/LK ° = number of Blood Volume processed during a leukapheresis.

* not significantly different from data of the G-CSF group using a Wilcoxon non parametric test.

Table III.

| HGF | Study A | | | Study B | | |
|----------------------------------|--------------|--------------|----------|---------------|---------------|----------|
| | G-CSF | G+GM-CSF | | G-CSF | G+GM-CSF | |
| <i>dose (µg/kg/d)</i> | 5 | 5 + 5 | | 5 | 2.5 + 2.5 | |
| Median values and (range) | | | | | | |
| Global results | | | <i>P</i> | | | <i>P</i> |
| (n) | 27 | 29 | | 27 | 26 | |
| LK ₁ | 3.1 (1-45) | 7.3 (1-31) | .03 | 3.7 (0.6-30) | 7.07 (1-24.5) | .04 |
| LK ₂ | 3.7 (1-13) | 6.3 (1-59) | .04 | 3 (0.6-14) | 8.02 (1-30) | .02 |
| LK ₃ | 2.7 (2-8) | 5.7 (1-12) | .04 | 3.32 (2.9-7) | 6.71 (1-9) | .2 |
| Total CD34/kg | 9 (3-45) | 18.3 (3-90) | .09 | 8.1 (2-34) | 15.85 (2-35) | .09 |
| CD34 LK ₁ /BV | 1.1 (0.3-12) | 2.4 (0.3-13) | .02 | 1.48 (0.1-12) | 2.41 (0.3-10) | .04 |
| MM | | | | | | |
| (n) | 14 | 11 | | 15 | 14 | |
| LK ₁ | 3.2 (1-18) | 9.3 (3-31) | .03 | 3.7 (1-30) | 8.8 (1-24) | .01 |
| LK ₂ | 5.7 (1.2-13) | 10.4 (4-59) | .04 | 5.6 (1.5-14) | 10.5 (1-30) | .02 |
| LK ₃ | 2.9 (2.3-8) | 4.5 (3-12) | .10 | 3.4 (3-7) | 7.3 (1-9) | .10 |
| Total CD34/kg | 11 (3-29) | 23.8 (11-90) | .005 | 9.26 (3-34) | 24.6 (2-35) | .03 |
| CD34 LK ₁ /BV | 1.17 (0.3-8) | 2.81 (1-13) | .02 | 1.35 (0.1-12) | 2.95 (0.4-10) | .01 |
| NHL/HD/CLL | | | | | | |
| (n) | 9 | 9 | | 10 | 10 | |
| LK ₁ | 2.6 (1-35) | 4.2 (1-22) | NS | 3.36 (1-14) | 3.61 (1-10) | NS |
| LK ₂ | 2.6 (1-4) | 3.8 (2-5) | NS | 2.37 (1-3) | 2.4 (1-4) | NS |
| LK ₃ | 2.6 (2-3.5) | 0.8 | / | / | 1.8 | / |
| Total CD34/kg | 6.4 (3-35) | 5.4 (3-26) | NS | 5.4 (2-14) | 5.2 (3-10) | NS |
| CD34 LK ₁ /BV | 0.9 (0.4-12) | 1.9 (0.3-7) | NS | 1.6 (0.3-6.5) | 1.4 (0.3-5) | NS |

CD34+ PBSC are harvested during one or two or three maximum large volume leukapheresis (LK). LK₁ = 1 st Leukapheresis, LK₂ = 2 nd Leukapheresis, LK₃ = 3 rd Leukapheresis. CD34 LK1/BV= CD34+ cells harvest in the 1 st LK in function of number of blood volume. P values were determined with a Wilcoxon test.

Datas of Solids Tumors are not detailed.

Table IV.

| | Study A | | Study B | |
|--|---------------------------------------|---------------------------------------|---------------------------------------|--------------------------------------|
| HGF | G-CSF | G+GM-CSF | G-CSF | G+GM-CSF |
| <i>dose (µg/kg/d)</i> | 5 | 5 + 5 | 5 | 2.5 + 2.5 |
| Median values and (range) | | | | |
| Number of administred courses | | | | |
| MM | | 3 (2-6) | | 3 (3-4) |
| NHL/HD/CLL | | 5 (2-16) <i>P = .02</i> | | 6 (3-20) <i>P = .005</i> |
| CD34 LK₁ /BV | | | | |
| MM | 1.2 (0.3-8) | 2.80 (1-13) | 1.35 (0.1-12) | 2.95 (0.4-10) |
| NHL/HD/CLL | 0.9 (0.4-12) <i>P = .12</i> | 1.9 (0.3-7) <i>P = .047</i> | 1.6 (0.3-6.5) <i>P = .8</i> | 1.4 (0.3-5) <i>P = .01</i> |

Table V.

| | Study A | | | Study B | | |
|---------------------------------|-------------------------|-----------------------|-----------|-------------------------|------------------------|-----------------------|
| Number of failure (%) | 4/56 (7.14 %) | | | 3/53 (5.66 %) | | <i>P</i> <i>NS</i> |
| HGF | G-CSF | G+GM-CSF | | G-CSF | G+GM-CSF | |
| <i>dose (µg/kg/d)</i> | 5 | 5 + 5 | | 5 | 2.5 + 2.5 | |
| | 2/27 (7.4 %) | 2/29 (6.9%) | <i>NS</i> | 2/27 (7.4 %) | 1/26 (3.8 %) | <i>NS</i> |

Number and percentage of patients who received the HGF and impaired the mobilization
 p = Fisher's exact test

Figure 1.




Figure 2.





Figure 3.




Figure 4.





Figure 5.



A

B

