

Estrogen receptor beta inhibits proliferation and invasion of breast cancer cells

**Gwendal Lazennec^{*}, Damien Bresson^{*}, Annick Lucas, Corine Chauveau
and Françoise Vignon[¶]**

Inserm U540 "Molecular and Cellular Endocrinology of Cancers",
60, rue de Navacelles - 34090 Montpellier, France

Running title: ER β and breast cancer

^{*} Both authors have equally contributed to this work.

[¶] **Corresponding Author:**

Dr Françoise Vignon

INSERM U540 " Molecular and Cellular Endocrinology of Cancers ",

60, rue de Navacelles - 34090 Montpellier, France

Tel: (33) 4 67 04 37 63; Fax: (33) 4 67 54 05 98

E-mail: vignon@u540.montp.inserm.fr

ABSTRACT

Recent studies indicate that the expression of ER β in breast cancer is lower than in normal breast, suggesting that ER β could play an important role in carcinogenesis. To investigate this hypothesis, we engineered estrogen-receptor negative MDA-MB-231 breast cancer cells to reintroduce either ER α or ER β protein with an adenoviral vector. In these cells, ER β (as ER α) expression was monitored using RT-PCR and Western blot. ER β protein was localized in the nucleus (immunocytochemistry) and able to transactivate estrogen-responsive reporter constructs in the presence of estradiol. ER β and ER α induced the expression of several endogenous genes such as pS2, TGF α or the cyclin kinase inhibitor p21, but in contrast to ER α , ER β was unable to regulate c-myc proto-oncogene expression. The pure antiestrogen ICI 164,384 completely blocked ER α and ER β estrogen-induced activities. ER β inhibited MDA-MB-231 cell proliferation in a ligand-independent manner, whereas ER α inhibition of proliferation is hormone-dependent. Moreover, ER β and ER α , decreased cell motility and invasion. Our data bring the first evidence that ER β is an important modulator of proliferation and invasion of breast cancer cells and support the hypothesis that the loss of ER β expression could be one of the events leading to the development of breast cancer.

INTRODUCTION

Estrogens modulate sexual gland development and reproductive functions but have also beneficial effects on cardiovascular, nervous systems or bone integrity (1). Besides, estrogens are potent mitogens in the mammary gland, where they regulate the growth, development and functioning of normal as well as cancerous breast (2, 3). Epidemiological evidences and numerous animal studies have indicated that estrogens play a role in the proliferation and progression of breast cancer; the removal of the ovaries or the treatment with antiestrogens oppose their deleterious effects (4, 5). Though there are growing evidences that estrogens can operate through non genomic pathways (6-8), estrogen receptors, ER α (NR3A1) and ER β (NR3A2), which belong to a large family of nuclear receptors, are mediating the genomic action of estrogens by acting as ligand dependent transcription factors (9, 10). Most human breast cancers, at least initially, express ER α and the presence of ER α is generally considered as an indication of hormone dependence, eventhough only 60% of ER-positive tumors will respond to adjuvant therapy with tamoxifen (11).

Although ER α has been cloned more than 10 years ago (12), the presence of ER β has been unrecognized till recently (13, 14). ER α and ER β have diverged early during evolution (15) and differ mostly in the N-terminal A/B domain, and to a lesser extent in the ligand binding domain (E domain). These differences suggest that the two receptors could serve distinct actions. Indeed, the activation functions AF-1 and AF-2 located respectively in the A/B and LBD domains display activities that are promoter and cell-specific (16-18). Cowley et al. (17) have shown that the AF-1 activity of ER β is weak compared with that of ER α on estrogen-responsive reporters, whereas their AF-2 activities are similar. In turn, when both AF-1 and AF-2 functions are active in a particular cell and/or on a particular promoter, the activity of ER α greatly exceeds that of ER β , whereas ER α and ER β activities are similar when only

AF-2 is required. The weaker activity of ER β in many promoter and cell contexts has also been reported by several groups (18-20).

Moreover, ER α and ER β knock-out mice have been generated and demonstrated striking different patterns (21, 22). ER β knock-out mice show significantly reduced fertility in female, with ovaries exhibiting follicular arrest and anovulation. However, these mutant mice have a normal mammary gland development and lactate (21). On the contrary, ER α knock-out mice have an impaired fertility for both sexes and exhibit an estrogen-insensitive mammary gland and genital tract (22), suggesting possible overlapping and distinct action on the expression of genes regulating the important biological functions. Concerning the rodent mammary gland, both estrogen receptors are expressed in the rat mammary gland but the presence and cellular distribution of the two receptors are distinct (23). In prepubertal rats, ER α is detected in 40% of the epithelial cell nuclei. During puberty and pregnancy, ER α expression is strongly decreased, whereas ER α is present in 70% of epithelial cells during lactation. About 60-70% of epithelial cells express ER β at all stages of breast development. Cells coexpressing both receptors represent up to 60% of the epithelial cells during lactation but are rare during pregnancy. Moreover, more than 90% of ER β expressing cells do not proliferate (23).

In agreement with these observations, recent studies in humans indicate that the ER β expression is decreased between normal and neoplastic breast, as well as colon and ovarian cancer (24-30), suggesting that ER β could be an inhibitor of tumorigenesis. In order to test this hypothesis, we have engineered a receptor negative breast cancer cell line to express functional ER β . In this cell line, ER β was able to activate the transcription of synthetic promoters in transient transfection experiments as well as natural endogenous promoters. Interestingly ER β had major effects both on the proliferation, motility and morphology of the cells, suggesting that ER β could effectively act as an inhibitor of breast cancer development.

MATERIALS AND METHODS

Plasmids.

The reporter plasmid ERE2-TK-CAT contains two copies of the consensus ERE cloned upstream of the minimal herpes simplex virus thymidine kinase promoter. CMV-hER α and CMV-hER β correspond to the wild-type ER α and ER β cDNAs cloned into CMV5. CMV-GAL reporter was used as an internal control and corresponds to the β -galactosidase gene under the control of the CMV promoter.

Recombinant adenovirus construction and propagation.

The complete coding sequence of wild-type hER β or hER α cDNAs were subcloned in BamHI site of the pACsk12CMV5 shuttle vector. To obtain recombinant viruses, permissive HEK-293 cells (human embryonic kidney cells) were cotransfected with the recombinant pACsk12CMV5-hER plasmid and with pJM17, which contains the remainder of the adenoviral genome as previously described (31-33). *In vivo* recombination of the plasmids generates infectious viral particles (Ad-hER α or Ad-hER β). DNA from these viruses were screened for the presence of the hER cDNA by PCR with hER primers, and titered virus stocks were used to infect MDA-MB-231 cells.

Cell Culture and Transient Transfection.

HEK-293 cells were cultured in DMEM-F12 supplemented with 10% FCS (fetal calf serum) in the presence of 5% CO₂. MDA-MB-231 cells (human breast cancer cells) were cultured in Leibovitz L-15 medium containing 10% FCS. $3 \cdot 10^5$ cells were plated in 6-well plates in phenol red -free DMEM-F12 supplemented with 10% CDFCS (charcoal-dextran treated FCS) 24h before transfection. Transfections were performed by lipofection (lipofectamine, Life Technologies, Rockville, MA) using 4 μ g of CAT reporter construct, 1 μ g of the internal reference β -galactosidase reporter plasmid (CMV-GAL) and CMV-hER expression vectors or

recombinant viruses per well. Transactivation ability was determined by CAT activity on the whole cell extract as previously described (34).

Whole cell extract preparation and western blot.

MDA-MB-231 cells were lysed in TEG (10 mM Tris-HCl, pH 7.4, 1.5 mM EDTA, and 10% glycerol) containing protease inhibitors (5 µg/ml aprotinin, leupeptin and pepstatin A and 0.1 mM phenylmethylsulfonyl fluoride). Then, cells were sonicated and the cellular debris were pelleted by centrifugation at 13000g for 20 minutes in microfuge tubes. 45 µg of whole cell extract proteins were subjected to SDS-PAGE followed by electrotransfer onto a nitrocellulose membrane. The blot was probed with anti hER α (SRA-1000) or hER β antibody (1:1000) (CWK-F12, (35)) and then incubated with goat anti-mouse IgG horseradish peroxidase conjugated antibody (1 µg/ml). ECL kit from Amersham (Arlington, IL, USA) was used for detection.

Gel Mobility Shift Assays.

Briefly, 30000 cpm of the [³²P]-labeled (AGCTCTTTGATCAGGTCACCTGTGACCTGACTTT) ERE double strand oligonucleotide was combined with 1 µg poly (dI-dC) and 5 µg of MDA-MB-231 whole cell extract (WCE). When indicated, anti-hER α (Stressgen, SRA-1000) or anti-hER β antibodies (CWK-F12, a kind gift of Pr B.S. Katzenellenbogen, for more details, see (35)) were added. The reaction buffer contained 20 mM Hepes, pH 7.9; 1 mM DTT; 50 mM KCl; 10% glycerol; 2.5 mM MgCl₂. Protein-DNA complexes were separated from the free probe by non-denaturing gel electrophoresis with 4% polyacrylamide gels in 0.5 X TBE.

Detection of ER α and ER β protein by immunocytochemistry.

MDA-MB-231 cells were seeded in 10% CDFCS DMEM-F12 on sterile coverslips in six-well plates and infected with the different adenoviruses. 48h after infection, the cells were fixed (formaldehyde 3.7% 12 min/methanol 5 min/ acetone 2min) and washed with PBS. The coverslips were incubated for 30 min with PBS containing non-immune rabbit serum (1:40).

Then the cells were incubated with the primary antibody (ER α : SRA-1000 1:2000 (Stressgen); ER β : CWK-F12 1:3000) in PBS for 60 min at room temperature. The cells were then incubated with the secondary antibody (rabbit anti-mouse peroxidase conjugate, Sigma, 13000) in PBS-BGG (bovine gamma globulin) for 30 min at room temperature. Finally, the cells were incubated with a DAB (diaminobenzidine) chromogen solution (0.66mg/ml in PBS+0.08% H₂O₂ (30 vol)) for 10 min at room temperature. The cells were counterstained with hematoxylin.

RNA Isolation, Northern Blot and RT-PCR.

Total RNA was isolated from MDA-MB-231 cells using the TRIzol reagent from Life Technologies (Rockville, MA) as described by the manufacturer. Probes were amplified by RT-PCR using specific primers:

ER α : AAAAGACCGAAGAGGAGGGAGAAT/ATCCGGAACCGAGATGATGTAG,

ER β : GCCGCCCATGTGCTGAT/GGACCCCGTGATGGAGGACTT,

c-myc: TACCCTCTCAACGACAGCAGCTCGCCCAAC/TCTTGACATTCTCCTCGGTGTCCGAGGACC,

p21: CGAGTGGGGGCATCATCAAAAAC/TGTTACAGGAGCTGGAAGGTGTTTG,

pS2: TGACTCGGGGTCGCCTTTGGAG/GTGAGCCGAGGCACAGCTGCAG,

TGF α : CCTGTTCGCTCTGGGTATTGTGTTG/CGTGGTCCGCTGATTTCTTCTCTAG).

Reverse transcription was performed using random primers and GenAmp (Roche, Basel, Switzerland) RT-PCR kit. The amplifying primers are described above. The PCR was performed with Platinum Taq polymerase (Life Technologies) and 1:40 of reverse transcription reaction. Cycles of 30s at 94°C, 1 min at 60°C, 1.30 min at 72°C were done 29 times. A tenth of each PCR was electrophoresed on agarose gel. For Northern blot analysis, 20 μ g of total RNA were electrophoresed and then hybridized with the different probes.

Cell proliferation studies.

Cells were maintained for 24h in 10% CDFCS DMEM-F12 and then seeded at 30000 cells/well in 24-well dishes in 10% CDFCS DMEM-F12. Cells were infected overnight with the different viruses. The next morning, the medium was removed and replaced with fresh 10% CDFCS DMEM-F12 medium. Treatment with E2 or ICI 164,384 began at the same time. After 2, 4 or 6 days of E2, ICI 164,384 or both compounds treatments, the cells were trypsinized and counted using a Coultronics Coulter counter.

Wound healing assay.

Cells were plated in 6-well dishes in DMEM-F12 containing 5% CDFCS. 24h after plating, the cells were infected with the different viruses overnight. The next morning, ethanol or E2 treatment started. After 20h of treatment, wound induced migration was triggered by scraping the cells at day 1 with a blue tip and the wound was pictured immediately. 18h after the wound (day 2), the cells were pictured again. The % of wound filling was calculated by measuring on the pictures the remaining gap space. The ratio of the gap space at day 2 over the gap space at day 1 gives the percentage of wound filling.

Invasion assay.

MDA-MB-231 cells infected with Ad5, Ad-hER α or Ad-hER β (MOI 100) were plated 24h after infection in the upper compartment of a 24-well Transwell (Costar) on a polycarbonate filter (8 μ m pore size) which was first coated with 30 μ g of matrigel (Becton Dickinson). The lower compartment of the well was filled with DMEM-F12 supplemented with 10% CDFCS and 30 μ g/ml fibronectin (Sigma). As a control, the same cells were layered on 24-well plates. Cells were treated with ethanol vehicle or E2 (10⁻⁸ M). After 36h of migration, cells which have migrated to the lower side of the filter and cells present in the control plates were trypsinized and counted using a Coulter counter.

Morphology analysis

MDA-MB-231 cells were cultured in DMEM-F12 supplemented with 10% CDFCS. After infection with Ad5, Ad-hER α or Ad-hER β (MOI 100), cells were treated with control vehicle ethanol or E2 (10^{-8} M) for 48h and then pictured using a Zeiss phase contrast microscope.

RESULTS

Adenoviruses elicit a high infection of MDA-MB-231 cells

Replication-deficient adenoviruses encoding hER α or hER β cDNA sequences were constructed and used to infect ER α -negative MDA-MB-231 cells (Fig. 1A). To control the efficiency of infection of MDA-MB-231 cells, we first treated the cells with an adenovirus coding for the β -galactosidase protein (Fig. 1B). We observed a very efficient infection of the cells when increasing the MOI (multiplicity of infection) from 1 to 100, leading to an infection of about 80% of the cells at the highest MOI.

Adenoviral mediated expression of ER α and ER β

ER α and ER β expression was examined following infection of MDA-MB-231 cells (Fig. 2A, 2B and 2D). We could not detect any expression of ER α or ER β in MDA-MB-231 cells infected with the non recombinant virus Ad5 as well as in non infected cells (data not shown). On the contrary, after infection with Ad-hER β virus, a high expression of ER β could be seen at RNA (Fig. 2A) and protein (Fig. 2B) levels. Similarly, expression of an equivalent amount of ER α was detected in Ad-hER α infected cells (Fig. 2A and 2B). To further check the functionality of the expressed ER β protein, we analyzed its ability to bind to an ERE (estrogen responsive element) DNA sequence by performing gel shift experiments (Fig. 2C). A specific binding could only be seen when ER α or ER β extracts were used (lanes 3 and 5). Moreover, the ER β shifted complex had a faster migration rate than the ER α one. The specificity of the shifted complex could be further demonstrated by using ER α and ER β specific antibodies (lanes 4 and 6). We then determined the cellular localization of ER α and ER β expressed proteins by immunocytochemistry (Figure 2D). ER α - and ER β -infected cells

displayed a clear and exclusive nuclear staining when using ER α and ER β antibodies respectively. These data confirm our previous findings with ER α (31) and suggest that ER β protein is correctly expressed at RNA and protein levels in MDA-MB-231 cells, is addressed to the nucleus and is able to bind to DNA.

ER β is able to activate estrogen-sensitive reporter genes

To further assess the functionality of the receptors produced, we analysed their ability to transactivate estrogen-sensitive reporter genes (Fig. 3). As a control, we transfected regular plasmids encoding hER α or hER β along with the ERE2-TK-CAT reporter (Fig. 3A). We observed a strong activation of the reporter by ER α in the presence of E2. ER β was also able to activate the transcription in the presence of E2, but the stimulation was half that obtained with ER α (Fig. 3A). We then tested the ability of our recombinant Ad-hER β virus to activate the ERE2-TK-CAT reporter (Fig. 3B). When increasing MOI of Ad-hER β were used, a strong ligand-dependent activation of the reporter occurred, demonstrating that the adenovirally produced ER β exhibits a classical pattern of activation. However, at low MOI, ER β was less active than ER α , whereas the use of higher MOI of Ad-hER β virus elicited a good activation of the reporter. We then checked the sensitivity of ER β to estrogen stimulation (Fig. 3C). We observed a characteristic dose-response curve for ER β , similar to that obtained for ER α . A slight shift in the sensitivity to E2 was observed for ER β , which reached its maximal activity at 10^{-8} M, whereas ER α activity was maximal at 10^{-9} M. Similar results have been obtained by others showing that ER β has a weaker activity than ER α at low concentrations of E2 (36). In order to demonstrate that the expressed receptors were triggering estrogen effect, we analyzed their transactivation ability in the presence of the pure antiestrogen ICI 164, 384 (Fig. 3D). As expected, ICI 164, 384 could not stimulate ER α or

ER β activity but completely shut down both the basal and E2-induced activities of both receptors, suggesting that the basal activity of both receptors was most probably due to remaining traces of E2 in the stripped serum.

ER α and ER β have common but also distinct target genes

Very little is known about the specific target genes of ER β . Therefore, we examined in these ER β -positive cells the expression of 4 genes, TGF α (Transforming growth factor alpha), p21, c-myc proto-oncogene and pS2 genes, which are also regulated in ER α -infected cells in the presence of E2 (Fig. 4A, B). ER α and ER β were able to activate the expression of pS2, p21 and TGF α genes in an estrogen-dependent manner. ER β was 2 to 3 fold less potent than ER α to stimulate pS2, p21 and TGF α expression than ER α . pS2 activation was maximum at 48h of E2 treatment for both receptors. For TGF α and p21 genes the maximal activation was reached at 24h for both receptors, suggesting that these genes exhibit an earlier response than pS2. Very interestingly, ER α almost completely abolished the expression of c-myc in the presence of E2, whereas ER β had no significant effect. These data suggest that ER α and ER β effects on target genes differ both in the amplitude of regulation and in the nature of the genes regulated. To evaluate whether antiestrogens could also modulate the expression of these genes, we performed the same experiments in the presence of ICI 164,384, alone or in combination with E2 (Fig. 4C). ICI 164,384 was able to decrease the basal level of expression of pS2, p21 and TGF α and most interestingly completely reverse the induction of pS2, p21 and TGF α genes by E2 in ER α and ER β infected cells.

ER α and ER β are potent inhibitors of the proliferation

The main question was to determine if ER β expression could modulate the proliferation rate of MDA-MB-231 cells. Control cells (non-infected or Ad5 infected) had a similar growth pattern in the absence or in the presence of estrogens (Fig. 5A, left panel). When MDA-MB-231 cells were infected with Ad-hER α virus (Fig. 5A, middle panel), they proliferated at the same rate as naive cells in the absence of estrogens. But when E2 was added, a strong inhibition (50%) of the proliferation occurred, which is in agreement with our previous work (31). Very interestingly, ER β was also able to inhibit the proliferation of MDA-MB-231 cells, but this effect was totally ligand-independent: a 40% inhibition occurred whether or not estrogens were present (Fig. 5A, right panel). This is to our knowledge, the first direct evidence that ER β can be involved in the proliferation control of breast cancer cells. To determine the effects of pure antiestrogens on cell proliferation, we performed experiments in the presence of ICI 164,384 (Fig. 5B). ICI 164,384 had no effect by itself on the proliferation of naive, ER α or ER β expressing cells. However, ICI 164,384 completely reversed E2-triggered inhibition in Ad-hER α infected cells. Moreover, ICI 164,384 or in combination with E2 could not modulate ER β expressing cells proliferation rate. These data confirm that inhibition of the proliferation by ER α is ligand-dependent, whereas the inhibition by ER β was ligand-independent.

ER expression has profound effects on invasion, motility and morphology of the cells

It was of interest to determine whether ER β expression might affect cell motility and therefore modulate the invasiveness of the cells. To address this issue, we performed wound healing-induced migration experiments (Figure 6A, B). Infected cells were forced to migrate through the space created by scraping the monolayer with a tip. After 18h of migration, Ad5-

infected MDA-MB-231 cells had filled 85% of the wound (Figure 6B). On the contrary, ER α -infected cells had only partially (50%) filled the wound. This lack of migration was not significantly affected when E2 was added. ER β was also able to inhibit the cell motility. This inhibition occurred in the absence or in the presence of E2 (filling of only 30-40% of the space), suggesting that ER β was a more potent inhibitor of motility than ER α . We then evaluated the migration ability of these cells using the classic Transwell *in vitro* assay. In this assay, cells are encouraged to migrate from the upper compartment coated with matrigel to the lower compartment coated with fibronectin, which serves as a chemoattractant. After 36h of migration, we observed that Ad-hER α and Ad-hER β migrating cells represent respectively 70% and 50% of the control migrating cells (Figure 6C). Addition of E2 did not change the migration rate of any kind of cells. In conclusion, motility and invasion assays are in close agreement suggesting that ER β is a more potent inhibitor of cell migration than ER α .

In correlation with these observations on the reduction of cell motility and invasion following ER expression, we have tested whether cell morphology was affected. Strikingly, ER β led to a change in the morphology of the cells (Fig. 7). Infected cells lose their fibroblastic appearance and acquired an "epithelioid-like" shape. The cells were enlarged and more rounded. ER α expression also modified the morphology of the cells and led to a more flattened shape of the cells. This change was even more pronounced in the presence of estradiol creating a characteristic structure of "branching" cells.

DISCUSSION

The use of recombinant adenoviruses has enabled us to express ER β in breast cancer cells devoid of detectable endogenous ERs. ER β protein appears to be fully functional as shown by DNA binding, cellular localization, transient transfection experiments and regulation of estrogen-regulated endogenous genes. Thus, our data suggest that this novel model exhibits all the interesting features required for the study of ER β action in breast cancer cells and could thus be predictive of its role in human tumors.

In MDA-MB-231 cells, expressed ER β regulated the activity of reporter constructs and endogenous genes. The weaker activity of ER β on reporter genes compared to ER α is likely due to a lack of ER β AF-1 activity as suggested by several studies (17, 19, 20, 36). Indeed, depending on the cellular and promoter context, AF-1 function has a negligible or high activity, which in turn leads to a greatly enhanced activity of ER α when gene regulation requires both AF-1 and AF-2. On the contrary, when only AF-2 is active, both receptors exhibit similar activities.

Interestingly, we show here that ER β can induce the expression of pS2, p21 and TGF α , whereas it has no effect c-myc expression. We and others have previously shown that in cellular models in which ER α was exogenously expressed, ER α could induce cathepsin D, pS2, p21 and TGF α expression in the presence of E2 (31, 37-39), whereas it was able to down regulate c-myc, TGF β 2, BRCA-1, BRCA-2 and c-fos/c-jun expression (31, 37, 40). This suggests that ER α and ER β target genes are partially overlapping but that there are also target genes only regulated by one type of receptor. To date, only a limited number of promoters regulated specifically by one E2 liganded-ER isotype have been identified. This is the case of the osteopontin (41) and hTERT (catalytic subunit of human telomerase) (42) promoters which are up-regulated by ER α and not by ER β . There is only one demonstration

of a gene exclusively regulated by ER β and not by ER α in the presence of estrogens. This is the case of methallothionein II gene which is specifically up-regulated by ER β in SAOS-2 cells but is regulated by ER α and ER β in LNCaPLN3 cells (43). Interestingly, c-myc RNA levels (whose expression is generally correlated with the proliferation rate) were not affected by ER β in the presence of E2 in contrast to what is observed in ER α -MDA-MB-231 cells. p21 RNA levels were increased by ER β in the presence of E2. p21 expression is definitely induced in numerous growth arrested cells (44), even if there are no growth abnormalities in p21-null mice (45). Moreover, p21 is also involved in some cases in the differentiation process, without affecting proliferation (46). Therefore, p21 up-regulation observed in E2-treated ER β expressing cells might be related to differentiation, as suggested by the morphological changes observed. The change in morphology from a "fibroblastic" to an "epithelioid-like" shape of MDA-MB-231 cells infected with ER β has been reported for other engineered cell lines, such as MDA-MB-231 cells stably expressing PR (47). Interestingly, the proliferation of these cells was inhibited by the addition of progesterone, the corresponding receptor ligand. It will be of interest to evaluate whether ER β expression leads to changes in adhesion properties of the cells and in particular to determine whether adhesion molecule expression is altered.

Our work represents the first direct evidence that ER β is involved in the control of the proliferation of breast cancer cells. Surprisingly, ER β inhibition of the proliferation was ligand-independent, whereas ER β was able to regulate reporter genes and endogenous gene expression in a ligand-dependent manner. Exogenous ER α expression using stable or retroviral infected cell lines has already been reported (31, 48-51). All these studies have shown an E2-dependent decreased proliferation of ER α expressing cells, ranging from a modest to a high level of inhibition. Therefore, our data suggest that ER α and ER β inhibit the

proliferation through distinct mechanisms. To date, only one study has reported the stable expression of ER β (52). These authors used rat-1 cells and compared ER α and ER β transfectants. ER β did not affect the proliferation, but in this model, in disagreement with all other studies, ER α had no ability either to repress proliferation in the presence of estradiol. More interestingly, in contrast to ER α , the effect of exogenous expression of ER β on proliferation seems to be relevant to the clinical situation. Indeed, numerous studies have shown that the ER β /ER α ratio was decreased between normal to cancerous tissues, as in breast, colon and ovarian cancers (24-30), suggesting that ER β could play a negative role on tumorigenesis. Roger et al. (27) have shown that ER β protein was expressed in 85% of epithelial cells of normal mammary gland and this expression was not significantly altered in non-proliferative breast benign disease (BBD). On the contrary, ER β expression was decreased in proliferative BBD and was nearly completely shut down in high grade *carcinoma in situ* (DCIS), suggesting that the presence of ER β is associated with non proliferative states of the disease. What is still under question is whether ER β expression in breast cancers could be considered as a good prognostic indicator. In invasive breast cancer, other studies have shown that ER β protein expression was associated with less invasive and proliferative tumors (negative axillary node status, low grade, low S-phase fraction) suggesting that ER β might be a good prognostic indicator (53). This conclusion was also supported by Omoto et al. (54), even though they could not see a significant correlation between ER β expression and other known clinical parameters. Finally, in terms of adjuvant hormonal therapy (AHT), the conclusions are rather contradictory at present as some studies suggest that ER β expressing tumors are associated to a better survival of patients under AHT (55) whereas other results suggest that ER β is up-regulated in tamoxifen resistant tumors and could be involved in tamoxifen resistance (56, 57).

In agreement with previous work (58), our data show that ER α and ER β activities on an ERE-containing reporter and on estrogen regulated genes can be inhibited by the pure antiestrogen ICI 164,384. Moreover, several studies have underlined the differences between ER α and ER β in terms of response to estrogens or anti-estrogens on AP-1 sites. Indeed, ER β is able to potentiate AP-1 containing reporters in the presence of antiestrogens but not in the presence of estrogens. ER α stimulates AP-1 activity in the presence of estrogens and anti-estrogens in endometrial cells (58-60), but antiestrogens have no effect on AP-1 activity in breast cancer cells (60, 61). Of particular note, ER β is overall more potent than ER α on AP-1 sites, whereas the contrary occurs on EREs (17, 19, 20, 36, 58).

We also show that ER α and ER β inhibit migration and invasion in a ligand-independent manner. These effects of ER β are in close agreement with a previous report showing that ER α inhibits the migration of ER α -negative breast cancer cells (48, 62). In the context of breast cancer, such a reduction of invasion and motility would certainly lead to less aggressive cancers with a lower rate of metastasis. These results fit also well with numerous reports describing that ER-positive breast cancer cells are generally less invasive than ER-negative breast cancer cells (63-66) and that ER β expressing tumors are less metastatic (53). Moreover, reintroduction of ER α in ER-negative breast cancer cells decreases their invasion and metastatic potential (48). Thus, both ER α and ER β are able to reverse the invasive phenotype of MDA-MB-231 cells into less invasive cells, mimicking the situation of ER-positive breast cancer cells.

In conclusion, our results strongly support the idea that ER β could be a potent proliferation gatekeeper as well as an inhibitor of cell motility and invasion. The decreased expression of ER β observed between normal and cancerous breast could be one of the events leading to an uncontrolled proliferation of the cells. Our data suggest that the use of ER β itself or of some

of its target genes could be of interest to design a gene therapy approach against hormone-unresponsive breast cancer.

ACKNOWLEDGMENTS.

We thank Pr B.S. Katzenellenbogen for the gift of ER β antibody, ER α and ER β cDNAs and ERE2-TK-CAT construct. We are also grateful to Dr P. Moullier and AFM (Association Française contre les Myopathies) for their support to produce the viruses. We acknowledge the participation of A. Licznar and M. Lacroix to some experiments during their graduate courses. We thank Dr P. Roger for critically reviewing this manuscript and J.Y. Cance for the photographic work. This work was supported by grants from ARC (Association pour la Recherche contre le Cancer, grant n°5405), la Ligue Nationale contre le Cancer (Comité du Gard), Inserm and CNRS.

ABBREVIATIONS.

Ad: adenovirus, Ad5: non recombinant adenovirus, Ad hER α or β : recombinant adenovirus with hER α or β , AF: activation function, BGG: bovine gamma globulin, DAB: diaminobenzidine, ERE: estrogen responsive element, hER α or β : human estrogen receptor α or β , WCE: whole cell extract.

LEGENDS TO FIGURES

Fig 1: Schematic representation of adenovirus construction and high infection efficiency of MDA-MB-231 cells

A. Ad-hER α and Ad-hER β viruses were constructed as described in Materials and Methods using *in vivo* recombination in HEK-293 cells. The recombination occurs between the shuttle vector pAC_{SK12}-CMV₅ carrying hER α or hER β cDNAs and pJM17 adenoviral sequences. **B.** MDA-MB 231 cells were grown in 6-well plates and infected overnight with no virus (A), or Ad-GAL virus at MOI (multiplicity of infection) 1 (B), 10 (C), or 25 (D), 50 (E), 100 (F). β -galactosidase activity was monitored after 48 h of expression. The upper panel corresponds to a picture of the entire plate and the lower panel to a 200 fold magnification of each well.

Fig. 2: Adenoviral expression of ER α and ER β in MDA-MB-231 cells

A. MDA-MB-231 cells were infected with the non recombinant (Ad5), Ad-hER α or Ad-hER β viruses at MOI 100. After 1 to 48h of treatment with 10⁻⁸M E2, hER α and hER β expression was monitored by RT-PCR using primers located in the ligand binding domain. The PCR products have a size of 542 bp and 703 bp for ER α and ER β respectively. **B.** hER α and hER β protein expression was analysed by Western blot using hER α (α Ab) or hER β (β Ab) specific antibodies. **C.** WCE from non infected cells (C, lane 1), non recombinant viruses (Ad5, lane 2), Ad-hER α (hER α) (lanes 3-4) or Ad-hER β (hER β) (lanes 5-6) infected MDA-MB 231 cells were used for gel shift assay using a consensus ERE as a probe. Supershifts were performed using specific anti hER α (α Ab) (lane 4) or anti-hER β (β Ab) (lane 6) antibodies. **D.** MDA-MB-231 cells were infected with Ad5, Ad-hER α or Ad-hER β (hER β) at MOI 25 and ER α and ER β expression was visualized by immunocytochemistry using ER α (α Ab) and ER β (β Ab) specific antibodies.

Fig. 3: hER α and hER β can activate the transcription of estrogen-sensitive reporter genes

A. Empty CMV5 vector (CMV), CMV-hER α (hER α) or CMV-hER β (hER β) vectors were cotransfected in MDA-MB-231 cells with ERE2-TK-CAT reporter constructs and CMV-GAL internal reporter plasmid. Cells were grown for 36 h in the presence of control vehicle ethanol (C) or 10^{-8} M E2. Results are expressed as the percentage of CAT activity in non infected cells (NI) and represent the mean \pm SD (n = 5) of CAT activity after normalization for β -galactosidase activity. **B.** Non infected (NI) or Ad5, Ad-hER α , or Ad-hER β infected MDA-MB-231 cells were transfected with ERE2-TK-CAT and CMV-GAL reporter constructs. Increasing MOI of Ad-hER α and Ad-hER β viruses (0.1/1/10/100) were used. Cells were grown for 36 h in the presence of control vehicle ethanol (C) or 10^{-8} M E2. Results are expressed as the percentage of CAT activity in non infected cells (NI) and represent the mean \pm SD (n = 6) of CAT activity after normalization for β -galactosidase activity. **C.** MDA-MB-231 cells were infected with Ad-hER α and Ad-hER β at MOI 100 and treated with increasing concentrations of E2. Results are expressed as the percentage of CAT activity in non infected cells (NI) and represent the mean \pm SD (n = 6) of CAT activity after normalization for β -galactosidase activity. **D.** MDA-MB-231 cells were either transfected with empty CMV5 vector (CMV), CMV-hER α (hER α) or CMV-hER β (hER β) vectors or infected with Ad5, Ad-hER α or Ad-hER β viruses along with ERE2-TK-CAT and CMV-GAL reporter constructs. Cells were grown for 36 h in the presence of control vehicle ethanol (C), 10^{-8} M E2, ICI 164,384 (10^{-6} M) or the combination of E2 and ICI 164, 384 (10^{-8} M and 10^{-6} M respectively). Results are expressed as the percentage of CAT activity in non infected cells (NI) and represent the mean \pm SD (n = 6) of CAT activity after normalization for β -galactosidase activity.

Fig. 4: Modulation of endogenous gene expression by hER α and hER β

A. MDA-MB-231 cells were infected at MOI 100 with the different viruses. 24 h after infection, the E2 treatment began sequentially. All cells were harvested at the same moment following different times of E2 exposure and RNA extracted. 20 μ g of total RNA were used for Northern blot and hybridized with TGF α , p21, c-myc or pS2 probes. Equal loading was checked with an RNA 28S probe. Data of a representative experiment are shown here. **B.** Quantification of Northern experiments after normalization by 28S RNA levels. Results are the mean \pm SD (n = 3) of 3 experiments. **C.** The same experiments were performed in the presence of control vehicle ethanol (C), E2 (10^{-8} M) (E), ICI 164,384 (10^{-6} M) alone (I) or in combination (EI). Data of a representative experiment are shown here and the quantification after normalization with 28S RNA is indicated below. Results are expressed in arbitrary units of scan.

Fig. 5: hER α and hER β are able to repress the proliferation of MDA-MB-231 cells

MDA-MB-231 cells were either non infected (NI) or infected with Ad5, Ad-hER α or Ad-hER β viruses at MOI 100. **A.** The cells were treated with ethanol vehicle or E2 (10^{-8} M) 24 h after the beginning of the infection. Proliferation rate was determined by counting the cells at day 2, 4 and 6. Results represent the mean \pm SD of 4 determinations. **B.** The effect of the pure antiestrogen ICI 164,384 was evaluated by treating the cells either with ICI 164,384 (10^{-6} M) alone or in combination with E2 (10^{-8} M). On day 4, cells were counted and results represent the mean \pm SD of 3 determinations.

Fig. 6: hER β is a strong inhibitor of motility and invasion

A. MDA-MB-231 cells were infected with Ad5, Ad-hER α or Ad-hER β viruses at MOI 100. 24 h after the beginning of the infection, the cells were then treated with ethanol (control) or E2 (10^{-8} M). After 48h of ligand treatment, cells were scratched with a blue tip and pictured

($t=0$). The wound was pictured again 18h after the scratch ($t=18h$). Pictures of a representative assay are shown here. **B.** Results are shown as the % of wound filling after 18h of migration and represent the mean \pm SD of 3 experiments. **C.** MDA-MB-231 cells were infected with Ad5, Ad-hER α or Ad-hER β (MOI 100). Cells were plated on transwell or on control plates and treated with ethanol vehicle or E2 (10^{-8} M) 24h after infection. Cells which have migrated to the lower side of the filter and cells present in the control plates were counted after 36h of migration. The percentage of control migrating cells was set up to 100. Results are expressed as the percentage of control migrating cells and represent the mean \pm SD of four experiments.

Fig. 7: hER α and hER β alter the morphology of MDA-MB-231 cells

MDA-MB-231 cells were infected with Ad5, Ad-hER α Ad-hER β viruses at MOI 100. 24 h after the beginning of the infection, the cells were then treated with ethanol (control) or E2 (10^{-8} M). After 48h of ligand treatment, cells were pictured under a phase contrast microscope.

REFERENCES

1. **Gustafsson JA** 1999 Estrogen receptor beta--a new dimension in estrogen mechanism of action. *J Endocrinol* 163:379-83
2. **Russo J, Russo IH** 1997 Differentiation and breast cancer. *Medicina (B Aires)* 57:81-91
3. **Haslam SZ, Counterman LJ, Nummy KA** 1993 Effects of epidermal growth factor, estrogen, and progestin on DNA synthesis in mammary cells in vivo are determined by the developmental state of the gland. *J Cell Physiol* 155:72-8
4. **Hilakivi-Clarke L, Clarke R, Lippman M** 1999 The influence of maternal diet on breast cancer risk among female offspring. *Nutrition* 15:392-401.
5. **Liao DZ, Pantazis CG, Hou X, Li SA** 1998 Promotion of estrogen-induced mammary gland carcinogenesis by androgen in the male Noble rat: probable mediation by steroid receptors. *Carcinogenesis* 19:2173-80.
6. **Pietras RJ, Szego CM** 1999 Cell membrane estrogen receptors resurface. *Nat Med* 5:1330.
7. **Collins P, Webb C** 1999 Estrogen hits the surface. *Nat Med* 5:1130-1.
8. **Ruehlmann DO, Mann GE** 2000 Rapid non-genomic vasodilator actions of oestrogens and sex steroids. *Curr Med Chem* 7:533-41.
9. **Nuclear Receptors Nomenclature C** 1999 A unified nomenclature system for the nuclear receptor superfamily [letter]. *Cell* 97:161-3
10. **Beato M, Herrlich P, Schutz G** 1995 Steroid hormone receptors: many actors in search of a plot. *Cell* 83:851-7
11. **McGuire WL** 1975 Endocrine therapy of breast cancer. *Annu Rev Med* 26:353-63

12. **Green S, Walter P, Kumar V, Krust A, Bornert JM, Argos P, Chambon P** 1986 Human oestrogen receptor cDNA: sequence, expression and homology to v-erbA. *Nature* 320:134-139
13. **Kuiper G, Enmark E, Peltouhikko M, Nilsson S, Gustafsson JA** 1996 Cloning of a novel estrogen receptor expressed in rat prostate and ovary. *P.N.A.S.* 93:5925-5930
14. **Mosselman S, Polman J, Dijkema R** 1996 ER-beta - identification and characterization of a novel human estrogen receptor. *Febs Lett.* 392:49-53
15. **Kelley ST, Thackray VG** 1999 Phylogenetic analyses reveal ancient duplication of estrogen receptor isoforms. *J Mol Evol* 49:609-14
16. **Tora L, White J, Brou C, Tasset D, Webster N, Scheer E, Chambon P** 1989 The human estrogen receptor has two independent nonacidic transcriptional activation functions. *Cell* 59:477-87
17. **Cowley SM, Parker MG** 1999 A comparison of transcriptional activation by ER alpha and ER beta. *J Steroid Biochem Mol Biol* 69:165-75
18. **McInerney EM, Weis KE, Sun J, Mosselman S, Katzenellenbogen BS** 1998 Transcription activation by the human estrogen receptor subtype beta (ER-beta) studied with ER-beta and ER-alpha receptor chimeras. *Endocrinology* 139:4513-4522
19. **Hall JM, McDonnell DP** 1999 The estrogen receptor beta-isoform (ERbeta) of the human estrogen receptor modulates ERalpha transcriptional activity and is a key regulator of the cellular response to estrogens and antiestrogens. *Endocrinology* 140:5566-78
20. **Delaunay F, Pettersson K, Tujague M, Gustafsson JA** 2000 Functional differences between the amino-terminal domains of estrogen receptors alpha and beta. *Mol Pharmacol* 58:584-90.

21. **Krege JH, Hodgin JB, Couse JF, Enmark E, Warner M, Mahler JF, Sar M, Korach KS, Gustafsson JA, Smithies O** 1998 Generation and reproductive phenotypes of mice lacking estrogen receptor beta. *PNAS* 95:15677-82
22. **Couse JF, Korach KS** 1999 Estrogen receptor null mice: what have we learned and where will they lead us? *Endocr Rev* 20:358-417
23. **Saji S, Jensen EV, Nilsson S, Rylander T, Warner M, Gustafsson J** 2000 Estrogen receptors alpha and beta in the rodent mammary gland. *PNAS* 97:337-342
24. **Foley EF, Jazaeri AA, Shupnik MA, Jazaeri O, Rice LW** 2000 Selective loss of estrogen receptor beta in malignant human colon. *Cancer Res* 60:245-8
25. **Campbell-Thompson M, Lynch IJ, Bhardwaj B** 2001 Expression of estrogen receptor (ER) subtypes and ERbeta isoforms in colon cancer. *Cancer Res* 61:632-40.
26. **Iwao K, Miyoshi Y, Egawa C, Ikeda N, Noguchi S** 2000 Quantitative analysis of estrogen receptor-beta mRNA and its variants in human breast cancers. *Int J Cancer* 88:733-6.
27. **Roger P, Sahla ME, Makela S, Gustafsson JA, Baldet P, Rochefort H** 2001 Decreased Expression of Estrogen Receptor {beta} Protein in Proliferative Preinvasive Mammary Tumors. *Cancer Res* 61:2537-2541
28. **Rutherford T, Brown WD, Sapi E, Aschkenazi S, Munoz A, Mor G** 2000 Absence of estrogen receptor-beta expression in metastatic ovarian cancer. *Obstet Gynecol* 96:417-21.
29. **Leygue E, Dotzlaw H, Watson PH, Murphy LC** 1998 Altered estrogen receptor alpha and beta messenger RNA expression during human breast tumorigenesis. *Cancer Res* 58:3197-3201
30. **Pujol P, Rey JM, Nirde P, Roger P, Gastaldi M, Laffargue F, Rochefort H, Maudelonde T** 1998 Differential expression of estrogen receptor-alpha and -beta

- messenger RNAs as a potential marker of ovarian carcinogenesis. *Cancer Res* 58:5367-5373
31. **Lazennec G, Katzenellenbogen BS** 1999 Expression of human estrogen receptor using an efficient adenoviral gene delivery system is able to restore hormone-dependent features to estrogen receptor-negative breast carcinoma cells. *Mol Cell Endocrinol* 149:93-105
 32. **Graham FL, Smilet J, Russel WC, Nairn R** 1977 Characteristics of a human cell line transformed by DNA from adenovirus type 5. *J Gen Virol* 36:59-72
 33. **McGregory W.J., Bautista D.S., Graham FL** 1988 A simple technique for the rescue of early region 1 mutations into infectious human adenovirus type 5. *Virology* 163:614-617
 34. **Lazennec G, Alcorn JL, Katzenellenbogen BS** 1999 Adenovirus-mediated delivery of a dominant negative estrogen receptor gene abrogates estrogen-stimulated gene expression and breast cancer cell proliferation. *Mol Endocrinol* 13:969-80
 35. **Choi I, Ko C, Park-Sarge O-K, Nie R, Hess R, Graves C, Katzenellenbogen BS** 2001 Human estrogen receptor beta-specific monoclonal antibodies: Characterization and use in studies of estrogen receptor beta protein expression in reproductive tissues. *Mol Cell Endocrinol* in press:
 36. **Pettersson K, Delaunay F, Gustafsson JA** 2000 Estrogen receptor beta acts as a dominant regulator of estrogen signaling. *Oncogene* 19:4970-8.
 37. **Jeng MH, Jiang SY, Jordan VC** 1994 Paradoxical regulation of estrogen-dependent growth factor gene expression in estrogen receptor (ER)-negative human breast cancer cells stably expressing ER. *Cancer Lett* 82:123-128
 38. **Thomas TJ, Faaland CA, Adhikarakunnathu S, Watkins LF, Thomas T** 1998 Induction of p21 (CIP1/WAF1/SID1) by estradiol in a breast epithelial cell line

- transfected with the recombinant estrogen receptor gene: a possible mechanism for a negative regulatory role of estradiol. *Breast Cancer Res Treat* 47:181-93
39. **Touitou I, Vignon F, Cavailles V, Rochefort H** 1991 Hormonal regulation of cathepsin D following transfection of the estrogen or progesterone receptor into three sex steroid hormone resistant cancer cell lines. *J Steroid Biochem Mol Biol* 40:231-7
 40. **Wang W, Smith Rr, Burghardt R, Safe SH** 1997 17 beta-Estradiol-mediated growth inhibition of MDA-MB-468 cells stably transfected with the estrogen receptor: cell cycle effects. *Mol Cell Endocrinol* 133:49-62
 41. **Vanacker JM, Pettersson K, Gustafsson J, Laudet V** 1999 Transcriptional targets shared by estrogen receptor- related receptors (ERRs) and estrogen receptor (ER) alpha, but not by ERbeta. *Embo J* 18:4270-4279
 42. **Misiti S, Nanni S, Fontemaggi G, Cong YS, Wen J, Hirte HW, Piaggio G, Sacchi A, Pontecorvi A, Bacchetti S, Farsetti A** 2000 Induction of hTERT expression and telomerase activity by estrogens in human ovary epithelium cells. *Mol Cell Biol* 20:3764-71
 43. **Harris HA, Henderson RA, Bhat RA, Komm BS** 2001 Regulation of Metallothionein II Messenger Ribonucleic Acid Measures Exogenous Estrogen Receptor- β Activity in SAOS-2 and LNCaPLN3 Cells. *Endocrinology* 142:645-652
 44. **Cariou S, Donovan JC, Flanagan WM, Milic A, Bhattacharya N, Slingerland JM** 2000 Down-regulation of p21WAF1/CIP1 or p27Kip1 abrogates antiestrogen-mediated cell cycle arrest in human breast cancer cells. *PNAS* 97:9042-6
 45. **Weiss RH, Joo A, Randour C** 2000 p21Waf1/Cip1 Is an Assembly Factor Required for Platelet-derived Growth Factor-induced Vascular Smooth Muscle Cell Proliferation. *J Biol Chem* 275:10285-10290

46. **Di Cunto F, Topley G, Calautti E, Hsiao J, Ong L, Seth PK, Dotto GP** 1998 Inhibitory function of p21Cip1/WAF1 in differentiation of primary mouse keratinocytes independent of cell cycle control. *Science* 280:1069-72
47. **Lin VC, Ng EH, Aw SE, Tan MG, Bay BH** 2000 Progesterone induces focal adhesion in breast cancer cells MDA-MB-231 transfected with progesterone receptor complementary DNA. *Mol Endocrinol* 14:348-58
48. **Garcia M, Derocq D, Freiss G, Rochefort H** 1992 Activation of estrogen receptor transfected into a receptor-negative breast cancer cell line decreases the metastatic and invasive potential of the cells. *PNAS* 89:11538-11542
49. **Jiang SY, Jordan VC** 1992 Growth regulation of estrogen receptor-negative breast cancer cells transfected with complementary DNAs for estrogen receptor. *J Natl Cancer Inst* 84:580-91
50. **Kushner PJ, Hort E, Shine J, Baxter JD, Greene GL** 1990 Construction of cell lines that express high levels of the human estrogen receptor and are killed by estrogens. *Mol Endocrinol* 4:1465-1473
51. **Zajchowski DA, Sager R** 1991 Induction of estrogen-regulated genes differs in immortal and tumorigenic human mammary epithelial cells expressing a recombinant estrogen receptor. *Mol Endocrinol* 5:1613-23
52. **Cheng J, Malayer JR** 1999 Responses to stable ectopic estrogen receptor-beta expression in a rat fibroblast cell line. *Mol Cell Endocrinol* 156:95-105
53. **Jarvinen TA, Pelto-Huikko M, Holli K, Isola J** 2000 Estrogen receptor beta is coexpressed with ERalpha and PR and associated with nodal status, grade, and proliferation rate in breast cancer. *Am J Pathol* 156:29-35

54. **Omoto Y, Inoue S, Ogawa S, Toyama T, Yamashita H, Muramatsu M, Kobayashi S, Iwase H** 2001 Clinical value of the wild-type estrogen receptor beta expression in breast cancer. *Cancer Lett* 163:207-12.
55. **Mann S, Laucirica R, Carlson N, Younes PS, Ali N, Younes A, Li Y, Younes M** 2001 Estrogen receptor beta expression in invasive breast cancer. *Hum Pathol* 32:113-118.
56. **Speirs V, Malone C, Walton DS, Kerin MJ, Atkin SL** 1999 Increased expression of estrogen receptor beta mRNA in tamoxifen-resistant breast cancer patients. *Cancer Res* 59:5421-4
57. **Speirs V, Kerin MJ** 2000 Prognostic significance of oestrogen receptor beta in breast cancer. *Br J Surg* 87:405-9
58. **Paech K, Webb P, Kuiper G, Nilsson S, Gustafsson JA, Kushner PJ, Scanlan TS** 1997 Differential ligand activation of estrogen receptors ER-alpha and ER-beta at AP1 sites. *Science* 277:1508-1510
59. **Webb P, Nguyen P, Valentine C, Lopez GN, Kwok GR, McInerney E, Katzenellenbogen BS, Enmark E, Gustafsson JA, Nilsson S, Kushner PJ** 1999 The estrogen receptor enhances AP-1 activity by two distinct mechanisms with different requirements for receptor transactivation functions. *Mol Endocrinol* 13:1672-85
60. **Webb P, Lopez GN, Uht RM, Kushner PJ** 1995 Tamoxifen activation of the estrogen receptor/AP-1 pathway: potential origin for the cell-specific estrogen-like effects of antiestrogens. *Mol Endocrinol* 9:443-56
61. **Philips A, Chalbos D, Rochefort H** 1993 Estradiol increases and anti-estrogens antagonize the growth factor- induced activator protein-1 activity in MCF7 breast cancer cells without affecting c-fos and c-jun synthesis. *J Biol Chem* 268:14103-8

62. **Platet N, Cunat S, Chalbos D, Rochefort H, Garcia M** 2000 Unliganded and liganded estrogen receptors protect against cancer invasion via different mechanisms [In Process Citation]. *Mol Endocrinol* 14:999-1009
63. **Tong D, Czerwenka K, Sedlak J, Schneeberger C, Schiebel I, Concin N, Leodolter S, Zeillinger R** 1999 Association of in vitro invasiveness and gene expression of estrogen receptor, progesterone receptor, pS2 and plasminogen activator inhibitor-1 in human breast cancer cell lines. *Breast Cancer Res Treat* 56:91-7.
64. **Thompson EW, Paik S, Brunner N, Sommers CL, Zugmaier G, Clarke R, Shima TB, Torri J, Donahue S, Lippman ME, et al.** 1992 Association of increased basement membrane invasiveness with absence of estrogen receptor and expression of vimentin in human breast cancer cell lines. *J Cell Physiol* 150:534-44.
65. **Sommers CL, Byers SW, Thompson EW, Torri JA, Gelmann EP** 1994 Differentiation state and invasiveness of human breast cancer cell lines. *Breast Cancer Res Treat* 31:325-35
66. **Madsen MW, Briand P** 1990 Relationship between tumorigenicity, in vitro invasiveness, and plasminogen activator production of human breast cell lines. *Eur J Cancer* 26:793-7