

**Sugar sensing by enterocytes combines polarity,
membrane bound detectors and sugar metabolism.**

Maude Le Gall, Vanessa Tobin, Emilie Stolarczyk, Véronique Dalet, Armelle
Leturque, Edith Brot-Laroche

► **To cite this version:**

Maude Le Gall, Vanessa Tobin, Emilie Stolarczyk, Véronique Dalet, Armelle Leturque, et al.. Sugar sensing by enterocytes combines polarity, membrane bound detectors and sugar metabolism.. Journal of Cellular Physiology, Wiley, 2007, 213 (3), pp.834-43. 10.1002/jcp.21245 . inserm-00170149

HAL Id: inserm-00170149

<https://www.hal.inserm.fr/inserm-00170149>

Submitted on 4 Sep 2009

HAL is a multi-disciplinary open access archive for the deposit and dissemination of scientific research documents, whether they are published or not. The documents may come from teaching and research institutions in France or abroad, or from public or private research centers.

L'archive ouverte pluridisciplinaire **HAL**, est destinée au dépôt et à la diffusion de documents scientifiques de niveau recherche, publiés ou non, émanant des établissements d'enseignement et de recherche français ou étrangers, des laboratoires publics ou privés.

1 **Sugar sensing by enterocytes combines polarity, membrane bound**
2 **detectors and sugar metabolism**

3
4 **Maude Le Gall^{1,2,3,#,*}, Vanessa Tobin^{1,2,3,#}, Emilie Stolarczyk^{1,2,3}, Véronique Dalet^{1,2,3},**
5 **Armelle Leturque^{1,2,3}, Edith Brot-Laroche^{1,2,3}**

6
7 1 INSERM, UMR S 872, Centre de Recherche des Cordeliers, Paris F-75006 France.

8 2 Université Pierre et Marie Curie-Paris6, UMR S 872, Paris F-75006 France.

9 3 Université Paris Descartes, UMR S 872, Paris F-75006 France.

10

11 #MLG and VT have equally contributed to the work.

12

13 * **Correspondence to** : Maude Le Gall, UMRS 872 Centre de Recherche des Cordeliers, 15 rue de
14 l'Ecole de Médecine, Paris, F-75006 France. Tel : + 33 1 42 34 68 99. Fax: + 33 1 43 25 16 15. Email:
15 mlegall@bhdc.jussieu.fr

16

17 Running head : Sugar signalling pathways in enterocytes

18 **Keywords**: glucose signalling, enterocyte, GLUT2, sugar metabolism, sweet taste receptor.

19 Total number of text figures and tables : 6 figures

20 Funded by: ALFEDIAM Merck Lipha, Institut Benjamin Delessert, AIP ATC Nutrition;

21 Grant Number: ASEO22129DSA.

22

23 Abbreviations : SI: sucrase isomaltase, L-PK: liver-pyruvate kinase, SGLT1: sodium glucose
24 transporter, GPCR: G protein-coupled receptor, T1R: Taste Receptor type 1, 3OMG: 3-O-
25 methyl glucose, N-AGA: N-acetylglucosamine, G-6-Pase : glucose-6-phosphatase, EGFP:
26 enhanced green fluorescent protein, ChREBP: carbohydrate response element binding protein,
27 ChoRE: carbohydrate response element, SREBP: sterol response element binding protein,
28 LXR: liver X receptor, PKA: protein kinase A, ADRP, adipose differentiation-related protein.

1
2
3
4
5
6
7
8
9
10
11
12
13
14
15
16
17
18
19
20
21

ABSTRACT

Sugar consumption and subsequent sugar metabolism are known to regulate the expression of genes involved in intestinal sugar absorption and delivery. Here we investigate the hypothesis that sugar-sensing detectors in membranes facing the intestinal lumen or the bloodstream can also modulate intestinal sugar absorption. We used wild-type and GLUT2-null mice, to show that dietary sugars stimulate the expression of sucrase-isomaltase (SI) and L-pyruvate kinase (L-PK) by GLUT2-dependent mechanisms, whereas the expression of GLUT5 and SGLT1, did not rely on the presence of GLUT2. By providing sugar metabolites, sugar transporters, including GLUT2, fuelled a sensing pathway. In Caco2/TC7 enterocytes, we could disconnect the sensing triggered by detector from that produced by metabolism, and found that GLUT2 generated a metabolism-independent pathway to stimulate the expression of SI and L-PK. In cultured enterocytes, both apical and basolateral fructose could increase the expression of GLUT5, conversely, basolateral sugar administration could stimulate the expression of GLUT2. Finally, we located the sweet-taste receptors T1R3 and T1R2 in plasma membranes, and we measured their cognate Galpha Gustducin mRNA levels. Furthermore, we showed that a T1R3 inhibitor altered the fructose-induced expression of SGLT1, GLUT5 and L-PK. Intestinal gene expression is thus controlled by a combination of at least three sugar-signalling pathways triggered by sugar metabolites and membrane sugar receptors that, according to membrane location, determine sugar sensing polarity. This provides a rationale for how intestine adapts sugar delivery to blood and dietary sugar provision.

INTRODUCTION

1
2
3 Intestinal sugar delivery depends on the levels of expression of disaccharidases (i.e. sucrase-
4 isomaltase, SI) and sugar transporters. Indeed, dietary glucose is transported across the apical
5 membrane of enterocytes by the energy-dependent sodium-glucose co-transporter 1 (SGLT1)
6 and dietary fructose by the facilitative transporter GLUT5 (Levin, 1994). At the basolateral
7 membrane, glucose and fructose exit the intestinal epithelia mainly via GLUT2-dependent
8 facilitated diffusion (Levin, 1994) to reach the bloodstream. In GLUT2-null mice, glucose
9 delivery seems to be mediated via vesicular trafficking (Stumpel et al., 2001) and the
10 mechanism by which fructose reaches the blood stream is probably dependent on the presence
11 of GLUT5 in the basolateral membrane (Blakemore et al., 1995). In the case of a sugar rich
12 meal, GLUT2 can be recruited into the apical membrane where it complements SGLT1 and
13 GLUT5 transport capacities (Kellett and Brot-Laroche, 2005). Furthermore, the small
14 intestine adapts to repeated sugar ingestion by increasing transporter expression over the
15 course of a few days (Burant and Saxena, 1994; Goda, 2000; Inukai et al., 1993). Thus, the
16 regulation of the transcription, expression and location of sugar transporters organizes sugar
17 absorption. In pathological conditions this regulation can amplify sugar delivery creating a
18 vicious circle. Indeed, in rat, streptozotocin-induced diabetes increases SGLT1, GLUT2 and
19 GLUT5 mRNA levels and sugar absorption (Burant et al., 1994; Ferraris et al., 1993),
20 possibly causing exacerbated postprandial blood glucose excursion. Knowledge of the
21 mechanisms of sugar sensing at the luminal and basolateral enterocyte membranes is thus
22 essential for an understanding of the mechanism of induction of sugar-sensitive genes.

23 The regulation by sugars of gene expression is highly conserved through evolution, and is
24 found in bacteria (Jacob and Monod, 1961), yeast (Johnston, 1999), plants (Villadsen and
25 Smith, 2004), and mammals (Girard et al., 1997; Towle, 2005). In pancreatic β cells, glucose
26 is known to alter the expression of more than 150 genes that can be grouped in functional
27 clusters, including insulin secretion, energy metabolism, membrane transport, signalling
28 pathways, gene transcription and protein synthesis/degradation (Schuit et al., 2002). Fructose
29 is known to regulate 50 genes in the small intestine of neonate rats, where it affects genes
30 encoding for ion and hexose-transporters as well as enzymes of hexose metabolism (Cui et
31 al., 2004).

32 Considering the number and functional variety of sugar regulated genes, and given their
33 location in diverse sugar-sensitive tissues, a single signalling pathway is unlikely to explain

1 all cellular adaptations to life in a sugar-containing environment. Our current understanding
2 of how cells respond to sugar is mainly based on studies in yeast and involves signalling
3 pathways either initiated from hexose metabolites directly or from the direct engagement of
4 hexoses with their cognate sugar receptors at the cell surface (Holsbeeks et al., 2004). In
5 mammals, studies have been primarily focused on glucose metabolism and related signals
6 (Girard et al., 1997; Towle, 2005). However, non metabolisable sugar analogues can stimulate
7 SGLT1 expression (Miyamoto et al., 1993). Some sugar receptors have been identified in the
8 plasma membrane of mammalian cells. A first group of mammalian sugar receptors is
9 composed of sugar transporters, which can also function as sugar detectors. This group
10 includes GLUT2 and SGLT3. Indeed, in hepatocytes GLUT2 can fuel sugar metabolism and
11 also trigger a receptor-dependent protein signalling cascade (Guillemain et al., 2000;
12 Guillemain et al., 2002). SGLT3 functions in cholinergic neurons neighboring enterocytes
13 and while it does not transport glucose, it induces membrane currents upon sodium-dependent
14 glucose binding (Diez-Sampedro et al., 2003). A second group of sugar detectors contains
15 members of the G Protein–Coupled Receptor (GPCR) family. Among them are the sweet
16 taste receptors of type 1, T1R, located on the tongue. Heterodimeric T1R3/T1R2 and T1R3
17 homodimers form sweet taste receptors that bind fructose or sucrose, leading to adenylate-
18 cyclase signalling cascades (Nelson et al., 2001; Xu et al., 2004; Zhao et al., 2003).
19 Biochemical data suggest that GPCRs participate to sugar signalling in an enteroendocrine
20 cell line, but their molecular identification has yet to be determined (Rozenfurt, 2006). In
21 enterocytes, cell polarity further complicates the regulation of sugar sensitive genes. Indeed,
22 detectors may conceivably be activated by dietary sugars at the apical membrane and by
23 blood glucose at the basolateral membrane of enterocytes.

24 In this study we consider both metabolism-driven sugar signalling and sugar detection
25 initiated signalling pathways. By monitoring the induction of sugar sensitive genes both *in*
26 *vivo* and in Caco-2/TC7 cells we report here the identification of distinct sugar sensing
27 signalling pathways with a striking polar distribution in enterocytes. These sugar sensory
28 mechanisms offer insight not only into the modes of sugar absorption and delivery, but also
29 unveil important avenues for the development of novel pharmacological compounds for
30 improved glycaemic control.

MATERIALS AND METHODS

Mice: Wild-type mice were from C57Bl/6 strain (Janvier, France). GLUT2-null mice (RIP GLUT1XGLUT2^{-/-}) (Guillam et al., 1997) were bred in the transgenic animal facility of IFR58 (Paris, France). All animal procedures are complied with recommendations for the use of laboratory animals from the French administration. Mice were fed for 5 days with the experimental diets, which contained either low amounts of sugar, or 65% (W/W) glucose or fructose as previously described (Gouyon et al., 2003a). The jejunum is here defined as the part of the small intestine starting at the Treitz ligament and excluding its last third. Intestinal samples, taken from fed mice, were rapidly everted and contents washed out in ice cold PBS before mucosa scrapings were taken. Mucosa was dispersed in RNA extraction buffer and snap frozen in liquid nitrogen.

Cell culture: Caco-2/TC7 cells were seeded at 6×10^5 cells/cm² either on six-well, solid or porous (3µm high pore density) supports (Becton Dickinson, Meylan, France). Cells were grown in complete Dulbecco's modified Eagle's medium (25 mmol.L⁻¹ glucose DMEM, Gibco, Paisley, U.K) supplemented with 20% heat-inactivated (30min, 56°C) fetal calf serum (FCS) (AbCys, Paris, France). Media were renewed every 24 hours for at least 10 days to allow differentiation of the cells (Chantret et al., 1994; Mahraoui et al., 1994).

Post-confluent, differentiated cells were switched from standard growth media (DMEM 25 mmol.L⁻¹ glucose) to glucose-free DMEM supplemented with 10% heat-inactivated FCS, and contained less than 1 mmol.L⁻¹ glucose (low sugar medium). According to need, media were supplemented with sugar for 2 to 4 days as indicated in the legend of the figures. For sugar metabolim assay; 25 mmole.L⁻¹ 3-O-Methylglucoside or 250 mmole.L⁻¹ N-AGA were added at both apical and basal sides of the cells for 48 hours. For sweet taste receptor functional assay, after differentiation, cells were grown in glucose and glutamine free DMEM supplemented with 10% dialyzed FCS. Inhibitor, 1 mmol.L⁻¹ lactisole (Sigma Aldrich, Saint Quentin, France) and substrate 25 mmol.L⁻¹ fructose, were added during the last 48 hours of culture at both apical and basal poles.

When grown on porous support, cell viability and cell monolayer integrity after treatments were estimated by measure of the transepithelial electrical resistance (TEER), which is a witness of tight junction integrity and of ion pump function in cell membranes (Grasset et al., 1984).

1 *Immunofluorescence*: Differentiated Caco-2/TC7 cells grown on filters were fixed with 4%
2 Paraformaldehyde (Sigma Aldrich, Saint Quentin, France), permeabilized with 0.2% Triton
3 (Sigma Aldrich, Saint Quentin, France) and labeled with antibodies targeting T1R2 (T-20: sc-
4 22456) and T1R3 (N-20: sc-22458) (Santa Cruz, Biotechnology, Tebu France). Images were
5 produced by confocal microscopy (Zeiss LSM510 software).
6

7 *Human GLUT2, GLUT5 and sucrase-isomaltase promoter constructs*: Caco-2/TC7 cells were
8 transfected with the following promoter regions inserted into p205 plasmid driving the
9 reporter gene luciferase (Rodolosse et al., 1996) : -1100/+300 of the hGLUT2 promoter
10 (generous gift of GI Bell, University of Chicago, Chicago, IL, USA), -3600/+60 of the hSI
11 promoter (Rodolosse et al., 1997) and -2500/+21 of the hGLUT5 promoter (Mahraoui et al.,
12 1994). Populations of stably transfected cells were established. Protein assays were made with
13 the BCA kit (Pierce, Interchim, Montluçon, France). Maximal variations of protein
14 concentrations between different cell cultures and different culture conditions were below
15 20%. Luciferase activities were measured using the Luciferase assay kit (Promega,
16 Charbonnières les Bains, France) in a Lumat LB9501 luminometer (Berthold Detection
17 System, Pforzheim, Germany). Results were expressed as relative light units $\text{RLU}\cdot\text{sec}^{-1}\cdot\mu\text{g}^{-1}$
18 protein.
19

20 *EGFP-GLUT2-loop and -C-terminus peptide constructs*: The coding regions of the
21 intracellular loop between transmembrane domain 6 and 7 (amino acids 237-301) and the C-
22 terminus GLUT2 domain (amino acids 481-521) of rat GLUT2 were cloned in frame with
23 EGFP in pEGFP-C (Clontech BD Biosciences, le Pont de Claix, France), as previously
24 described (Guillemain et al., 2000). The EGFP-GLUT2 domain cassettes were placed
25 downstream the SV40 promoter of pGL3 (Promega, Charbonnières les Bains, France),
26 allowing the expression of EGFP-GLUT2 domains in differentiated Caco-2/TC7 cells. Stably
27 transfected EGFP positive cells were established and sorted by FACS (Epics Altra, Beckman
28 Coulter, Roissy, France). The cells were secondarily and transiently transfected with the
29 hGLUT2 promoter using the lipofectin transfection kit (Life Technologies, Cergy Pontoise,
30 France).
31

32 *Messenger RNA*: Total RNA from the jejunum of mice or from Caco2/TC7 cells were
33 extracted using TriReagent (MRC, Interchim, Montluçon, France). Mouse and human

1 GLUT2, L-Pyruvate kinase (L-PK), Glucose-6-phosphatase (G-6-Pase), Sucrase-Isomaltase
 2 (SI) and human SGLT1 and GLUT5 mRNAs were quantified by reverse transcription and
 3 real-time PCR using the Light-Cycler System according to the manufacturer's procedures
 4 (Roche Molecular Biochemicals, Meylan, France) as previously described (Gouyon et al.,
 5 2003a; Guillemain et al., 2002). The primers used were for hGLUT5 forward 5'-
 6 TCTCCTTGCAAACGTAGATGG-3' and reverse 5'-GAAGAAGGGCAGCAGAAGG-3', for hSGLT1
 7 forward 5'-TGGCAATCACTGCCCTTTA-3' and reverse 5'-TGCAAGGTGTCCGTGTAAT-3' and for
 8 hADRP forward 5'-GTGAGATGGCAGAGAACGGTGTG- 3' and reverse 5'-
 9 TGCCCTTTGGTCTTGTCCA-3'. All primer pairs amplified a single amplicon as indicated by the
 10 unique melting temperature of the PCR product. Moreover, we verify the size and specificity
 11 of the amplicon by restriction enzyme analysis and electrophoresis. The large ribosomal
 12 protein L19 was used as a control gene since its expression level varied by less than 20% in
 13 all the culture conditions applied to the cells. Results were expressed as ratios of gene mRNA
 14 over L19 mRNA levels. Northern blots were performed to measure mouse intestinal GLUT5
 15 and SGLT1 mRNA. Density analyses (Gel Analyst 3.02 software) were expressed as the ratio
 16 of mRNA to 18S rRNA. For human taste receptors identification, primers were selected using
 17 published data (Rozenfurt et al., 2006) or predicted sequences available in the NCBI
 18 database: NM_152232 for T1R2, NM_152228 for T1R3, and XM_001129050, XM_294370
 19 and X_M939789 for Gustducin. T1R2: forward 5'-GTATGAAGTGAAGGTGATAGGC-3' and
 20 reverse 5'-GGGTAGACCACCCTCTTGG-3'; T1R3: forward 5'-CAAGTTCTTCAGCTTCTTCCTC-3'
 21 and reverse 5'-GTACATGTTCTCCAGGAGCTGC-3'; Gustducin: forward 5'-
 22 GCCAAATACATTTGAAGATGCAGG-3' and reverse 5'-GCACTTCTGGGATTACATAATC-3'. Note
 23 that for T1R2 and T1R3 nested primers were also used. Nested T1R2: forward 5'-
 24 TGCGCTTCGCGGTGGAGG-3' and reverse 5'-CAGCCGAGGAGGCTGTGC-3'; nested T1R3: forward
 25 5'-GGTCAGCTACGGTGCTAGC-3' and reverse 5'-AGCCTGAGGCGTTGCACTG-3'.

26

27 *Sugar transepithelial transfer:* The glucose or fructose content of the apical and basal culture
 28 media was assayed with an enzymatic assay kit (Sigma Aldrich, Saint Quentin, France). The
 29 non-specific transpithelial transfer of sugar across Caco-2/TC7 cells grown on porous support
 30 was measured using (1-³H NEN) L-glucose. L-glucose transfer was lower than 1.5 % of the
 31 D-isomers (3 independent experiments, data not shown) indicating that epithelia were tight.
 32 Using fructose media helped to distinguish fructose and glucose transporting GLUTs.

33

- 1 *Statistics* : All statistical analyses were made using ANOVA and student T tests (PRISM
- 2 software).

RESULTS

Regulation of intestinal genes by dietary sugars

In vivo experiments were first conducted to determine if glucose and fructose employ distinct signalling pathways to induce sugar-regulated genes. We focused our study on sugar transporters (GLUT5, GLUT2, SGLT1) and sugar metabolism enzymes (SI and L-PK). As previously observed for GLUT2 (Gouyon et al., 2003a), glucose- or fructose-rich diets efficiently stimulated the mRNA accumulation of these sugar-regulated genes in the jejunal mucosa of wild-type (WT) mice (Figure 1). SGLT1, SI and L-PK mRNA increased to similar levels under both glucose and fructose dietary regimens (Figure 1). On the other hand, GLUT5 mRNA increased in mice fed fructose- but not glucose-rich diets (compare Figure 1A to 1B), in agreement with previous studies (Burant and Saxena, 1994; Gouyon et al., 2003b) indicating that cells discriminate glucose and fructose signals to regulate the expression of GLUT5. As glucose and fructose are both catabolized through glycolysis in enterocytes, metabolic signalling via glycolysis cannot fully account for the differential effect. We used GLUT2-null mice to document the contribution of this glucose/fructose transporter to the regulation of sugar-sensitive genes. The mRNA levels were similar for the different genes analysed in GLUT2-null and WT mice fed low carbohydrate diets. GLUT2-null and WT mice exhibited indistinguishable increases of GLUT5 mRNA in response to fructose-rich diet. Similarly, the small but significant induction of SGLT1 by sugar-rich diets was similar in GLUT2-null and WT mice. By contrast, SI and L-PK mRNA inductions were not observed or were dramatically reduced in GLUT2-null mice. Therefore, a GLUT2-dependent signalling pathway is required to obtain SI and L-PK gene regulation by dietary glucose and fructose. Interestingly, feeding GLUT2-null mice with fructose produced a partial accumulation of L-PK mRNA (Figure 1B), indicating that fructose and glucose signalling pathways can differ in the intestine. These *in vivo* data indicate that enterocytes exploit several sugar signalling pathways to adapt the expression of sugar target genes to the dietary environment.

Effect of sugars on sensitive genes in Caco-2/TC7 enterocytes

Enterocyte detection of dietary sugars could occur at their apical membranes exposed to the intestinal lumen content or their basolateral membranes exposed to the blood. To assess the sugar detection capacity of each of these membranes in a controlled manner, we used differentiated human colon carcinoma, Caco-2/TC7 cells that display the morphological characteristics and functional properties of intestinal absorptive cells (Delie and Rubas, 1997).

1 Caco-2/TC7 cells cultured on solid support have only their apical poles in contact with culture
2 media, whereas cultures grown on porous support permit sugar provision to either apical or
3 basolateral membranes. This system was thus employed to differentially expose enterocytes
4 to dietary sugars and to determine polarity in sugar detection and signalling.

5 As shown in Figure 2, fructose stimulated hGLUT5 promoter activity by 2 fold in Caco-
6 2/TC7 cells regardless of whether they were grown on solid or porous support. By contrast,
7 neither glucose nor fructose could stimulate the hGLUT2 promoter activity in cells grown on
8 solid support (Figure 2), although the same hGLUT2 promoter fragment has been shown
9 capable of being activated by glucose in pancreatic MIN6 cells (Cassany et al., 2004).
10 However, a net stimulation of hGLUT2 promoter activity was observed in Caco-2/TC7 cells
11 grown on porous support in agreement with the capacity of dietary sugars to induce GLUT2
12 *in vivo*. Thus growing enterocytes on porous support with their basal poles capable of
13 detecting sugars unmasks polarity in sugar signalling toward GLUT2 gene regulation.

14 We took advantage of the restricted localization of SI at the apical membrane of enterocytes
15 to further study the mechanisms of polarity in sugar detection and signalling. The addition of
16 sucrose into apical medium stimulated hSI promoter activity significantly (Figure 2), but no
17 activation was obtained when sucrose addition was restricted to media exposed to the
18 basolateral membrane. Furthermore, sucrose cannot be simply hydrolyzed to fructose and
19 glucose at the basolateral membrane as it lacks SI activity. These data also indicate that
20 epithelial tight junctions were functional. Importantly, the glucose and fructose moieties of
21 sucrose stimulated the hSI promoter activity irrespective of the compartment of addition
22 (Figure 2). Thus, enterocytes growing on porous support display polarity in their responses to
23 dietary sugars.

24 When administered independently to the apical or basal compartments, glucose and fructose
25 concentrations equilibrated between the apical and basolateral media within 12 hours (Figure
26 3A). Therefore, in our culture conditions, polarized sugar signals could only be detected
27 within 12 hours (i.e. before sugar equilibration is achieved). We measured the time course of
28 hGLUT2 promoter activity induced by sugar (Figure 3B, left panel) and unfortunately, the
29 addition of sugar required 48 hours to maximally stimulate the hGLUT2 promoter and little to
30 no stimulation was observed at 12 hours (Figure 3B, left panel). Accordingly, the addition of
31 glucose to either pole of the cell for 48h (i.e. long after sugar equilibration is achieved),
32 produced a similar stimulation of the hGLUT2 promoter (not shown).

1 By contrast, 12 hours of sugar removal were sufficient to significantly decrease hGLUT2
2 promoter activity and near basal levels were obtained after 24 hours (Figure 3B right panel).
3 After 12 hours, hGLUT2 promoter activity was significantly higher in cells deprived of apical
4 sugar but supplied with glucose at the basal pole (Figure 3C) whereas strict apical provision
5 of sugar did not activate the hGLUT2 promoter significantly within the same time course
6 (Figure 3C). This is congruent with the inability of sugars to stimulate the hGLUT2 promoter
7 when the basolateral membranes of Caco-2/TC7 cells are attached on solid support (Figure 2).
8 These results suggested that basolateral glucose provides a sugar signal able to maintain the
9 transcription of hGLUT2. Moreover, the difference in hGLUT2 promoter stimulation between
10 strictly basal versus basal and apical sugar supply might reveal a role for sugar metabolic
11 fluxes in these signalling processes. Lastly, we found the activation of the hGLUT2 promoter
12 exhibited saturation at 20 mmol.L⁻¹ (data not shown) in accordance with the kinetic
13 parameters of the GLUT2 transporter (Gould et al., 1991), the main sugar transporter in the
14 basolateral membrane of enterocytes (Cheeseman and O'Neill, 1998).

15 **Sugar sensitive genes are induced by metabolism**

16 Sugar metabolism is an essential component of the regulation of sugar sensitive genes in
17 hepatocytes and adipocytes. We investigated the contribution of metabolic signalling to the
18 expression of selected glucose-sensitive genes in enterocytes, as GLUT2, L-PK and glucose-
19 6-phosphatase (G-6-Pase). The glucose analogue, 3-O-methylglucose (3OMG) is a substrate
20 for SGLT1 and GLUT2 that can be transported but not metabolized. Provision of 3OMG to
21 Caco-2/TC7 cells did not increase hGLUT2 promoter activity even when cells were cultured
22 on porous support (Figure 4A), suggesting a metabolic requirement for the induction of this
23 glucose-sensitive gene. However, the structural difference introduced by glucose methylation
24 might not be suitable for all sugar-signalling machineries. The contribution of metabolism
25 was thus further analyzed by inhibiting glucose phosphorylation into glucose-6-phosphate. N-
26 acetylglucosamine (N-AGA) is an efficient inhibitor of hexokinases, it did not affect the
27 expression of genes that were not regulated by sugar like Adipose Differentiation-Related
28 Protein (ADRP) (Figure 4B) or Cyclophilin A (not shown). N-AGA completely prevented
29 glucose-dependent G-6-Pase mRNA increase as expected from earlier reports demonstrating
30 that G-6-Pase promoter activity is strictly regulated by glucose metabolism (Massillon, 2001)
31 (Figure 4B). However, N-AGA only partly prevented GLUT2 and L-PK mRNA increases
32 (Figure 4B). These data confirmed that an active metabolic pathway is necessary to sustain a
33 proper transcriptional response to glucose in enterocytes. Interestingly, N-AGA also reduced

1 GLUT2 mRNA levels in the absence of added extracellular glucose, indicating that
2 metabolism contributes to basal promoter activity due to the low levels of glucose provided
3 by the serum (Figure 4B). However, inhibition of glucose metabolism did not abolish sugar-
4 dependent increases of GLUT2 and L-PK mRNAs, suggesting either the existence of some
5 additional sugar signalling pathways or an incomplete block of hexokinases by N-AGA.

6 **Additional sugar signalling pathways exist in enterocytes**

7 *GLUT2-loop mediated pathway*

8 Both our data in mouse jejunal mucosa and our data in Caco-2/TC7 cells show that
9 metabolism is necessary but not sufficient to fully explain the response of sugar sensitive
10 genes to glucose or fructose in enterocytes. Therefore, the possibility that GLUT2 could
11 function as a sugar detector and thereby mediate sugar signalling was tested. Indeed, in
12 mhAT3F hepatoma cells, we previously showed that GLUT2 is capable of triggering glucose-
13 dependent signalling by interacting with the karyopherin alpha2 protein through its large
14 cytoplasmic loop between transmembrane domain 6 and 7 (Guillemain et al., 2000). We here
15 asked if different GLUT2 constructs could functionally alter dietary sugar sensing. Two
16 intracellular domains of GLUT2 were expressed in Caco-2/TC7 enterocytes, to test their
17 effect on sugar-sensitive gene expression. Cells that stably expressed EGFP-GLUT2-loop,
18 EGFP-GLUT2-Cterminus or EGFP alone, were obtained by cell sorting and cells were
19 selected for similar levels of transgene expression (Figure 5A, upper panel). As expected,
20 EGFP-GLUT2-loop exhibited a nuclear location whereas EGFP or EGFP-GLUT2-Cterminus
21 did not present any specific location (Figure 5A, lower panel). In cells expressing EGFP
22 alone, as in non-transfected cells, fructose and glucose increased endogenous hGLUT2
23 mRNA levels by 4- and 3-fold respectively (compare Figure 5B and 2). EGFP-GLUT2-
24 Cterminus expression did not significantly change endogenous hGLUT2 mRNA
25 accumulation in response to sugar (Figure 5B). By contrast, EGFP-GLUT2-loop expression
26 reduced endogenous hGLUT2 mRNA accumulation by 60% in response to fructose and by
27 90% in response to glucose as compared to controls (Figure 5B). Similar results were
28 obtained for hL-PK mRNA (Figure 5C) and hSI mRNA (not shown) in response to glucose.

29 To determine whether these strong inhibitions were due to reduced transcription, we
30 measured transiently transfected hGLUT2 promoter activity in the stable lines expressing
31 EGFP-GLUT2-loop and EGFP-GLUT2-Cterminus (Figure 5D). In accordance with the
32 mRNA accumulation data, EGFP alone and EGFP-GLUT2-Cterminus did not interfere with

1 the fructose-dependent stimulation of hGLUT2 promoter activity, whereas EGFP-GLUT2-
2 loop partially inhibited it.

3 We assumed that the remaining stimulations of GLUT2 and L-PK observed in EGFP-
4 GLUT2-loop expressing cells were due to a sugar metabolism pathway. N-AGA could further
5 inhibit endogenous GLUT2 mRNA accumulation in EGFP-GLUT2-loop cells (Figure 5E), as
6 well as L-PK mRNA accumulation. These results indicate that both metabolic and GLUT2-
7 dependent signalling are necessary for the regulation of sugar sensitive genes.

8

9 *Sweet-taste receptor mediated sugar signalling*

10 As sweet-taste receptors of the GPCR family have been implicated in fructose and sucrose
11 signalling, we questioned the putative role of T1R sweet taste receptors in enterocytes
12 challenged by fructose. Functional assays were performed in differentiated Caco-2/TC7
13 enterocytes cultured in the presence of T1R3 ligands. Lactisole, a T1R3 inhibitor, decreased
14 the stimulation by fructose of human SGLT1, GLUT5, and L-PK mRNA expression (Figure
15 6A), underlining the involvement of T1R3 in fructose signalling. By contrast, neither GLUT2
16 mRNA levels (Figure 6A) nor hGLUT2 promoter activity (not shown) were changed by
17 lactisole, suggesting that T1R3 did not control GLUT2 expression. By RT-PCR, the T1R3
18 receptor expression was revealed, as expected from functional assays. Its cognate G protein,
19 Gustducin ($G_{\alpha\text{gust}}$) also known as GNAT3 (Guanine nucleotide binding protein, alpha
20 transducing 3) was identified (Figure 6B). Moreover, neither T1R3 nor Gustducin mRNA
21 expressions were modified by glucose or fructose (Figure 6B). T1R2 and T1R3 proteins were
22 recovered in plasma membranes of fully differentiated Caco-2/TC7 enterocytes by confocal
23 analysis (Figure 6C). We could locate the sweet-taste receptors mainly at the basolateral
24 membranes of enterocyte cells (Figure 6C).

25

DISCUSSION

In this study we show the convergence of several sugar signalling pathways on the regulation of intestinal sugar-sensitive target genes. In addition to the better understood metabolic signalling mechanisms, we identified GLUT2 and GPCRs to be important sugar sensing signalling initiators governing enterocyte gene regulation. We identified a pathway strictly requiring GLUT2, since it was absent in GLUT2-null mice and could be blocked in dominant-negative fashion in cells expressing a large amount of the cytoplasmic loop domain of GLUT2. Particularly important was finding the polar nature of GLUT2 sugar sensing and signalling from the basolateral and not the apical enterocyte membrane. Thus as enterocytic GLUT2 senses hyperglycaemia it could further increase sugar delivery to the blood by the up-regulation of sugar transport and metabolism genes.

The GLUT2 pathway had different and complementary effects compared to the well-studied metabolic pathways with respect to the genes they regulated, suggesting differences in the molecular mechanics of these two pathways. In mammalian cells, sugar metabolites, including glucose-6-phosphate or xylulose-5-phosphate can mediate the regulation of sugar-sensitive gene expression (Vaulont et al., 2000). Both metabolites are thought to accumulate in the cells and to convey sugar signals to the transcription machinery. In hepatocytes, xylulose-5-phosphate has been proposed to activate protein phosphatase 2A (Nishimura and Uyeda, 1995), which in turn would dephosphorylate a transcription factor named carbohydrate response element binding protein (ChREBP). ChREBP would then be transported into the nucleus to stimulate the transcription of glucose-sensitive genes (Uyeda et al., 2002). ChREBP is very abundant in the intestine (Uyeda et al., 2002), but it might not be the only transcription factor to regulate glucose-sensitive genes. Indeed, all glucose-sensitive promoters do not contain ChREBP binding sites (ChoRE, carbohydrate response element). For instance, in hepatocytes, the GLUT2 promoter rather binds SREBP-1C (sterol response element binding protein) on a sterol responsive element (Im et al., 2005). SREBP-1C also stimulates the transcription of lipogenic genes in response to glucose and insulin (Wang et al., 2003). In addition, in pancreatic cells, SREBP-1C was emphasized over ChREBP in the activation of the transcription of GLUT2 and Insulin Receptor Substrate by glucose (Wang et al., 2005). The metabolite that activates SREBP-1C has not yet been identified, and the specific role of SREBP-1C in glucose-induced gene expression in the intestine is unknown. Sugar transport and metabolism in enterocytes rely on the abundance and kinetic parameters

1 of several transporters. Among them, the high K_m and high V_{max} transporter GLUT2 is
2 abundant in the basolateral membranes allowing a large influx of sugars and providing
3 abundant fuel for the metabolic pathway in enterocytes.

4 In hepatocytes, GLUT2 not only triggers a metabolic-signal but it also triggers a protein
5 signalling cascade (Guillemain et al., 2002). In Caco-2/TC7 cells, high expression of the large
6 cytoplasmic loop between transmembrane domain 6 and 7 of GLUT2 inhibited the
7 stimulation by sugars of GLUT2 and L-PK and its expression was restricted to the nucleus as
8 in hepatoma cells. Since the inhibition by the GLUT2-loop was partial and could be further
9 inhibited by hexokinase inhibition, we concluded that the metabolic and GLUT2 pathways are
10 both required in enterocytes. The nuclear importer karyopherin alpha2 can bind to the GLUT2
11 loop, furthermore the rate of its nuclear shuttling is regulated by glucose in hepatoma cells
12 (Cassany et al., 2004). This nuclear importer is overexpressed in diabetic kidney suggesting
13 that its expression is regulated by sugar (Kohler et al., 2001). Moreover, in Caco-2/TC7 cells
14 unmasking their basal pole, karyopherin alpha2 expression was recovered in cells that
15 exhibited GLUT2-triggered sugar signalling pathway (not shown). Karyopherin alpha2
16 imports proteins into the nucleus, and could trigger the ultimate steps of the effect of sugars
17 on the transcription machinery. The convergence of and the cross-talk between metabolic-
18 and GLUT2-pathways appeared to be an important way sugar-sensitive genes can be
19 regulated. Transcriptions factors involved in glucose sensing such as ChREBP (Uyeda et al.,
20 2002) and liver X receptor LXR (Mitro et al., 2007) are known to be translocated to the
21 nucleus in response to extracellular sugar (Helleboid-Chapman et al., 2006; Kawaguchi et al.,
22 2001). We speculate that if these transcription factors bind to the karyopherin alpha2 then it
23 will constitute the ultimate step of the GLUT2-triggered sugar signalling pathway.

24 GPCRs offer additional possibilities for sugar detection at the plasma membrane and for
25 sugar-dependent signal transduction. Indeed, GPCRs that bind fructose and sucrose have been
26 shown to transduce sugar signals in the tongue and several taste receptors have been identified
27 in an enteroendocrine cell line (Nelson et al., 2001; Rozengurt, 2006). Studies in knockout
28 mice support a role of T1R2/T1R3 heterodimers and T1R3 homodimers in sweet preferences
29 (Zhao et al., 2003). In enterocytes, the use of lactisole (an inhibitor of T1R3) suggests that
30 T1R-dependent signalling is involved in the regulation of some fructose-stimulated genes.
31 The signal transduction mechanisms used by these G protein-coupled sweet receptors are less
32 clear (Wettschureck and Offermanns, 2005). The $G\alpha$ Gustducin, mainly expressed in taste
33 cells, is believed to be able to couple receptors to phosphodiesterase resulting in a decrease of

1 cyclic nucleotide levels (Ozeck et al., 2004). However, studies also implicate that a receptor-
2 activated G_s can activate adenylyl cyclase to generate cAMP (Striem et al., 1989). Sugar
3 provision to enterocytes increases intracellular cAMP levels suggesting that the regulation of
4 SGLT1 (Dyer et al., 2003) and GLUT5 (Gouyon et al., 2003b; Mahraoui et al., 1994;
5 Mesonero et al., 1995) rely on adenylate cyclase/PKA pathways. Moreover, we report here
6 the presence of the T1R3 cognate $G\alpha$ Gustducin in Caco-2/TC7 cells. We speculate that
7 sweet taste receptor pathways contribute to fructose effects on GLUT5 and L-PK mRNA
8 accumulations, which were still detected in GLUT2-null mice.

9 Finally, we present here evidence that the sugar signal can be polarized in enterocytes.
10 Whereas the apical supply of fructose elicited stimulation of GLUT5 mRNA expression, the
11 basolateral supply of sugars generated a transcriptional response of GLUT2. *In vivo*, in
12 neonatal rat intestine, plasma fructose was shown to regulate the expression of GLUT2 (Cui
13 et al., 2003). However, the experimental design *in vivo* did not distinguish between direct
14 effects of plasma fructose from secondary signals. Our data in Caco-2/TC7 cells support the
15 idea that sugars can directly signal from the basolateral membrane. Thus detectors located at
16 the basolateral membrane trigger a polarized sugar signal. Here, we show that T1R2 and
17 T1R3 are both located in basolateral membranes of differentiated enterocytes. Moreover,
18 GLUT2 itself is the main sugar transporter in the basolateral membrane of enterocytes
19 (Cheeseman and O'Neill, 1998). The physiological role of the stimulation by basolateral
20 sugar provision of the expression of glucose transporters is unclear, however it could be a
21 thrifty mechanism well suited to an environment in which nutrient supplies are rare and need
22 to be fully absorbed. This mechanism however could become a threat for organisms facing a
23 nutritionally rich environment that would permanently stimulate sugar uptake capacities in the
24 intestine. In the diabetic state, hyperglycaemia is maintained during interprandial periods and
25 this sustains the transcription of sugar target genes via a basolateral signal and might
26 contribute to glucose toxicity in the long term. At the apical membrane, SGLT1 transports
27 glucose and galactose and GLUT5 transports fructose, but these transporters are saturated
28 rapidly as sugar concentrations rise. After a sugar-rich meal, GLUT2 transiently translocates
29 into the apical membrane and improves the transport capacity of the intestine (Kellett and
30 Brot-Laroche, 2005). Hence, GLUT2, depending on its relative abundance in the apical and
31 basolateral membranes, is likely to trigger sugar signals from the blood or the intestinal lumen
32 content. Whereas GLUT2 may be transiently expressed in the apical enterocyte membrane, it
33 is permanently high in the basolateral membrane, and may thereby determine the polarity of
34 sugar detection.

1 At least three sugar-signalling pathways have now been identified suggesting the capacity for
2 fine tuning of intestinal sugar delivery. The molecular identification of sugar-detectors
3 embedded in the enterocyte plasma membrane provides new therapeutic targets and strategies
4 toward controlling intestinal sugar delivery for the improved health of patients suffering
5 hyperglycemic episodes.

ACKNOWLEDGMENTS

1
2 We thank Pr J. Chambaz and the members of late UMRS505 INSERM-UPMC for constant
3 support and fruitful discussions. We are most grateful to B. Ballif for critical reading of the
4 manuscript and to L. Arnaud for encouragements and assistance. We acknowledge the
5 contribution of C. Lasne and N. Mokhatari-Comuce for animal care. We thank L. Caillaud
6 (IFR58) and K. Ahamed (IFR58) for assistance with the experiments, A.M. Faussat (IFR58)
7 for FACS sorting and C. Klein (IFR58) for confocal microscopy analysis. EBL and AL are
8 grateful to CNRS, INSERM and UPMC for support. MLG is an INSERM fellow and
9 recipient of a grant from the Benjamin Delessert Institute. VT and ES are recipients of the
10 French MRT fellowship.

BIBLIOGRAPHY

- 1
2 Blakemore SJ, Aledo JC, James J, Campbell FC, Lucocq JM, Hundal HS. 1995. The GLUT5
3 hexose transporter is also localized to the basolateral membrane of the human
4 jejunum. *Biochem J* 309 (Pt 1):7-12.
- 5 Burant CF, Flink S, DePaoli AM, Chen J, Lee WS, Hediger MA, Buse JB, Chang EB. 1994.
6 Small intestine hexose transport in experimental diabetes. Increased transporter
7 mRNA and protein expression in enterocytes. *J Clin Invest* 93(2):578-585.
- 8 Burant CF, Saxena M. 1994. Rapid reversible substrate regulation of fructose transporter
9 expression in rat small intestine and kidney. *Am J Physiol* 267(1 Pt 1):G71-79.
- 10 Cassany A, Guillemain G, Klein C, Dalet V, Brot-Laroche E, Leturque A. 2004. A
11 karyopherin alpha2 nuclear transport pathway is regulated by glucose in hepatic and
12 pancreatic cells. *Traffic* 5(1):10-19.
- 13 Chantret I, Rodolosse A, Barbat A, Dussaulx E, Brot-Laroche E, Zweibaum A, Rousset M.
14 1994. Differential expression of sucrase-isomaltase in clones isolated from early and
15 late passages of the cell line Caco-2: evidence for glucose-dependent negative
16 regulation. *J Cell Sci* 107 (Pt 1):213-225.
- 17 Cheeseman CI, O'Neill D. 1998. Basolateral D-glucose transport activity along the crypt-
18 villus axis in rat jejunum and upregulation induced by gastric inhibitory peptide and
19 glucagon-like peptide-2. *Exp Physiol* 83(5):605-616.
- 20 Cui XL, Jiang L, Ferraris RP. 2003. Regulation of rat intestinal GLUT2 mRNA abundance by
21 luminal and systemic factors. *Biochim Biophys Acta* 1612(2):178-185.
- 22 Cui XL, Soteropoulos P, Tolias P, Ferraris RP. 2004. Fructose-responsive genes in the small
23 intestine of neonatal rats. *Physiol Genomics* 18(2):206-217.
- 24 Delie F, Rubas W. 1997. A human colonic cell line sharing similarities with enterocytes as a
25 model to examine oral absorption: advantages and limitations of the Caco-2 model.
26 *Crit Rev Ther Drug Carrier Syst* 14(3):221-286.
- 27 Diez-Sampedro A, Hirayama BA, Osswald C, Gorboulev V, Baumgarten K, Volk C, Wright
28 EM, Koepsell H. 2003. A glucose sensor hiding in a family of transporters. *Proc Natl*
29 *Acad Sci U S A* 100(20):11753-11758.
- 30 Dyer J, Vayro S, King TP, Shirazi-Beechey SP. 2003. Glucose sensing in the intestinal
31 epithelium. *Eur J Biochem* 270(16):3377-3388.
- 32 Ferraris RP, Casirola DM, Vinnakota RR. 1993. Dietary carbohydrate enhances intestinal
33 sugar transport in diabetic mice. *Diabetes* 42(11):1579-1587.
- 34 Girard J, Ferre P, Foufelle F. 1997. Mechanisms by which carbohydrates regulate expression
35 of genes for glycolytic and lipogenic enzymes. *Annu Rev Nutr* 17:325-352.
- 36 Goda T. 2000. Regulation of the expression of carbohydrate digestion/absorption-related
37 genes. *Br J Nutr* 84 Suppl 2:S245-248.
- 38 Gould GW, Thomas HM, Jess TJ, Bell GI. 1991. Expression of human glucose transporters in
39 *Xenopus* oocytes: kinetic characterization and substrate specificities of the
40 erythrocyte, liver, and brain isoforms. *Biochemistry* 30(21):5139-5145.
- 41 Gouyon F, Caillaud L, Carriere V, Klein C, Dalet V, Citadelle D, Kellett GL, Thorens B,
42 Leturque A, Brot-Laroche E. 2003a. Simple-sugar meals target GLUT2 at enterocyte
43 apical membranes to improve sugar absorption: a study in GLUT2-null mice. *J*
44 *Physiol* 552(Pt 3):823-832.
- 45 Gouyon F, Onesto C, Dalet V, Pages G, Leturque A, Brot-Laroche E. 2003b. Fructose
46 modulates GLUT5 mRNA stability in differentiated Caco-2 cells: role of cAMP-
47 signalling pathway and PABP (polyadenylated-binding protein)-interacting protein
48 (Paip) 2. *Biochem J* 375(Pt 1):167-174.

- 1 Grasset E, Pinto M, Dussaulx E, Zweibaum A, Desjeux JF. 1984. Epithelial properties of
2 human colonic carcinoma cell line Caco-2: electrical parameters. *Am J Physiol* 247(3
3 Pt 1):C260-267.
- 4 Guillam MT, Hummler E, Schaerer E, Yeh JI, Birnbaum MJ, Beermann F, Schmidt A, Deriaz
5 N, Thorens B. 1997. Early diabetes and abnormal postnatal pancreatic islet
6 development in mice lacking Glut-2. *Nat Genet* 17(3):327-330.
- 7 Guillemain G, Loizeau M, Pincon-Raymond M, Girard J, Leturque A. 2000. The large
8 intracytoplasmic loop of the glucose transporter GLUT2 is involved in glucose
9 signaling in hepatic cells. *J Cell Sci* 113 (Pt 5):841-847.
- 10 Guillemain G, Munoz-Alonso MJ, Cassany A, Loizeau M, Faussat AM, Burnol AF, Leturque
11 A. 2002. Karyopherin alpha2: a control step of glucose-sensitive gene expression in
12 hepatic cells. *Biochem J* 364(Pt 1):201-209.
- 13 Helleboid-Chapman A, Helleboid S, Jakel H, Timmerman C, Sergheraert C, Pattou F,
14 Fruchart-Najib J, Fruchart JC. 2006. Glucose regulates LXRAalpha subcellular
15 localization and function in rat pancreatic beta-cells. *Cell Res* 16(7):661-670.
- 16 Holsbeeks I, Lagatie O, Van Nuland A, Van de Velde S, Thevelein JM. 2004. The eukaryotic
17 plasma membrane as a nutrient-sensing device. *Trends Biochem Sci* 29(10):556-564.
- 18 Im SS, Kang SY, Kim SY, Kim HI, Kim JW, Kim KS, Ahn YH. 2005. Glucose-stimulated
19 upregulation of GLUT2 gene is mediated by sterol response element-binding protein-
20 1c in the hepatocytes. *Diabetes* 54(6):1684-1691.
- 21 Inukai K, Asano T, Katagiri H, Ishihara H, Anai M, Fukushima Y, Tsukuda K, Kikuchi M,
22 Yazaki Y, Oka Y. 1993. Cloning and increased expression with fructose feeding of rat
23 jejunal GLUT5. *Endocrinology* 133(5):2009-2014.
- 24 Jacob F, Monod J. 1961. Genetic regulatory mechanisms in the synthesis of proteins. *J Mol*
25 *Biol* 3:318-356.
- 26 Johnston M. 1999. Feasting, fasting and fermenting. Glucose sensing in yeast and other cells.
27 *Trends Genet* 15(1):29-33.
- 28 Kawaguchi T, Takenoshita M, Kabashima T, Uyeda K. 2001. Glucose and cAMP regulate the
29 L-type pyruvate kinase gene by phosphorylation/dephosphorylation of the
30 carbohydrate response element binding protein. *Proc Natl Acad Sci U S A*
31 98(24):13710-13715.
- 32 Kellett GL, Brot-Laroche E. 2005. Apical GLUT2: a major pathway of intestinal sugar
33 absorption. *Diabetes* 54(10):3056-3062.
- 34 Kohler M, Buchwalow IB, Alexander G, Christiansen M, Shagdarsuren E, Samoilova V,
35 Hartmann E, Mervaala EM, Haller H. 2001. Increased importin alpha protein
36 expression in diabetic nephropathy. *Kidney Int* 60(6):2263-2273.
- 37 Levin RJ. 1994. Digestion and absorption of carbohydrates--from molecules and membranes
38 to humans. *Am J Clin Nutr* 59(3 Suppl):690S-698S.
- 39 Mahraoui L, Takeda J, Mesonero J, Chantret I, Dussaulx E, Bell GI, Brot-Laroche E. 1994.
40 Regulation of expression of the human fructose transporter (GLUT5) by cyclic AMP.
41 *Biochem J* 301 (Pt 1):169-175.
- 42 Massillon D. 2001. Regulation of the glucose-6-phosphatase gene by glucose occurs by
43 transcriptional and post-transcriptional mechanisms. Differential effect of glucose and
44 xylitol. *J Biol Chem* 276(6):4055-4062.
- 45 Mesonero J, Matosin M, Cambier D, Rodriguez-Yoldi MJ, Brot-Laroche E. 1995. Sugar-
46 dependent expression of the fructose transporter GLUT5 in Caco-2 cells. *Biochem J*
47 312 (Pt 3):757-762.
- 48 Mitro N, Mak PA, Vargas L, Godio C, Hampton E, Molteni V, Kreuzsch A, Saez E. 2007. The
49 nuclear receptor LXR is a glucose sensor. *Nature* 445(7124):219-223.

- 1 Miyamoto K, Hase K, Takagi T, Fujii T, Taketani Y, Minami H, Oka T, Nakabou Y. 1993.
2 Differential responses of intestinal glucose transporter mRNA transcripts to levels of
3 dietary sugars. *Biochem J* 295 (Pt 1):211-215.
- 4 Nelson G, Hoon MA, Chandrashekar J, Zhang Y, Ryba NJ, Zuker CS. 2001. Mammalian
5 sweet taste receptors. *Cell* 106(3):381-390.
- 6 Nishimura M, Uyeda K. 1995. Purification and characterization of a novel xylulose 5-
7 phosphate-activated protein phosphatase catalyzing dephosphorylation of fructose-6-
8 phosphate,2-kinase:fructose-2,6-bisphosphatase. *J Biol Chem* 270(44):26341-26346.
- 9 Ozeck M, Brust P, Xu H, Servant G. 2004. Receptors for bitter, sweet and umami taste couple
10 to inhibitory G protein signaling pathways. *Eur J Pharmacol* 489(3):139-149.
- 11 Rodolosse A, Carriere V, Chantret I, Lacasa M, Zweibaum A, Rousset M. 1997. Glucose-
12 dependent transcriptional regulation of the human sucrase-isomaltase (SI) gene.
13 *Biochimie* 79(2-3):119-123.
- 14 Rodolosse A, Chantret I, Lacasa M, Chevalier G, Zweibaum A, Swallow D, Rousset M. 1996.
15 A limited upstream region of the human sucrase-isomaltase gene confers glucose-
16 regulated expression on a heterologous gene. *Biochem J* 315 (Pt 1):301-306.
- 17 Rozengurt E. 2006. Taste receptors in the gastrointestinal tract. I. Bitter taste receptors and
18 alpha-gustducin in the mammalian gut. *Am J Physiol Gastrointest Liver Physiol*
19 291(2):G171-177.
- 20 Rozengurt N, Wu S, Chen MC, Huang C, Sternini C, Rozengurt E. 2006. Co-localization of
21 the {alpha} subunit of gustducin with PYY and GLP-1 in L cells of human colon. *Am*
22 *J Physiol Gastrointest Liver Physiol*.
- 23 Schuit F, Flamez D, De Vos A, Pipeleers D. 2002. Glucose-regulated gene expression
24 maintaining the glucose-responsive state of beta-cells. *Diabetes* 51 Suppl 3:S326-332.
- 25 Striem BJ, Pace U, Zehavi U, Naim M, Lancet D. 1989. Sweet tastants stimulate adenylate
26 cyclase coupled to GTP-binding protein in rat tongue membranes. *Biochem J*
27 260(1):121-126.
- 28 Stumpel F, Burcelin R, Jungermann K, Thorens B. 2001. Normal kinetics of intestinal
29 glucose absorption in the absence of GLUT2: evidence for a transport pathway
30 requiring glucose phosphorylation and transfer into the endoplasmic reticulum. *Proc*
31 *Natl Acad Sci U S A* 98(20):11330-11335.
- 32 Towle HC. 2005. Glucose as a regulator of eukaryotic gene transcription. *Trends Endocrinol*
33 *Metab* 16(10):489-494.
- 34 Uyeda K, Yamashita H, Kawaguchi T. 2002. Carbohydrate responsive element-binding
35 protein (ChREBP): a key regulator of glucose metabolism and fat storage. *Biochem*
36 *Pharmacol* 63(12):2075-2080.
- 37 Vaulont S, Vasseur-Cognet M, Kahn A. 2000. Glucose regulation of gene transcription. *J Biol*
38 *Chem* 275(41):31555-31558.
- 39 Villadsen D, Smith SM. 2004. Identification of more than 200 glucose-responsive
40 *Arabidopsis* genes none of which responds to 3-O-methylglucose or 6-deoxyglucose.
41 *Plant Mol Biol* 55(4):467-477.
- 42 Wang H, Kouri G, Wollheim CB. 2005. ER stress and SREBP-1 activation are implicated in
43 beta-cell glucolipotoxicity. *J Cell Sci* 118(Pt 17):3905-3915.
- 44 Wang H, Maechler P, Antinozzi PA, Herrero L, Hagenfeldt-Johansson KA, Bjorklund A,
45 Wollheim CB. 2003. The transcription factor SREBP-1c is instrumental in the
46 development of beta-cell dysfunction. *J Biol Chem* 278(19):16622-16629.
- 47 Wettschureck N, Offermanns S. 2005. Mammalian G proteins and their cell type specific
48 functions. *Physiol Rev* 85(4):1159-1204.

- 1 Xu H, Staszewski L, Tang H, Adler E, Zoller M, Li X. 2004. Different functional roles of
2 T1R subunits in the heteromeric taste receptors. Proc Natl Acad Sci U S A
3 101(39):14258-14263.
- 4 Zhao GQ, Zhang Y, Hoon MA, Chandrashekar J, Erlenbach I, Ryba NJ, Zuker CS. 2003. The
5 receptors for mammalian sweet and umami taste. Cell 115(3):255-266.
6

FIGURE LEGENDS

Figure 1: Dietary sugars regulate intestinal genes in mice

Messenger RNA samples were from jejunal mucosa of wild-type (white bars) or GLUT2-null (black bars) mice fed a protein-, glucose- or fructose-rich diet for 5 days. Northern blots (mSGLT1, mGLUT5) and RT-PCR (mL-PK, mSI) were performed. Results represent the average \pm SEM from at least 6 mice and are presented as the ratio of mRNA levels in sugar over low carbohydrate (protein) dietary conditions. mRNA levels under a protein diet were set at 1 (dotted line). (*) Statistical significance ($P < 0.05$) comparing sugar- to protein-fed mice; (#) Statistical significance ($P < 0.05$) comparing wild-type and GLUT2 null mice; (NS) Non significant.

Figure 2: Polarized sugar supply affects the expression of sugar sensitive genes in human enterocytes

Caco-2/TC7 cells stably expressing either hGLUT5-, hGLUT2- or hSI-promoter luciferase constructs, were cultured on either solid or porous support for 17 days. After differentiation of the cells, culture media were without sugar addition (white bars) or supplemented with either 25 mmole.L⁻¹ glucose (G, black bars), 25 mmole.L⁻¹ fructose (F, hatched bars), 12.5 mmole.L⁻¹ sucrose (S, large hatched bars), or a 12.5 mmole.L⁻¹ equimolar, glucose and fructose mix (FG, grey hatched bars). Sugars were added to the apical and/or basal compartments for 4 days as indicated. Luciferase activities in RLU.sec⁻¹. μ g⁻¹ protein were normalized to basal as measured in cells cultured in media without sugar addition for 4 days. (*) Statistical significance ($P < 0.05$) compared to low sugar conditions; (NS) Non significant; (NA) Non-applicable for cells cultured on solid support.

Figure 3: Basolateral sugar sensing regulates hGLUT2 promoter activity

(A) Time course of sugar transfer across Caco-2/TC7 cells relative to the compartment of sugar application (left panel apical supply, right panel basolateral supply). Apical sugar is denoted by filled diamonds, basolateral sugar is denoted by open squares. Sugar decrease in the compartment of origin is denoted by dotted lines while sugar appearance in the opposite compartment is indicated by a solid line. Sugar concentrations reached equilibrium within 12 hours. For panels B and C, Caco-2/TC7 cells stably expressing an hGLUT2-promoter luciferase construct were used. (B, left panel) Time course of hGLUT2-promoter activity in cells apically and basolaterally provided with 25 mmole.L⁻¹ glucose (filled squares; G25) or

1 fructose (filled circles, F25) after having been cultured for 4 days without sugar addition
 2 (Low sugar (LS) is denoted by open diamonds). (B, right panel) Time course of hGLUT2-
 3 promoter activity in cells deprived of glucose in apical and basolateral compartments. (C)
 4 hGLUT2-promoter activity in cells after glucose-deprivation either in the apical, basal or
 5 apical and basal compartments for 12 hours. Note that the promoter is sensitive to basally
 6 applied glucose. Luciferase activities in $\text{RLU}\cdot\text{sec}^{-1}\cdot\mu\text{g}^{-1}$ protein were normalized to basal
 7 amounts measured in cells cultured in media without sugar addition for 4 days (B, left panel)
 8 or in cells cultured in media containing glucose (B, right panel and C). (*) Statistical
 9 significance ($P < 0.05$) comparing to $t=0\text{h}$; (NS) Non significant.

10

11 ***Figure 4: Metabolic pathways control sugar-sensitive gene expression***

12 (A) Differentiated Caco-2/TC7 cells stably expressing the hGLUT2-promoter luciferase
 13 construct were cultured on porous support, either without sugar addition (white bars) or with
 14 $25\text{ mmole}\cdot\text{L}^{-1}$ glucose (black bars) or with $25\text{ mmole}\cdot\text{L}^{-1}$ 3-O-Methylglucose (3OMG, hatched
 15 bar) for 48 hours. Luciferase activities $\text{RLU}\cdot\text{sec}^{-1}\cdot\mu\text{g}^{-1}$ protein were normalized to values
 16 without glucose addition and are represented as the mean \pm SEM ($n=6$). (B) RT-PCR
 17 analysis of hADRP, hGLUT2, hL-PK and hG-6-Pase mRNA in differentiated Caco-2/TC7
 18 cells cultured on porous support in the presence or absence of $25\text{ mmole}\cdot\text{L}^{-1}$ glucose and/or
 19 $250\text{ mmole}\cdot\text{L}^{-1}$ N-acetylglucosamine (N-AGA) for 48 hours. hADRP/L19, hGLUT2/L19, hL-
 20 PK/L19 and hG-6-Pase/L19 ratios were normalized to the control level in cells cultured
 21 without added sugar. Results are representative of three independent experiments and are
 22 represented as means \pm SEM ($n=3$). (*) Statistical significance ($P < 0.05$) compared to
 23 control; (#) Statistical significance ($P < 0.05$) compared N-AGA treated cells; (NS) Non
 24 significant.

25 ***Figure 5: GLUT2 is a sugar detector controlling gene expression***

26 (A) Upper panel: FACS profiles of Caco-2/TC7 cells untransfected (parental) and stably
 27 expressing EGFP, EGFP-GLUT2-Cterminus or EGFP-GLUT2-loop constructs. Lower panel:
 28 Epifluorescence of EGFP, EGFP-GLUT2-Cterminus or EGFP-GLUT2-loop proteins in Caco-
 29 2/TC7 transfected cells. Toto-3 is a nuclear marker. Endogenous hGLUT2 (B) and hL-PK (C)
 30 mRNA expression is shown from control cells (untransfected or EGFP expressing cells) or
 31 from cells expressing EGFP-GLUT2 domains, cultured on porous support in glucose (black
 32 bars, B and C) or fructose (hatched bars, B) for 4 days. hGLUT2/L19 and hL-PK/L19 mRNA

1 ratios were normalized to control levels reached in cells in media without sugar addition
2 (dotted line). (D) Caco-2/TC7 cells stably expressing EGFP-GLUT2 domains were
3 secondarily and transiently transfected with the hGLUT2 promoter luciferase construct.
4 Luciferase activities ($\text{RLU}\cdot\text{sec}^{-1}\cdot\mu\text{g}^{-1}$ protein) measured in 4 days fructose stimulated cells
5 were normalized to conditions without sugar addition (dotted line). (E) Endogenous human
6 GLUT2 mRNA abundance in the presence of $250\text{ mmole}\cdot\text{L}^{-1}$ N-acetylglucosamine (N-AGA)
7 in control or in EGFP-GLUT2-loop Caco-2/TC7 cells for 48 hours. hGLUT2/L19 mRNA
8 ratios were normalized to mRNA levels in cells without sugar addition. Data were expressed
9 as means \pm SEM (n=6). (*) Statistical significance ($P < 0.05$) compared to control
10 conditions; (#) statistical significance ($P < 0.05$) comparing control and EGFP-GLUT2-loop
11 cells; (NS) non significant.

12 **Figure 6: Identification of GPCR-mediated sugar signalling in enterocytes**

13 (A) Human GLUT2, GLUT5, L-PK and SGLT1 mRNA abundance was measured in Caco-
14 2/TC7 cells using quantitative RT-PCR. Fully differentiated cells were maintained for 2 days
15 in sugar free medium (white bars), before challenge with $25\text{ mmole}\cdot\text{L}^{-1}$ fructose (F, hatched
16 bars) or $25\text{ mmole}\cdot\text{L}^{-1}$ fructose and $1\text{ mmole}\cdot\text{L}^{-1}$ lactisole (F+Lact, grey and hatched bars) for
17 48 hours. Messengers/L19 mRNA ratios were normalized to basal levels in cells in media
18 without sugar addition. Data were expressed as the mean \pm SEM of 2 independent
19 experiments. (*) Statistical significance ($P < 0.05$) compared to control conditions; (#)
20 Statistical significance ($P < 0.01$) comparing F and F+Lact; (NS) non significant. (B) Semi-
21 quantitative RT-PCR of T1R3 sweet taste receptor and $G\alpha_{\text{gust}}$ in differentiated Caco-2/TC7
22 cells cultured without sugar addition (Low Sugar, LS) or in media supplemented with 25
23 $\text{mmole}\cdot\text{L}^{-1}$ fructose (F), $25\text{ mmole}\cdot\text{L}^{-1}$ glucose (G) for 4 days. (C) Confocal microscopy
24 analysis of filter-grown differentiated Caco-2/TC7 cells labelled with T1R2 and T1R3
25 antibodies. Serial Z sections of Caco-2/TC7 cells displayed similar pattern of basolateral
26 location for T1R2 and T1R3 receptors. White bars represent $10\mu\text{m}$.

27
28
29
30
31