



**HAL**  
open science

## Comparison of viro-immunological marker changes between HIV-1 and HIV-2-infected patients in France.

Julia Drylewicz, Sophie Matheron, Estibaliz Lazaro, Florence Damond, Fabrice Bonnet, François Simon, François Dabis, Françoise Brun-Vézinet, Geneviève Chêne, Rodolphe Thiébaud

### ► To cite this version:

Julia Drylewicz, Sophie Matheron, Estibaliz Lazaro, Florence Damond, Fabrice Bonnet, et al.. Comparison of viro-immunological marker changes between HIV-1 and HIV-2-infected patients in France.: Comparison between HIV-1 and HIV-2 in France. *AIDS*, Lippincott, Williams & Wilkins, 2008, 22 (4), pp.457-68. 10.1097/QAD.0b013e3282f4ddfc . inserm-00258131

**HAL Id: inserm-00258131**

**<https://www.hal.inserm.fr/inserm-00258131>**

Submitted on 6 Jul 2009

**HAL** is a multi-disciplinary open access archive for the deposit and dissemination of scientific research documents, whether they are published or not. The documents may come from teaching and research institutions in France or abroad, or from public or private research centers.

L'archive ouverte pluridisciplinaire **HAL**, est destinée au dépôt et à la diffusion de documents scientifiques de niveau recherche, publiés ou non, émanant des établissements d'enseignement et de recherche français ou étrangers, des laboratoires publics ou privés.

Editorial Manager(tm) for AIDS  
Manuscript Draft

Manuscript Number: AIDS-D-07-00945R1

Title: Comparison of Viro-Immunologic Markers Changes between HIV-1 and HIV-2 Infected Patients in France from the ANRS CO 3 and CO 5 cohorts

Article Type: Original paper

Section/Category:

Keywords: HIV-1, HIV-2, CD4, CD8, HIV viral load, Longitudinal study

Corresponding Author: Dr. Rodolphe THIÉBAUT, MD, PhD

Corresponding Author's Institution: INSERM U875 & U593

First Author: Julia Drylewicz, MSc

Order of Authors: Julia Drylewicz, MSc; Sophie Matheron, MD, PhD; Estibaliz Lazaro, MD; Florence Damond, MD, PhD; Fabrice Bonnet, MD, PhD; François Simon, MD, PhD; François Dabis, MD, PhD; Françoise Brun-Vezinet, MD, PhD; Geneviève Chêne, MD, PhD; Rodolphe THIÉBAUT, MD, PhD

Manuscript Region of Origin:

Suggested Reviewers:

## Abstract

**Background:** HIV-2 is known to be less pathogenic than HIV-1, although the underlying mechanisms are still debated. We compared the changes over time in viro-immunologic markers in HIV-1 and HIV-2 infected patients living in France during natural history and after initiation of the first Combination of AntiRetroviral Treatment (CART).

**Method:** Patients were included in the ANRS CO3 HIV-1 cohort (N=6707) or the ANRS CO5 HIV-2 cohort (N=572). HIV-1 infected patients were matched to HIV-2 patients according to sex, age, HIV transmission group and period of treatment initiation. Changes in markers have been estimated with linear mixed models.

**Results:** Analyses were performed for three groups of patients: (1) those with estimated date of contamination (98 HIV-1 and 49 HIV-2 seroincident patients), (2) untreated seroprevalent patients (320 HIV-1 and 160 HIV-2) and (3) those who initiated a first CART (59 HIV-1 and 63 HIV-2). In group 1, CD4 T-cell decreased less rapidly in HIV-2 than HIV-1 patients (-9 vs.-49 cells/mm<sup>3</sup>/year,  $p<10^{-4}$ ). Estimated slopes in untreated group 2 were similar to those estimated in group 1 (-11 vs.-49 cells/mm<sup>3</sup>/year,  $p=0.003$ ). In group 3, baseline CD4 at CART initiation was not different according to the type of infection (269 vs.220 cells/mm<sup>3</sup>). During the first two months of treatment, CD4 count increased by +59 cells/mm<sup>3</sup>/month (95% Confidence Interval [CI]=34;84) for HIV-1 and +24 (CI=6;42) for HIV-2. The plasma viral load drop was 3-fold more important in HIV-1 patients: -1.56 log<sub>10</sub>/ml/month (CI=-1.83;-1.30) vs. -0.62 (CI=-0.84;-0.40) among HIV-2 patients ( $p<10^{-4}$ ).

**Conclusion:** Differences between the two infections during natural history are similar to those previously described in Africa. Paradoxically, once treatment is started, response is poorer in HIV-2 patients than in HIV-1 patients.

**Keywords** HIV-1, HIV-2, CD4, CD8, HIV viral load, Longitudinal study

## **Comparison of Viro-Immunologic Markers Changes between HIV-1 and HIV-2 Infected Patients in France from the ANRS CO 3 and CO 5 cohorts.**

**Short title:** Comparison between HIV-1 and HIV-2 in France

**Authors:** Julia Drylewicz<sup>1,2</sup>, Sophie Matheron<sup>3</sup>, Estibaliz Lazaro<sup>4</sup>, Florence Damond<sup>3</sup>, Fabrice Bonnet<sup>4</sup>, François Simon<sup>5</sup>, François Dabis<sup>1,2</sup>, Françoise Brun-Vezinet<sup>3</sup>, Geneviève Chêne<sup>1,2</sup>, Rodolphe Thiébaud<sup>1,2</sup> and ANRS CO5 HIV-2 study cohort and ANRS CO3 Aquitaine cohort study groups

**Affiliations:**

1. INSERM, U875 & U593, Bordeaux, F-33076, France
2. Bordeaux 2 University, ISPED, Bordeaux, F-33076, France
3. Bichat Claude Bernard Hospital, Assistance Publique des Hôpitaux de Paris, Paris, F-75018, France
4. Bordeaux University Hospital, Bordeaux, F-33000, France
5. Charles Nicolle Hospital, Rouen, F-76000, France

**Corresponding authors:**

Rodolphe Thiébaud

INSERM U875 & U593

ISPED – Université Bordeaux 2

146 Rue Léo Saignat 33076 BORDEAUX, France

Tel : (33) 5 57 57 45 21 Fax : (33) 5 56 24 00 81

Email : [rodolphe.thiebaud@isped.u-bordeaux2.fr](mailto:rodolphe.thiebaud@isped.u-bordeaux2.fr)

**Word count:** 275 (abstract), 3213 (text), 63 references, 4 tables, 2 figures

**Acknowledgments:**

We gratefully thank Delali Sefe for editing the English.

## Introduction

Human Immunodeficiency Virus type 2 (HIV-2) is endemic in West Africa and sporadic in the rest of the world [1-4]. Compared to individuals infected by HIV-1, those infected by HIV-2 have a slower clinical progression [5], a lower mortality rate for patients with high CD4+ T lymphocyte count (CD4) [6-7] and lower rates of transmission [8-11]. In West African countries, comparisons have shown a slower CD4 depletion [11-13] and a lower plasma viral load in HIV-2 infected patients [13-20]. At AIDS stage, HIV-2 infected patients tend to have higher CD4 count [15] and clinical manifestations may differ between the two infections [16,17]. Several hypotheses have been raised to explain the differences between the two infections: lower virulence of HIV-2 [21,22], lower replication capacity of HIV-2 [23-27], better immune control [28-33] and lower activation of immune system during HIV-2 infection [12,23,28,34]. These factors might be associated as cell activation is linked to viral load [34]. However, cell responsiveness to activation might also vary [5,35].

All reported differences in the rate of disease progression between the two infections are from cohort studies performed in sub-Saharan Africa. No direct comparison has been made in Europe or the United States during the course of the infection, although the environment may play a role in the difference of pathogenicity between the two infections. For instance, the level of lymphocyte activation is higher in Africa than in Europe [36], consequently the role of activation in the difference between the two infections may be weaker or even reinforced because of different background rates. Therefore, we hypothesized that the differences in viro-immunologic markers levels and evolution could be different in Europe compared to sub-Saharan Africa.

Here, we report the changes in plasma HIV RNA, CD4 and CD8 over time in the French national cohort of HIV-2 infected adult patients compared to individually matched HIV-1 infected patients from the French Aquitaine Cohort.

## Methods

### HIV cohorts

Data is taken from the ANRS CO5 HIV-2 cohort [37] and the ANRS CO3 Aquitaine cohort [38]. The ANRS CO5 HIV-2 cohort is an ongoing national prospective study initiated in 1994 in 111 clinical centres in France. Inclusion criteria to the cohort are HIV-2 infection only, age  $\geq 18$  years, residence in France planned for at least 1 year and informed consent available. The ANRS CO3 Aquitaine cohort is an ongoing prospective study initiated in 1987.

Inclusion criteria are HIV-1 infection in patients aged over 18 years, and informed consent available. In the two cohorts, clinical, epidemiological and therapeutic data are collected by standardized questionnaires at each visit to the hospital (every 3 to 6 months according to clinical, immuno-virological and therapeutic status).

### **Markers quantifications**

CD4 count was performed by flow cytometry in the two cohorts. Plasma HIV-1 RNA was quantified mainly by branched DNA assays (Chiron Quantiplex RNA HIV-1, Emeryville, CA, USA) with detection limits of 2.7 log<sub>10</sub> copies/ml (500 copies/ml) or 1.7 log<sub>10</sub> copies/ml (50 copies/ml). Although there is one commercial kit, designed for HIV-1, which can also quantify only HIV-2 subtype A RNA, there is no commercial assay specifically designed for HIV2 viral load [39]. Plasma HIV-2 RNA quantification was performed using HIV-2 strain NIHZ as a standard (Advanced Biotechnology Incorporated, Maryland, USA) with lower detection limits of either 2.4 log<sub>10</sub> copies/ml (250 copies/ml) [39] or 2.0 log<sub>10</sub> copies/ml (100 copies/ml) [41].

### **Study populations**

We defined three study populations in each cohort: (1) seroincident patients, (2) seroprevalent and (3) naive patients starting a Combined Anti Retroviral Treatment (CART: combination of 2 nucleoside inhibitors and 1 protease inhibitor or 3 nucleoside inhibitors). The seroincident group included all seropositive patients whose date of seroconversion was known or well estimated, based on the period between the last negative and the first positive antibody test of less than 3 years. This population was defined retrospectively according to the availability of negative serology in the patients already included in each cohort. Data was collected from date of seroconversion and censored after 3 years of follow-up to avoid any informative dropout [42]. In this group, no patient started an antiretroviral treatment or died before the censoring date. The seroprevalent group included all seropositive untreated patients, and without documented date of HIV infection. Data was collected from inclusion and was censored if patient started an antiretroviral treatment or died. The last group included all HIV antiretroviral-naive patients who started an antiretroviral therapy consisting of at least 3 antiretroviral drugs. Data was collected from the date of first CART regimen initiation and was censored if the antiretroviral treatment was modified or if the patient died. An intent-to-continue analysis was also performed and results were similar (data not shown).

HIV-1 infected patients were individually matched to HIV-2 infected patients according to factors known to be associated with HIV-1 disease progression [38,42-44]. We considered: sex, HIV transmission group (in 4 categories: heterosexual, homosexual, blood recipients and other), period of treatment initiation (in 2 categories: 1996-2000 and 2001-2005 according to generation of available treatment) and age (in 4 categories:  $\leq 30$  years, 31-40, 41-50 and  $>50$  years) at seroconversion (for group 1), at cohort inclusion (for group 2) and at first CART regimen initiation (for group 3). For each HIV-2 infected patient, one (for group 3) to two (for group 1 and 2) HIV-1 infected patients were randomly selected for matching among eligible candidates. We selected only one HIV-1 patient for each HIV-2 patient in group 3 because of the restricted number of available patients. All HIV-1 infected patients who were prescribed a non-nucleoside inhibitor in their CART regimen were excluded, this class of antiretroviral drugs being not active against HIV-2 infection [45].

We carried out two sub-analyses to account for additional factors. For group 2, in addition to sex, age and HIV transmission group, we constituted a new study population by matching for country of birth (West Africa, Europe and others). This sub-analysis was not feasible in group 1 and 3 because of the restricted number of available patients. In group 3, we performed an additional match according to the baseline plasma viral load at treatment initiation ( $>3.5$  vs.  $\leq 3.5$   $\log_{10}$  copies/ml).

### **Statistical analysis**

Changes in biological markers were studied using piecewise linear mixed models. The baseline ( $t = 0$ ) was the date of seroconversion for group 1, the date of inclusion for group 2 and the date of first CART regimen initiation for group 3. Trends in the evolution of markers were fitted using one slope (in unit/year) for the first two groups. For the last group of treated patients, two slopes were considered: one for the early change (in unit/month) and a second for the long-term trend (in unit/year). The time taken for the slope to change ( $t = 2$  months) was determined for all patients by a likelihood profile. The correlation between individual baseline value(s) and the subsequent slope(s) was handled through the unstructured covariance matrix of random effects. The left-censoring of plasma viral load due to undetectable values was taken into account using a maximum likelihood method as previously described [46]. Adjustment for the type of assay used to quantify viral load did not modify the estimates of the slopes (data not shown). Data analyses were conducted with SAS<sup>®</sup> 8.1 (SAS Institute, Cary, NC, USA).

## Results

### Study population

In January 2006, the ANRS CO5 HIV-2 cohort had recruited 572 patients. Of these, 89 were seroincident patients, of whom 49 were antiretroviral-naïve at inclusion in the cohort (group 1). Among the 483 seroprevalent patients, 160 had no history of any antiretroviral treatment (group 2). Of the 572 patients, 105 started an antiretroviral regimen, of whom 63 received CART (group 3). By January 2006, 6 707 HIV-1 infected patients have been recruited into the ANRS CO3 Aquitaine cohort. The date of seroconversion was well estimated for 1 464 patients including 962 who were antiretroviral-naïve. Among the entire cohort, 1 036 patients started the CART regimen without any previous exposure to antiretroviral. A total of 98 HIV-1 and 49 HIV-2 seroincident patients, 320 HIV-1 and 160 HIV-2 seroprevalent patients and 59 HIV-1 and 63 HIV-2 CART treated patients were included (Figure 1). Study populations are described in Table 1.

### Seroincident patients

Median delay between seroconversion and first available laboratory measure was significantly shorter for HIV-1 infected patients than for HIV-2 (Table 2): 4.1 years vs. 6.8 years ( $p < 10^{-4}$ ). Without administrative censoring, the median follow-up was 36 months and 81 months for HIV-1 and HIV-2 respectively. During the first three years of follow-up, a median of four biological measurements per patient were available among HIV-1 and HIV-2 patients. The proportion of undetectable viral load measures were 14% and 85% in HIV-1 and HIV-2 infected patients, respectively. At enrolment in the cohort, median viral load was 4.11  $\log_{10}$  copies/ml for HIV-1 and 2.09  $\log_{10}$  copies/ml for HIV-2 and median CD4 count was 399 cells/mm<sup>3</sup> and 585 cells/mm<sup>3</sup> for HIV-1 and HIV-2 respectively.

Mean slopes estimated using linear mixed models were as show in table 3. CD4 count and CD4 percentage significantly decreased in the HIV-1 group (-49 cells/mm<sup>3</sup>/year and -1.01%/year) but was quite stable in HIV-2 group (-9 cells/mm<sup>3</sup>/year and -0.04%/year). On average, plasma viral load was quite stable over time in the HIV-1 group and in HIV-2 (-0.02 and +0.06  $\log_{10}$  copies/ml/year, respectively). CD8 count did not change significantly in both groups (Table 2). Hence, the CD4:CD8 ratio decreased significantly in HIV-1 group (-0.06/year) whereas it did not change in HIV-2 group (0.02/year,  $p < 10^{-4}$ ).



### **Seroprevalent patients**

Median delay between inclusion into the study and the first measurement of CD4 count was 2 months for HIV-1 and 6 months for HIV-2. During the follow-up (median of 4.9 years for HIV-1 and 2.9 for HIV-2), a median of four measurements were available for HIV-1 and seven for HIV-2. At inclusion into the study, the proportion of patients with undetectable plasma viral load was 9% and 39% for HIV-1 and HIV-2 respectively.

At enrolment into the cohorts, median CD4 count was significantly lower in HIV-2 than in HIV-1 patients: 260 cells/mm<sup>3</sup> vs. 324 (p=0.007), probably reflecting a later enrolment of HIV-2 patients compared to HIV-1 patients. However, median plasma viral load was still significantly lower in HIV-2 (2.62 vs. 4.39 log<sub>10</sub> copies/ml, p<10<sup>-4</sup>) as well as median CD8 count (p=0.0005).

The estimated average decrease in CD4 was 4.5-fold more pronounced in HIV-1: -49 cells/mm<sup>3</sup>/year than HIV-2: -11, (p=0.003, table 3). The CD4 percentage decrease was not significantly different between the two groups (p=0.70). There was a small insubstantial increase in plasma viral load in the two groups: +0.20 log<sub>10</sub> copies/ml/year for HIV-1 and +0.14 for HIV-2. Therefore, plasma viral load was still very different in the two populations after one year of follow-up (difference of 1 log<sub>10</sub> copies/ml, p=0.005). The increase in CD8 count did not differ between the two groups (p=0.44). The CD4:CD8 ratio decreased over time in the two groups, but it was more pronounced in the HIV-1 group: -0.06/year vs. -0.02 (p=10<sup>-4</sup>). We performed a second match including country of birth as a matching variable and results were similar. In addition, we looked at any modification of the effect of the type of infection (HIV-1 or HIV-2) on the slopes of each marker according to the country of birth and none were significant.

### **Patients starting CART regimen**

At the initiation of CART, the observed median CD4 count was not significantly different in the two groups (table 2, p=0.06), as well as CD4 percentage (p=0.70). Plasma viral load was significantly higher in the HIV-1 group (p<10<sup>-4</sup>). During the first two months of CART, the decline in plasma viral load was 3-fold steeper in the HIV-1 group (-1.56 vs. -0.62 log<sub>10</sub> copies/ml/month, p<10<sup>-4</sup>). The increases in CD4 count and in CD4 percentage were more pronounced in the HIV-1 group (+59 cells/mm<sup>3</sup>/month vs. +24 for HIV-2, Table 3). CD8 count was stable and did not differ significantly between the two groups (p=0.26). The CD4:CD8 ratio increased significantly in the two groups: +0.11/month for HIV-1 vs. +0.06 for HIV-2.

After the first two months of CART, in HIV-1 infected patients, CD4 count and CD4 percentage continue to increase: +46 cells/mm<sup>3</sup>/year and +3.3%/year, respectively. Plasma HIV RNA: -1.13 log<sub>10</sub> copies/ml/year and CD8 count: -100 cells/mm<sup>3</sup>/year decreased slightly. Therefore, the CD4:CD8 ratio increased significantly: +0.16/year. In HIV-2 infected patients, all these markers were stable (Table 4). Indeed, there was no further increase in CD4 count (-2.88 cells/mm<sup>3</sup>/year). Among the 24 (60%) patients who reached a viral load below 2.7 log<sub>10</sub> copies/mL without rebound during the first 6 months of follow-up, the slope of CD4 count was also stable (+18 cells/mm<sup>3</sup>/year, p=0.50). In the 34 (77%) HIV-1 infected patients who achieved the same viral load target, the CD4 increase was still greater (+59 cells/mm<sup>3</sup>/year, p<0.0001).

Changes in markers were not modified according to the treatment type, i.e. with or without protease inhibitor (data not shown). In a secondary analysis, we matched patients according to plasma viral load >3.5 log<sub>10</sub> copies/ml (54% of HIV-1 and 24% of HIV-2 patients) and results were similar.

## Discussion

We compared the changes in viro-immunologic markers between individuals infected with HIV-2 and individuals infected with HIV-1, all being followed in France. During natural history of infection, the estimated rates of CD4<sup>+</sup> decrease were much more pronounced in HIV-1 infected patients compared to HIV-2 infected patients. Furthermore, the estimated slopes were noticeably similar in seroprevalent and seroincident patients (-49 cells/mm<sup>3</sup>/year for HIV-1 and -9 for HIV-2, Figure 2). The plasma viral load always remained higher in HIV-1 patients compared to HIV-2 patients with the difference varying from 1.1 log<sub>10</sub> copies/ml in seroincident patients to 2.2 log<sub>10</sub> copies/ml in the patients initiating CART. Differences in CD8 count mirrored to the differences in plasma viral load. Although a formal comparison with studies performed in Africa is difficult because of the great variability in the dates of enrolment since the onset of infection, reported differences in the present study look similar to those performed in Africa [12,13,18-20,23,24,26,33,47,48]. In seroprevalent cohorts of patients not treated with antiretrovirals, the reported differences in HIV RNA varied between 1.5 log<sub>10</sub> and 3.3 log<sub>10</sub>, and between 50 and 400 cells/mm<sup>3</sup> for CD4 count [23,24,47,48]. The average difference of each marker between the seroincident HIV-1 and HIV-2 groups in the present study were also similar to those reported in a seroincident cohort of female sex workers in Senegal [18].

A novel aspect of this study is the estimation of slopes for each marker. Here again, these estimations were similar to those reported in Senegal [48] with a decline of 13% in T-cell count in HIV-1 infected patients (16% in [48]) and 3.7% in HIV-2 infected patients (4.1% in [48]). However, Gottlieb et al. reported similar slopes in both infections when controlling for plasma viral load levels. In our study, however, neither baseline plasma viral load (according to the following categories: <2.7, 2.7-3.7, >3.7, p=0.44) nor baseline CD4 count (<200, 200-500, >500, p=0.17) influenced the effect of either HIV-1 or -2 on CD4 slopes in seroprevalent patients. In other words, the differences in CD4 count decline between the two infections were similar whatever the viral load or CD4 count at the time of enrolment into the cohort.

The difference in pathogenicity between the two types of virus may be independent of environment because, in this study, the differences between the natural history HIV-1 and HIV-2 infection were similar for patients from the same geographic area. However, this study did not explore the respective roles of the host and the virus in determining the differences between the two infections. Whether differences in pathogenicity are mainly due to viral

replication, viral infectivity, cell susceptibility to activation, or CTL response remains unknown.

The virological response to CART was weaker in HIV-2 patients whatever the initial HIV RNA level. This result has been previously reported in Africa, Europe and the United-States [49-54]. In Abidjan (Côte d'Ivoire), Adjé-Touré et al. [51] reported a median viral load decrease of  $-0.6 \log_{10}$  copies/ml and an increase of  $+80$  cells/mm<sup>3</sup> for CD4 count, two months after the beginning of therapy in HIV-2 treated patients. The cause of this poor immunological response to treatment is a matter of debate [49], the first hypothesis being the limited impact of antiretroviral drugs on *in vivo* HIV-2 replication. It is difficult to distinguish whether the poorer *in vivo* response in HIV-2 patients might be linked to the potency of antiretrovirals [51,53,55] or to pathogenic features of HIV-2 infection such as the low replicative capacity [27,57]. The 50% divergence in *protease* gene nucleotides between the two types of infection could explain the reduced susceptibility of HIV-2 to the protease inhibitors developed for HIV-1 infected patients [49,57]. It is also clear that the potency against HIV-2 differs for each individual protease inhibitor [55,56] and that resistance may occur [51,58-61]. Some of these resistances are similar to those observed in HIV-1 infection [54,62] but others differs (e.g. Q151M) leading to the hypothesis that the preferred pathway for resistance development may be different between the two viruses [63]. Indeed, it could be expected that a more potent regimen leading to better virological control would improve the global response to treatment. However, the CD4 count did not increase in response to treatment in HIV-2 patients with controlled viral load during the first 6 months. It should also be noted that HIV-2 patients started an antiretroviral treatment at the same CD4 count as than HIV-1 patients, so they started therapy after a longer duration of infection. This late initiation of antiretroviral therapy in HIV-2 infection might contribute to the poorer response to treatment in particular in the CD4 increase. Hence, the findings of the present study are in favour of an earlier initiation of HAART treatment in HIV-2 infected patients.

Several limits of this study should be recognized. First, the individual matching was limited to a restricted number of potential confounding factors. We were not able to match for country of birth in all analyses because of the restricted number of patients from West Africa in the Aquitaine cohort. However, we could perform such matching for the seroprevalent group and results were similar. Another limitation was the censoring of follow-up due to change in treatment or death. This may lead to biased estimates of the change in viro-immunological markers. However, the consistency in the estimates of the slopes for each marker between the seroprevalent and seroincident groups, although the censoring was only

administrative for this latter group (three years of follow-up), argues in favour of the validity of estimates. Finally, the large number of HIV-2 infected patients with undetectable viral load yielded to insufficient information to reliably estimate the slopes. Therefore, blunted variations in HIV RNA viral load (lower than the usual measurement of  $0.5 \log_{10}$  copies/ml) need to be explored with more sensitive assays.

In conclusion, this study, the first comparing the evolution of markers between HIV-1 and HIV-2 infected patients outside of Africa, found similar differences between the two infections in Europe and in Sub-Saharan Africa. Although the difference in viral load is consistent across all analyses, the biological mechanism is still a matter of debate. The reduced response to CART in HIV-2 infected patients raises the question of optimal antiretroviral drug regimens and the right time to initiate treatment in HIV-2 infection. A better understanding of the differences in pathogenicity between the two infections may lead to improvements in treating both of them.

## Appendix

### **Composition of the ANRS CO3 Aquitaine cohort:**

Scientific committee: R. Salamon (chair), J. Beylot, M. Dupon, M. Longy-Boursier, JL.

Pellegrin, JM. Ragnaud

Coordinator: F.Dabis

Epidemiology: G. Chêne, F. Dabis, S. Lawson-Ayayi, C. Lewden, R. Thiébaud, M. Winnock

Medical coordinator: Dr M. Bonareck (Bordeaux), Dr. F. Bonnal (Bordeaux), Dr F. Bonnet (Bordeaux), Dr N. Bernard (Bordeaux), Dr O. Caubet (Bordeaux), Dr L. Caunègre (Dax), Dr J. Ceccaldi (Libourne), Pr P. Couzigou (Bordeaux), Dr C. De La Taille (Bordeaux), Dr S. De Witte (Mont de Marsan), Pr M. Dupon (Bordeaux), Dr H. Dutronc (Bordeaux), Dr S. Farbos (Bayonne), Dr T. Galpérine (Bordeaux), Dr K. Lacombe (Bordeaux), Dr D. Lacoste (Bordeaux), Dr S. Lafarie (Bordeaux), Dr P. Lose (Dax), Pr D. Malvy (Bordeaux), Pr P. Mercié (Bordeaux), Pr P. Morlat (Bordeaux), Pr D. Neau (Bordeaux), Dr A. Ochoa (Bordeaux), Pr JL. Pellegrin (Bordeaux), Pr JM. Ragnaud (Bordeaux), Dr S. Tchamgoué (Libourne), Pr J.F. Viillard (Bordeaux)

Virological coordinator: Pr H. Fleury (Bordeaux), Pr ME. Lafon (Bordeaux), Dr B. Masquelier (Bordeaux), Dr I. Pellegrin (Bordeaux)

Clinical pharmacology: Pr D. Breilh (Bordeaux)

Pharmacovigilance: Dr G. Miremont-Salamé (Bordeaux)

Immunology: Dr P. Blanco (Bordeaux), Pr JF. Moreau (Bordeaux)

Data management and statistical analysis: E. Balestre, MJ. Blaizeau, M. Decoin, S. Delveaux, L. Dequae-Merchadou, C. Hannapier, S. Geffard, S. Labarrère, V. Lavignolle-Aurillac, B. Uwamaliya-Nziyumvira

Technical team: J. Lemonnier, D. Touchard, H. Zouari, B. Boulant, N. Boulant, D. Dutoit, L. Houinou

### **Composition of the ANRS CO5 HIV-2 cohort**

Investigator coordinator: S. Matheron (Hôpital Bichat-Claude Bernard, Paris)

Virological coordinators: F. Brun-Vezinet, F. Damond (Hôpital Bichat-Claude Bernard, Paris)

Methodological coordinator: G. Chêne, A. Bénard (INSERM U593, Bordeaux)

Clinical investigators: M. Beaufils, I. Lecomte, (Tenon, Paris) ; C. Boitard, L. Roudière, JP Viard (Necker, Paris) ; PM Girard, MC Meyohas, D. Bollens, D. Berriot, F. Besse, P. Tangre, M. Kirsteter (St-Antoine, Paris) ; JB Dubuisson, A. Compagnucci, L. Finkielsztejn (Port-Royal, Paris) ; D. Sicard, C. Bernasconi, O. Zak dit Zbar (Cochin, Paris) ; W. Rozenbaum, C. Lascoux-Combe, I. Donadieu, A. Coridian, S. Thevenet, S. Courtial-Destembert (Tenon, Paris) ; J.Gilquin (St-Joseph, Paris) ; JL Vildé, P.Longuet, W. Nouioua, G. Breton, C. Charlois (Bichat, Paris) ; JP. Coulaud, S. Matheron, A. Leprêtre, S. Mas, G. Moreau, P. Campa, E. Bouvet, S. Fegueux, V. Joly, S. Lariven, PH. Consigny, I. Fournier, R. Landman, C. Voyer, G. Benabdelmoumen, C. Gaudebout, P. Yéni (Bichat, Paris) ; F. Bricaire, R.Tubiana, V. Zeller, M. Pauchard (Pitié-Salpêtrière, Paris) ; F.Meier, E. Mortier, C. Chandemerle (L. Mourier, Colombes) ; C Perronne, JC Melchior, P de Truchis (R. Poincaré, Garches) ; D. Mechali, MA Khuong-Josse, T. Labergère, B.Taverne (Delafontaine, Saint-Denis) ; A. Krivitzky, M. Bentata, JP Pathé, F. Rouges, A Mosnier, P. Honoré-Berlureau (Avicenne, Bobigny) ; P. Morel, FJ Timsit, E.Spindler (St-Louis, Paris) ; M. Thomas, H. Mouas, V. Jeantils (J. Verdier, Bondy) ; O.Patey, C. Semaille, L. Richier(Villeneuve St-Georges) ; JF Delfraissy, C.Goujard, C. Rousseau, Y. Quertainmont, T. Rannou (Bicêtre, Le Kremlin Bicêtre) ; JP Clauvel, L. Oksenhendler, L. Gérard (St-Louis, Paris) ; M. Beumont-Mauriel, G. Cessot (Institut A. Fournier, Paris) ; P. Vinceneux, M. Bloch, E. Lafon, AM SimonPoli (L. Mourier, Colombes) ; P. Lagarde, F. David-Ouaknine, E. Froguel, P. Simon

(Lagny-sur-Marne) ; P. Mornet, Y. Welker (St-Germain-en-Laye) ; M. Ruel, K. Chemlal (M. Fourestier, Nanterre) ; F. Lionnet, P. Genet (V. Dupouy, Argenteuil) ; P. Polomeni, A. Leprêtre (E. Roux, Euabonne) ; T. Papo (Bichat, Paris) ; JM Decazes, D. Ponscarne, F. Bani-Sadr, N. Colin Deverdière (St-Louis, Paris) ; O. Blétry, D. Zucman, C. Majerholc (Foch, Suresnes) ; D. Champetier de Ribes, G. Force (Perpétuel Secours, Levallois) ; S. Herson, N. Amirat, A. Coutelier, M. Brançon (Pitié-Salpêtrière, Paris) ; PY Redelsperger, B. Ponge (M. Jacquet, Melun) ; JP Escande, N. Dupin (Tarnier, Paris) ; M. Kazatchkine, M. Karmochkine, L. Weiss, D. Batisse, C. Picketty, S. Marinier-Roger, D. Tisne-Dessus (HEGP, Paris) ; M. Chousterman, V. Garrait, L. Richier (Intercommunal, Créteil) ; D. Hillion, H. Masson (Poissy) ; O. Danne, L. Blum, M. Eouzan (Pontoise) ; A. Sobel, Y. Lévy, P. Lesprit, F. Bourdillon, C. Jung (H. Mondor, Créteil) ; G. Guermontprez, A. Dulioust, P. Chardon (CMC de Bligny, Briis-sous-Forges) ; G. Raguin, M. Karmochkine (La Croix St-Simon, Paris) ; F. Bournierias (St-Cloud) ; F. Granier, V. Perronne (F. Quesnay, Mantes-la-Jolie) ; C. Lejeune, P. Bellaiche (HEGP, Paris) ; M. Janowski, C. Winter (Intercommunal, Montreuil) ; C. Vilain, I. La Torre (Montargis) ; G. Naudin (Diaconnesses, Paris) ; G. Tobelem, J. Cervoni, JM Zini (Lariboisière, Paris) ; D. Sereni, C. Lascoux-Combe, F. PrévotEAU du Clary, C. Pintado (St-Louis, Paris) ; J. Molina, D. Ponscarne, M. Lafaurie (St-Louis, Paris) ; P. Brunet (Arpajon) ; B. Godeau, A. Schaeffer (H. Mondor, Créteil) ; G. Charpentier, A. Devidas, P. Chevojon, P. Kousignian, C. Petitdidier, Y. Lemercier, I. Turpault (Corbeil) ; E. Rouveix, S. Daum-Morelon (A. Paré, Boulogne) ; P. At Khen, I. Bouchard (Etablissement public de santé national, Fresnes) ; R. Stein Metzger, S. Kernbaum (Hôpital Américain, Neuilly) ; B. Gachot (Institut Pasteur, Paris) ; P. Bourdon, JL Delassus (R. Balanger, Aulnay) ; M. Gayraud, L. Bodard, L. Raffenne (Institut Mutualiste Montsouris, Paris) ; H. Sors, D. Israël-Biet (HEGP, Paris) ; R. Roué, P. Imbert (Bégin, St-Mandé) ; P. Galanaud, F. Boué, D. Emilie, AM Delavalle (A. Béclère, Clamard) ; L. Guillevin, O. Launay, L. Belarbi (Avicenne, Bobigny) ; G. Beck-Wirth, C. Beck (Mulhouse) ; J. Barrier, F. Raffi, E. Billau (Hotel-Dieu, Nantes) ; JF Stalder, B. Milpied (Hotel-Dieu, Nantes) ; R. Laurent, C. Drobacheff, D. Bourezane (St-Jacques, Besançon) ; C. Trépo, L. Cotte, C. Brochier (Hotel-Dieu, Lyon) ; AP Blanc, T. Allègre (Aix-en-Provence) ; JA Gastaud, MP Drogoul, M. Fabre (Régional, Marseille) ; G. Dien, C. Daniel, A. Hascouet (Saint-Brieuc) ; P. Granier (Fleyriat, Bourg-en-Bresse) ; P. Choutet, JM Bestnier, D. Vautier (Bretonneau, Tours) ; F. Caron, F. Borsa Lebas, Y. Debab (C. Nicolle, Rouen) ; C. Brambilla, P. Leclerc (Michalon, Grenoble) ; M. Mornet, G. Adam (J. Cœur, Bourges) ; JP Delmont, J. Moreau, P. Brouqui (Nord, Marseille) ; T. Cartier, F. Andrieux, C. Bouvier (Pontchaillou, Rennes) ; MH Delangle, B. Héry (St-Nazaire) ; B. Becq-Giaudon, JP Breux (J. Bernard, Poitiers) ; P. Arsac (Régional, Orléans) ; A. Lafeuillade (Chalucet, Toulon) ; JM Ragnaud (Pellegrin-Tripode, Bordeaux) ; MJ Chantereau, S. Durupatient (W. Morey, Chalon-sur-Saône) ; P. Perré (La Roche sur Yon) ; M. Langlois (Villefranche de Rouergue) ; M. Dupon, H. Dutronc (Pellegrin-Tripode, Bordeaux) ; JL Pellegrin, P. Mercié (Haut-Lévêque, Bordeaux Pessac) ; P. Simonet, B. Vialatte (Cannes) ; T. May, L. Boyer (Nancy Brabois, Vandoeuvre les Nancy) ; B. Manoury, M. Daumal (St-Quentin) ; F. Janbon, J. Reynes, M. Vidal (Montpellier) ; JL Schmit, A. Smail (Amiens-Nord, Amiens) ; H. Jardel, Y. Poinsignon, D. Le Pichon (P. Chubert, Vannes) ; J. Beytout, C. Jacomet, F. Gourdon (Hotel-Dieu, Clermont Ferrand) ; H. Portier, M. Buisson, M. Greusard (Bocage, Dijon) ; Y. Mouton, F. Ajana, V. Baclet (Dron, Tourcoing) ; L. Yver, Y. Turpin (Angoulême) ; M. Uzan, D. Garipuy (J. Ducuing, Toulouse) ; M. Bonnefoy, A. Riché (Angoulême).

Virological investigators: F. Brun-Vézinet, F. Damond (Bichat, Paris) ; F. Ferchal, P. Palmer (St-Louis, Paris) ; L. Morand-Joubert (St-Antoine, Paris) ; D. Cottalorda (F. de médecine, Nice) ; MP Schmitt (F. de médecine, Strasbourg) ; P. Billaudel, V. Ferre (Institut de biologie, Nantes) ; P. Lab, A. Bassignot (Besançon) ; P. Barin (Bretonneau, Tours) ; P. Boulanger, D.

Tardy (Université, Lyon) ; D. Raoult, C. Tamalet (La Timone, Marseille) ; P. Avril, M. Kerriguy (Pontchaillou, Rennes) ; J. Puel, A. Jaafar (Purpan, Toulouse) ; P. Buffet-Janvresse (C. Nicolle, Rouen) ; P. Seigneurin, D. Morand (Grenoble) ; P. Agius, M. Bourgoïn (J. Bourgoïn, Poitiers) ; P. Fleury, B. Masquelier (Pellegrin-Tripode, Bordeaux) ; D. Duverlie, M. Roussel (Amiens) ; H La Feuille, D. Henquell (F. de médecine, Clermond-Ferrand) ; A. Chiacha (St-Michel, Angoulême) ; D. Trevoux, D. Delarbre (Mulhouse) ; C. Brehant, M. Marty (La Rochelle) ; P. Watry, D. Bocket (Lille) ; B. Montes (St-Eloi, Montpellier).  
Data management and statistical analysis: P. Campa, A. Kissila, V. El Fouikar, A. Taieb



## References

1. Chang LW, Osei-Kwasi M, Boakye D, Aidoo S, Hagy A, Curran JW et al. HIV-1 and HIV-2 seroprevalence and risk factors among hospital outpatients in the Eastern Region of Ghana, West Africa. *J Acquir Immune Defic Syndr* 2002 ; **29** :511-516
2. Diop OM, Pison G, Diouf I, Enel C, Lagarde E. Incidence of HIV-1 and HIV-2 infections in a rural community in southern Senegal. *AIDS* 2000; **14**:1671-1672
3. Schim van der Loeff MF, Akum AA, Sarge-Njie R, Van der Sande M, Jaye A, Sabbaly S et al. *Int J Epidemiol* 2006; **35**:1322-1328
4. Semaille C, Barin F, Cazein F, Pillonel J, Lot F, Brand D et al. Monitoring the dynamics of the HIV epidemic using assays for recent infection and serotyping among new HIV diagnoses: experience after 2 years in France. *J Infect Dis* 2007 ; **196** :377-383
5. Rowland-Jones SL and Whittle DH. Out of Africa: what can we learn from HIV-2 about protective immunity to HIV-1? *Nature Immunology* 2007; **8**:329-331
6. Schim van der Loeff MF, Jaffar S, Aveika AA, Sabally S, Corrah T, Harding E et al. Mortality of HIV-1, HIV-2 and HIV-1/HIV-2 dually infected patients in a clinic-based cohort in the Gambia. *AIDS* 2002; **16**:1775-1783
7. Hansmann A, Schim van der Loeff MF, Kaye S, Awasana AA, Sarge-Njie R, O'Donovan D et al. Baseline plasma viral load and CD4 Cell percentage predict survival in HIV-1 and HIV-2 infected women in a community-based cohort in the Gambia. *JAIDS* 2005; **38**:335-341
8. Whittle CH, Ariyoshi K and Rowland-Jones S. HIV-2 and T cell recognition. *Current Opinion in Immunology* 1998; **10**:382-387
9. Adjololo-Johnson G, DeCock KM, Ekpini E, Velter KM, Sibailly T, Brattegaard K et al. Prospective comparison of mother to child transmission of HIV-1 and HIV-2 in Abidjan, Ivory Coast. *JAMA* 1994; **272**:462-466
10. Kanki PJ, Travers KU, Mboup S, Hsieh CC, Marlink RG, Gueye-Ndiaye A et al. Slower heterosexual spread of HIV-2 than HIV-1. *Lancet* 1994; **343**:943-946
11. Marlink R, Kanki P, Thior I, Travers K, Eisen G, Siby T et al. Reduced rate of disease development after HIV-2 infection as compared to HIV-1. *Science* 1994; **265**:1587-1590
12. Berry N, Ariyoshi K, Jaffar S, Sabally S, Corrah T, Tedder R et al. Low peripheral blood HIV-2 RNA in individuals with high CD4 percentage differentiates HIV-2 from HIV-1 infection. *J Hum Viro* 1998; **1**:457-468
13. Jaffar S, Wilkins A, Ngom PT, Sabally S, Corrah T, Bangali JE et al. Rate of decline of percentage CD4+ cells is faster in HIV-1 than in HIV-2 infection. *J Acquir Immune Defic Syndr Hum Retrovirol* 1997; **16**:327-332
14. Michel P, Balde AT, Roussilhon C, Aribot G, Sarthou JL and Gougeon ML. Reduced immune activation and T cell apoptosis in Human Immunodeficiency Virus type 2 compared with type 1: correlation of T cell apoptosis with  $\beta_2$  microglobulin concentration and disease evolution. *J Infect Dis* 2000; **181**:64-75
15. Martinez-Steele E, Awasana AA, Corrah T, Sabally S, van der Sande M, Jaye A et al. Is HIV-2 induced AIDS different from HIV-1 associated AIDS? Data from a West African clinic. *AIDS* 2007; **21**:317-324
16. Ndour M, Sow PS, Coll-Seck AM, Badine S, Ndour CT, Diakhaté N et al. AIDS caused by HIV1 and HIV2 infection: are there clinical differences? Results of AIDS surveillance 1986-97 at Fann Hospital in Dakar, Senegal. *Tro Med Int Health* 2000; **5**:687-691

17. Matheron S, Mendoza-Sassi G, Simon F, Olivares R, Coulaud JP, Brun-Vezinet F. HIV-1 and HIV-2 AIDS in African patients living in Paris. *AIDS* 1997; **11**:934-936
18. Popper SJ, Dieng Sarr A, Travers KU, Gueye-Ndiaye A, Mboup S, Essex ME et al. Lower Human Immunodeficiency Virus (HIV) type 2 viral load reflects the difference in pathogenicity of HIV-1 and HIV-2. *J Infect Dis* 1999; **180**:1116-1121
19. Alabi AS, Jaffar S, Ariyoshi K, Blanchard T, Schim van der Loeff M, Awasana AA et al. Plasma viral load, CD4 cell percentage, HLA and survival of HIV-1, HIV-2, and dually infected Gambian patients. *AIDS* 2003; **17**:1513-1520.
20. Shanmugam V, Switzer WM, Nkengasong JN, Garcia-Lerma G, Green TA, Ekpini E, et al. Lower HIV-2 plasma viral loads may explain differences between the natural histories of HIV-1 and HIV-2 infections. *J Acquir Immune Defic Syndr* 2000; **24**:257-263
21. Gilbert PB, McKeague IW, Eisen G, Mullins C, Guéye-NDiaye A, Mboup S et al. Comparison of HIV-1 and HIV-2 infectivity from a prospective cohort study in Senegal. *Statist Med* 2003; **22**:573-593
22. Looney DJ, Hayashi S, Nicklas M, Redfield RR, Broder S, Wong-Staal F et al. Differences in the Interaction of HIV-1 and HIV-2 with CD4. *JAIDS* 1990; **3**:649-657
23. Koblavi-Dème S, Kestens L, Hanson D, Otten RA, Borget MY, Bilé C, et al. Differences in HIV-2 plasma viral load and immune activation in HIV-1 and HIV-2 dually infected persons and those infected with HIV-2 only in Abidjan, Côte D'Ivoire. *AIDS* 2004; **18**:413-419
24. Popper SJ, Dieng Sarr A, Gueye-Ndiaye A, Mboup S, Essex ME and Kanki PJ. Low plasma Human Immunodeficiency Virus type 2 viral load is independent of proviral load: low viral production in vivo. *J Virol* 2000; **74**:1554-1557
25. Arien KK, Abraha A, Quiñones-Mateu ME, Kestens L, Vanham G and Arts EJ. The replicative fitness of primary Human Immunodeficiency Virus type 1 (HIV-1) group M, HIV-1 group O, and HIV-2 isolates. *J Virol* 2005; **79**:8979-8990
26. Jaye A, Sarge-Njie R, Schim van der Loeff M, Todd J, Alabi A, Sabally S, Corrah T et al. No differences in cellular immune responses between asymptomatic HIV type 1- and type 2- infected Gambian patients. *J Infect Dis* 2004; **189**:498-505
27. MacNeil A, Sarr AD, Sankale JL, Meloni ST, Mboup S, Kanki P. Direct evidence of lower viral replication rates in vivo in HIV-2 infection than HIV-1 infection. *J Virol* 2007; **81**:5325-5330
28. Hanson A, Dieng Sarr A, Shea A, Jones N, Mboup S, Kanki P et al. Distinct profile of T cell activation in HIV type 2 compared to HIV type 1 infection: differential mechanism for immunoprotection. *AIDS Research and Human Retroviruses* 2005; **21**:791-798
29. Duvall MG, Jaye A, Dong T, Brenchley JM, Alabi AS, Jeffries DJ et al. Maintenance of HIV-specific CD4+ T cell help distinguishes HIV-2 from HIV-1 infection. *J Immunol* 2006; **176**:6973-6981
30. Lopes AR, Jaye A, Dorrell L, Sabally S, Alabi A, Jones NA et al. Greater CD8+ TCR heterogeneity and functional flexibility in HIV-2 compared to HIV-1 infection. *J Immunol* 2003; **171**:307-316
31. Zheng NN, Kiviat NB, Sow PS, Hawes SE, Wilson A, Diallo-Agne H, et al. Comparison of Human Immunodeficiency Virus (HIV)-specific T-cell responses in HIV-1 and HIV-2-infected individuals in Senegal. *J Virol* 2004; **78**:13934-13942
32. Nuvor SV, Van der Sande M, Rowland-Jones Sarah, Whittle H and Jaye A. Natural killer cell function is well preserved in asymptomatic Human Immunodeficiency Virus type 2 (HIV-2) infection but similar to that of HIV-1 infection when CD4 T-cell counts fall. *J Virol* 2006; **80**:2529-2537

33. Chollet-Martin S, Simon F, Matheron S, Joseph CA, Elbim C and Gougerot-Pocidalo MA. Comparison of plasma cytokine levels in African patients with HIV-1 and HIV-2 infection. *AIDS* 1994; **8**:879-884
34. Deeks SG, Ktichen CMR, Liu L, Guo Hua, Gascon R, Narváez AB et al. Immune activation set point during early HIV infection predicts subsequent CD4+ T-cell changes independent of viral load. *Blood* 2004; **104**:942-947
35. Schindler M, Munch J, Kutsch O, Li H, Santiago ML, Bibollet-Ruche F et al. Nef-mediated suppression lineage that gave rise to HIV-1. *Cell* 2006; **125**:1055-1067
36. Eggena MP, Barugahare B, Okello M, Mutyala S, Jones N, Ma Y, Kitoy C et al. T-cell activation in HIV-seropositive Ugandans: differential associations with viral load, CD4+ T cell depletion and co-infection. *J Infect Dis* 2005; **191**:694-701
37. Matheron S, Puyeo S, Damond F, Simon F, Leprêtre A, Campa P et al. Factors associated with clinical progression in HIV-2 infected-patients: the French ANRS cohort. *AIDS* 2003; **17**:2593-2601
38. Lazaro E, Coureau G, Guedj J, Blanco P, Pellegrin I, Commenges D et al. Change in T-lymphocyte count after initiation of highly active antiretroviral therapy in HIV-infected patients with history of Mycobacterium avium complex infection. *Antivir Ther.* 2006; **11**:343-50.
39. Rodés B, Sheldo, J, Toro C, Cuesva L, Pérez-Pastrana E, Herrera I et al. Quantitative detection of plasma human immunodeficiency virus type 2 subtype A RNA by the Nuclisens EasyQ Assay. *J Clin Microbiol.* 2007; **45**:88-92
40. Damond F, Gueudin M, Pueyo S, Farfara I, Robertson DL, Descamps D et al. Plasma RNA viral load in Human Immunodeficiency Virus type 2 subtype A and subtype B infections. *J Clin Microbiol* 2002; **40**:3654-3659
41. Damond F, Collin G, Deschamps D, Matheron S, Pueyo S, Taieb A et al. Improved sensitivity of Human Immunodeficiency Virus type 2 subtype B plasma viral load assay. *J Clin Bio* 2005; **43**:4234-4236
42. Touloumi G, Pantazis N, Babiker AG, Walker SA, Katsarou O, Karafoulidou A et al. on behalf of the CASCADE Collaboration. Differences in HIV RNA levels before the initiation of antiretroviral therapy among 1864 individuals with known HIV-1 seroconversion dates. *AIDS* 2004; **18**:1697-1705
43. CASCADE Collaboration. Differences in CD4 cell counts at seroconversion and decline among 5739 HIV-1 infected individuals with well-estimated dates of seroconversion. *J Acquir Immune Defic Syndr* 2003; **34**:76-83
44. Thiébaud R, Jacqmin-Gadda H, Walker S, Sabin C, Prins M, Del Amo J, et al. and the CASCADE Collaboration. Determinants of response to first HAART regimen in antiretroviral-naïve patients with an estimated time since HIV seroconversion. *HIV Medicine* 2006; **1**:1-9
45. Witvrouw M, Pannecouque C, Van Laethem K, Desmyter J, De Clercq E, Vandamme, AM. Activity of non-nucleoside reverse transcriptase inhibitors against HIV-2 and SIV. *AIDS* 1999; **13**:1477-1483
46. Thiébaud R, Jacqmin-Gadda H. Mixed models for longitudinal left-censored repeated measures. *Computer Methods and Programs in Biomedicine* 2004; **74**:255-260
47. Gottlieb GS, Hawes SE, Diallo-Agne H, Stern JE, Critchlow CW, Kiviat NB et al. Lower levels of HIV RNA in semen in HIV-2 compared with HIV-1 infection: implications for differences in transmission. *AIDS* 2006; **20**:895-900
48. Gottlieb GS, Sow PS, Hawes SE, Ndoeye I, Redman M, Collsek AM, et al. Equal plasma viral load predict a similar rate of CD4+ T cell decline in human immunodeficiency virus (HIV) type-1 and HIV-2 infected individuals from Senegal, West Africa. *J Infect Dis* 2002; **185**:905-914

49. Matheron S, Damond F, Benard A, Taieb A, Campa P, Peytavin G, et al. and the ANRS CO5 HIV2 Cohort Study Group. CD4 cell recovery in treated HIV-2 infected adults is lower than expected: results from the French ANRS CO5 HIV-2 Cohort.[letter] *AIDS* 2006; **20**:459-462
50. Mullins C, Eisen G, Popper S, Dieng Sarr A, Sankalé JL, Berger JJ, et al. Highly Active Antiretroviral Therapy and viral response in HIV type-2 infection. *Clinical Infectious Disease* 2004; **38**:1771-1779
51. Adjé-Touré CA, Cheingsong R, Garcia-Lerma JG, Eholié S, Borget MY, Bouchez JM, et al. Antiretroviral therapy in HIV-2 infected patients: change in plasma load, CD4+ cell count and drug resistance profiles of patients in Abidjan, Côte d'Ivoire. *AIDS* 2003; **17**:S49-S54
52. Jallow S, Kaye S, Alabi A, Aveika A, Sarge-Njie R, Sabally S et al. Virological and immunological response to Combivir and emergence of drug resistance mutations in a cohort of HIV-2 patients in the Gambia.[letter] *AIDS* 2006; **20**:1455-1458
53. van der Ende ME, Prins JM, Brinkman K, Keuter M, Veenstra J, Danner SA et al. Clinical, immunological and virological response to different antiretroviral regimens in a cohort of HIV-2 infected patients. *AIDS* 2003; **17**:S55-S61
54. Smith NA, Shaw T, Berry N, Vella C, Okorafor L, Taylor D et al. Antiretroviral therapy for HIV-2 infected patients. *J Infect* 2001; **42**:126-133
55. Withrouw M, Pannecouque C, Switzer WM, Folks TM, De Clercq E and Heneine W. Susceptibility of HIV-2, SIV and SHIV to various anti-HIV-1 compounds: implications for treatment and postexposure prophylaxis. *Antivir Ther* 2004; **9**:57-65
56. Desbois D, Peytavin G, Matheron S, Damond F, Collin G, Bénard A et al. Phenotypic Susceptibility *in vitro* to Amprenavir, Atazanavir, Darunavir, Lopinavir, and Tipranavir of HIV-2 Clinical Isolates from the French ANRS HIV-2 Cohort. In: the 14<sup>th</sup> Conference on Retroviruses and Opportunistic Infections (CROI); Feb 25-28, 2007; Los Angeles [Poster # 615]
57. Blaak H., Van der Ende M.E., Boers P.H.M., Schuitemaker H, Osterhaus A.D. In vitro replication capacity of HIV-2 variants from long-term aviremic individuals *Virology* 2006; **353**:144-154
58. Rodés B, Holguin A, Soriano V, Dourana M, Mansinho K, Antunes F et al. Emergence of drug resistance mutations in human immunodeficiency virus type-2-infected subjects undergoing antiretroviral therapy. *J Clin Microbiol* 2000; **38**:1370-1374
59. Van der Ende ME, Guillon C, Boers PH, Ly TD, Gruters RA, Osterhaus AD et al. Antiviral resistance of biologic HIV-2 clones obtained from individuals on nucleoside reverse transcriptase inhibitor therapy. *J Acquir Immune Defic Syndr* 2000; **25**:11-18
60. Damond F, Matheron S, Peytavin G, Campa P, Taieb A, Collin G et al. Selection of K65R mutation in HIV-2 infected patients receiving tenofovir-containing regimen. *Antivir Ther* 2004; **9**:635-636
61. Descamps D, Damond F, Matheron S, Collin G, Campa P, Delarue S et al. High frequency of selection of K65R and Q151M mutations in HIV-2 infected patients receiving nucleoside reverse transcriptase inhibitors containing regimen. *J Med Virol* 2004; **74**:197-201
62. Colson P, Henry M, Tourres C, Lozachmeur D, Gallais H, Gastaut JA et al. Polymorphism and drug-selected mutations in the protease gene of human immunodeficiency virus type 2 from patients living in Southern France. *J Clin Microbiol* 2004; **42**:570-577

63. Boyer PL, Sarafianos SG, Clark PK, Arnold E and Hughes SH. Why do HIV-1 and HIV-2 use different pathways to develop AZT resistance? *PLoS Pathog* 2006; **2**:101-111

**Table 1: Characteristics of patients from ANRS CO 3 Aquitaine and ANRS CO 5 HIV-2 cohorts according to the study group**

Characteristics		Seroincident group		Seroprevalent group		Naive starting CART	
		HIV-1	HIV-2	HIV-1	HIV-2	HIV-1	HIV-2
<b>Total</b>		98	49	320	160	59	63
Sex	Male	26	13	128	64	27	28
	Female	72	36	192	96	32	35
Age*	≤ 30	48	24	250	125	6	6
	30-40	30	15	70	35	14	14
	40-50	18	9	0	0	21	25
	> 50	2	1	0	0	18	18
HIV transmission group	Heterosexual	52	26	286	143	52	54
	Blood recipients	36	18	0	0	2	4
	Homosexual	4	2	10	5	2	2
	Other	6	3	24	12	3	3
Country of birth	Europe	92	10	292	55	49	12
	West Africa	0	32	6	96	2	43
	Other	6	7	22	9	6	8
Year of first CART initiation	1996-1999	-	-	-	-	9	9
	2000-2005	-	-	-	-	50	54
First CART including PI		-	-	-	-	37	37
	Ritonavir	-	-	-	-	4	1
	Ritonavir + Lopinavir	-	-	-	-	9	13
	Ritonavir + Indinavir	-	-	-	-	0	8
	Ritonavir + Saquinavir	-	-	-	-	2	2
	Ritonavir + Telzir	-	-	-	-	1	1
	Nelfinavir	-	-	-	-	9	10
	Indinavir	-	-	-	-	7	1
	Saquinavir	-	-	-	-	1	0
	Atazanavir	-	-	-	-	4	0
	Lopinavir	-	-	-	-	0	1

\* at seroconversion for seroincident patients, at inclusion for seroprevalent patients and at first CART regimen initiation for patients starting CART

**Table 2:** Characteristics at enrolment into the cohorts and at the time of treatment initiation of patients from ANRS CO 3 Aquitaine and ANRS CO 5 HIV-2 cohorts according to the study group

Characteristics	Seroincident patients		Seroprevalent patients		Naive starting CART	
	HIV-1 (n=98)	HIV-2 (n=49)	HIV-1 (n=320)	HIV-2 (n=160)	HIV-1 (n=59)	HIV-2 (n=63)
Median delay between seroconversion and baseline (years) (IQR*)	4.1 (2.0-6.9)	6.8 (3.6-14.2)	-	-	-	-
Median CD4 cell count /mm <sup>3</sup> (IQR)	399 (229-605)	585 <sup>a</sup> (458-838)	324 (172-503)	260 <sup>b</sup> (116-421)	277 (141-370)	267 (163-381)
CD4 cell count (%)	>500 350-500 <350	40 (42) 31 (63) 11 (22)	81 (25) 60 (19) 179 (56)	21 (16) 23 (18) 85 (66)	7 (12) 6 (10) 44 (78)	5 (8) 16 (25) 42 (67)
Median CD4% (IQR)	21.6 (12.6-32.6)	37.0 <sup>a</sup> (28.0-43.0)	15. (7.0-23.3)	20.0 <sup>b</sup> (10.0-28.5)	17.8 (12.0-22.6)	15.0 (12.0-25.5)
Median CD8 cell count/mm <sup>3</sup> (IQR)	842 (631-1 116)	621 <sup>a</sup> (484-830)	788 (527-1 219)	696 <sup>b</sup> (461-993)	883 (580-1 202)	670 <sup>a</sup> (376-853)
CD8 cell count (%)	>1 000 ≤1 000	34 (35) 7 (14)	111 (35) 209 (35)	30 (23) 99 (77)	18 (32) 39 (68)	7 (13) 46 (87)
Median CD4/CD8 ratio	0.45 (0.25-0.71)	1.05 <sup>a</sup> (0.62-1.42)	0.37 (0.20-0.58)	0.36 (0.15-0.68)	0.27 (0.17-0.42)	0.31 (0.23-0.54)
CD4/CD8 ratio (%)	≥1 <1	16 (16) 82 (84)	26 (53) 295 (92)	14 (12) 115 (98)	1 (2) 56 (98)	4 (7) 49 (93)
Median viral load log <sub>10</sub> cp/ml (IQR)	4.11 (3.60-4.55)	2.40 <sup>a</sup> (2.00-2.40)	4.40 (3.75-5.10)	2.62 <sup>a</sup> (2.40-3.68)	4.64 (3.01-5.18)	2.92 <sup>a</sup> (2.40-3.72)
Viral load (%)	≤2.7 >2.7	2 (7) 27 (93)	41 (89) 5 (11)	8 (8) 98 (92)	70 (54) 59 (46)	10 (17) 47 (83)

\* IQR: InterQuartile Range

Comparison between HIV-1 and HIV-2, (a) p-value <10<sup>-4</sup>, (b) p-value <0.001

**Table 3:** Estimated mean (95% Confidence Interval) slopes by year from linear mixed models for seroincident and seroprevalent patients from ANRS CO 3 Aquitaine and ANRS CO 5 HIV-2 cohorts

	Seroincident patients			Seroprevalent patients		
	HIV-1 (N=98)	HIV-2 (N=49)	p-value*	HIV-1 (N=320)	HIV-2 (N=160)	p-value*
Change in CD4 cell count/mm <sup>3</sup>	-49 (-60;-38)	-9 (-18;1)	<10 <sup>-4</sup>	-49 (-60;-41)	-11 (-18;-3)	0.0018
Change in CD4 (%)	-1.01 (-1.95;-0.07)	-0.04 (-0.41;0.33)	0.004	-0.92 (-1.45;-0.40)	-0.58 (-0.95;-0.22)	0.11
Change in viral load in log <sub>10</sub> cp/ml	-0.02 (-0.19;0.14)	0.06 (0.05;0.08)	0.87	0.20 (0.13;0.28)	0.14 (0.09;0.18)	<10 <sup>-4</sup>
Change in CD8 cell count/mm <sup>3</sup>	-5 (-33;23)	-8 (-20;3)	0.46	12 (-6;31)	17 (1;33)	0.44
Change in CD4/CD8 ratio	-0.06 (-0.08;-0.04)	0.02 (-0.00;0.04)	<10 <sup>-4</sup>	-0.06 (-0.07;-0.04)	-0.02 (-0.03;-0.00)	10 <sup>-4</sup>

\* Comparison between HIV-1 and HIV-2




**Table 4:** Estimated mean (95% Confidence Interval) slopes from linear mixed models for patients starting CART from ANRS CO 3 Aquitaine and ANRS CO 5 HIV-2 cohorts

		naive starting CART			
		HIV-1 (N=59)	HIV-2 (N=63)	p-value*	
<b>Slope during the first two months</b>	CD4 cell count/mm <sup>3</sup> /month	60 (34;84)	25 (7;42)	0.06	
	CD4%/month	3.19 (2.18;4.21)	1.25 (-0.62;3.12)	0.11	
	Viral load in log <sub>10</sub> cp/ml/month	-1.56 (-1.83;-1.30)	-0.62 (-0.84;-0.40)	<10 <sup>-4</sup>	
	CD8 cell count/mm <sup>3</sup> /month	-51 (-109;6)	2 (-70;73)	0.26	
	CD4/CD8 ratio/month	0.11 (0.06 ;0.15)	0.06 (0.02;0.10)	0.22	
	<b>Slope after the first two months</b>	CD4 cell count/mm <sup>3</sup> /year	46 (8;84)	-3 (-38;32)	<10 <sup>-4</sup>
		CD4%/year	3.29 (1.38;5.19)	0.39 (-2.55;3.33)	0.02
		Viral load in log <sub>10</sub> cp/ml/year	-1.13 (-2.35;0.09)	0.02 (-0.27;0.32)	0.42
CD8 cell count/mm <sup>3</sup> /year		-100 (-183;17)	1 (-101;104)	0.08	
CD4/CD8 ratio/year		0.16 (0.07;0.25)	-0.02 (-0.11;0.06)	0.004	

\* Comparison between HIV-1 and HIV-2



**Figure 1:** Selection of the 3 studied groups from ANRS CO 3 Aquitaine and ANRS CO 5 HIV-2 cohorts.



**Figure 2:** Estimated mean (95% Confidence Interval) CD4 T-cell slopes (cells/mm<sup>3</sup>/year for seroincident, seroprevalent and 2<sup>nd</sup>e slope of naive starting CART patients and cells/mm<sup>3</sup>/month for 1<sup>st</sup>e slope) for seroincident, seroprevalent and naive starting CART patients from ANRS CO 3 Aquitaine and ANRS CO 5 HIV-2 cohorts.

