

**The transfer of VLDL-associated phospholipids to activated platelets depends upon cytosolic phospholipase A2 activity.**

Salam Ibrahim, Catherine Calzada, Valérie Pruneta-Deloche, Michel Lagarde, Gabriel Ponsin

► **To cite this version:**

Salam Ibrahim, Catherine Calzada, Valérie Pruneta-Deloche, Michel Lagarde, Gabriel Ponsin. The transfer of VLDL-associated phospholipids to activated platelets depends upon cytosolic phospholipase A2 activity.: Mechanism of VLDL phospholipid transfer to platelets. The Journal of Lipid Research, 2007, 48 (7), pp.1533-8. 10.1194/jlr.M600480-JLR200 . inserm-00266366

**HAL Id: inserm-00266366**

**<https://www.hal.inserm.fr/inserm-00266366>**

Submitted on 21 Mar 2008

**HAL** is a multi-disciplinary open access archive for the deposit and dissemination of scientific research documents, whether they are published or not. The documents may come from teaching and research institutions in France or abroad, or from public or private research centers.

L'archive ouverte pluridisciplinaire **HAL**, est destinée au dépôt et à la diffusion de documents scientifiques de niveau recherche, publiés ou non, émanant des établissements d'enseignement et de recherche français ou étrangers, des laboratoires publics ou privés.

Copyright

# **The transfer of very low density lipoprotein- associated phospholipids to activated human platelets depends upon cytosolic phospholipase A<sub>2</sub> activity**

Salam Ibrahim, Catherine Calzada, Valérie Pruneta-Deloche, Michel Lagarde and Gabriel Ponsin<sup>#</sup>

Université de Lyon, Lyon, F-69003 ; INSERM, U870, IFR62, Lyon, F-69008 ; INRA, UMR1235, Lyon, F-69008 ; INSA-Lyon, RMND, Villeurbanne, F-69621 ; Université Lyon 1, Lyon, F-69003 ; Hospices Civils de Lyon, Lyon, F-69008, France.

**Abbreviated title** : Mechanism of VLDL phospholipids transfer to platelets

**Abbreviations** : PL, phospholipid; PC, phosphatidylcholine; PE, phosphatidylethanolamine; [<sup>14</sup>C]PAPC, 1-palmitoyl-2-[1-<sup>14</sup>C]arachidonoyl-PC; PLA<sub>2</sub>, phospholipase A<sub>2</sub>; TG, triglyceride; LDL, low density lipoprotein; HDL, high density lipoprotein; VLDL, very low density lipoprotein; LPL, lipoprotein lipase; BAPTA-AM, 1,2 – Bis (2-aminophenoxy) ethane – N, N, N', N' – tetraacetic acid tetrakis (acetoxymethyl) ester; MAFP, methyl arachidonoyl fluorophosphonate; BEL, bromoenol lactone; U73122, 1-[6-[[17β-3-Methoxyestra-1,3,5(10)-trien-17-yl]-amino]hexyl]-1H-pyrrole-2,5-dione; TXB<sub>2</sub>, thromboxane B<sub>2</sub>; PLTP, phospholipid transfer protein.

**#Corresponding author :**

Gabriel Ponsin

UMR 870 INSERM/INSA-Lyon, RMND

Bâtiment Louis Pasteur – 20 Avenue A. Einstein

69621 Villeurbanne, France

Phone : 33 4 72 43 81 13

Fax : 33 4 72 43 85 24

Email : [gabriel.ponsin@insa-lyon.fr](mailto:gabriel.ponsin@insa-lyon.fr)

## ABSTRACT

We have previously reported that VLDL could transfer phospholipids (PL) to platelets and that these transfers were favored by thrombin or lipoprotein lipase (LPL)- mediated platelet activation. The present work was undertaken to identify the platelet metabolic pathway involved in this process. The transfer of radiolabelled PL from VLDL (200  $\mu$ M PL) to platelets ( $2 \times 10^8$  /mL) was measured after incubations of 1 h at 37°C, with or without thrombin (0.1 U/mL) or LPL (500 ng/mL). To discriminate between metabolic pathways, various inhibitors, including aspirin, a cyclooxygenase inhibitor (300  $\mu$ M), esculetin, a 12-lipoxygenase inhibitor (20  $\mu$ M), Methyl-Arachidonyl-Fluorophosphate (MAFP), a phospholipase A<sub>2</sub> (PLA<sub>2</sub>) inhibitor (100  $\mu$ M), 1,2-Bis (2-aminophenoxy) ethane-N,N,N',N'-tetraacetic acid tetrakis (acetoxymethyl) ester (BAPTA-AM), an intracellular Ca<sup>2+</sup> chelator (20  $\mu$ M), bromoenol lactone (BEL), a Ca<sup>2+</sup> independent-PLA<sub>2</sub> (iPLA<sub>2</sub>) inhibitor (100 nM), or 1-[6-[[17 $\beta$ -3-Methoxyestra-1,3,5(10)-trien-17-yl]-amino]hexyl]-1H-pyrrole-2,5-dione (U73122), a phospholipase C (PLC) inhibitor (20  $\mu$ M), were added to the incubation medium. Aspirin and esculetin had no effect, showing that PL transfer was not directly dependent upon cyclooxygenase or lipoxygenase pathways. The transfer of PL was inhibited by MAFP, U73122 or BAPTA-AM. Although MAFP inhibits both cytosolic PLA<sub>2</sub> (cPLA<sub>2</sub>) and iPLA<sub>2</sub>, only cPLA<sub>2</sub> is a calcium- dependent enzyme. Since intracellular calcium mobilization is favored by PLC and inhibited by BAPTA-AM, our data suggest that the transfer of PL from VLDL to platelets results from a cPLA<sub>2</sub> rather than a iPLA<sub>2</sub>- dependent process. This conclusion was confirmed by our observation that the inhibition of iPLA<sub>2</sub> by BEL had no effect on PL transfers.

**Supplementary key words:** Thrombin, LPL, metabolic inhibitors, cPLA<sub>2</sub>, iPLA<sub>2</sub>

In platelets, phospholipids (PL) are involved in several signal transduction pathways including those that depend upon the activities of phospholipase A<sub>2</sub> (PLA<sub>2</sub>) and phospholipase C (PLC) enzymes (1-4). Platelet activation stimulates the activity of PLA<sub>2</sub> that cleave fatty acids from the sn<sub>2</sub> position of PL. In particular, the cytosolic PLA<sub>2</sub> (cPLA<sub>2</sub>) reaction favors the release of arachidonic acid which is the precursor of prostaglandins and leukotrienes generated through the actions of cyclooxygenase and lipoxygenase, respectively (1-2). In addition, in activated platelets the formation of diacylglycerols resulting from the action of PLC stimulates several metabolic cascades leading to various effects including protein phosphorylation, granule secretion and release of fatty acids by di- and monoacylglycerol lipases (3-4). Thus, platelets actively degrade PL, which necessitates their permanent regeneration. Although PL may be resynthesized in platelets (5), a substantial part has been shown to be imported from circulating lipoproteins. In vitro, low (LDL)- and high density lipoproteins (HDL), the two major human plasma lipoprotein fractions transfer various PL species to platelets, including phosphatidylcholine (PC), phosphatidylethanolamine (PE) and sphingomyelin (6-8). More recently, we have considered the possibility that very low density lipoprotein (VLDL)- associated PL could also be transferred to platelets (9). Albeit less abundant than LDL and HDL in fasting conditions, VLDL may have elevated postprandial concentrations. VLDL are secreted by the liver in the circulation where they undergo hydrolysis of their core triglyceride (TG) content through the successive actions of lipoprotein lipase (LPL) and hepatic lipase, which ultimately results in the formation of LDL (10-11). During this process, the excess of VLDL surface components, including apolipoproteins, cholesterol and PL, is released from the particles. Although a large part of cholesterol and PL is transferred to HDL (12-13), our work demonstrated that VLDL- associated PL can also be transferred to platelets, and that these transfers are favored by LPL and platelet activation (9). This effect of LPL results from two different actions. Firstly, the LPL-mediated lipolysis of VLDL destabilizes the particle surface, thereby favoring the release of PL. Secondly, the fatty acids released during lipolysis favor platelet activation, as judged from their increased thromboxane production. That the transfer of PL from VLDL to platelets depended upon platelet activation was confirmed by our observation that thrombin stimulates both platelet thromboxane production and PL transfer.

Although platelets are able to import PL from various lipoprotein fractions, the underlying mechanisms of these transfers appeared to be complex. They are independent of lipoprotein binding and internalization (7). In agreement with this concept, the scavenger receptor B1 which can mediate the specific import of PL into various cells was shown to be absent in platelets (8). However, major differences emerge when comparing the PL transfers obtained from the different lipoprotein fractions. The transfer of LDL or HDL - derived PE into platelets, but not that of PC or sphingomyelin, was stimulated by platelet activators including thrombin, collagen and ADP and was dependent upon the secretion of an unidentified cellular protein factor (14). In contrast, there was no apparent specificity of the PL species transferred from VLDL to platelets (9). Both LPL and thrombin stimulated the import by platelets of VLDL- derived Palmitoyl-Arachidonyl-PC, Palmitoyl-Arachidonyl-PE and DiPalmitoyl-PC with similar efficiencies. Thus the regulation of the PL uptake by platelets appears to dramatically depend upon the lipoprotein used as the donor. While LDL might preferentially transfer certain PL species, namely PE, by a specific mechanism, VLDL could supply all types of PL to platelets without consideration of their nature. This concept prompted us to further characterize the metabolic pathway governing the transfer of PL from VLDL to platelets. In this work, using a variety of metabolic inhibitors, we present *in vitro* evidences showing that this transfer results from a cPLA<sub>2</sub>-dependent process.

## MATERIALS AND METHODS

### Materials

[1-<sup>14</sup>C] arachidonic acid (40–60 mCi/mmol) and 1-Palmitoyl-2-[1-<sup>14</sup>C] arachidonyl-phosphatidylcholine ([<sup>14</sup>C] PAPC; 40–60 mCi/mmol) were purchased from Perkin-Elmer (Boston, MA). Thrombin, bovine milk LPL (EC 3.1.1.34), 1,2 – Bis (2-aminophenoxy) ethane – N, N, N', N' – tetraacetic acid tetrakis (acetoxymethyl) ester (BAPTA-AM) and polyoxyethylen-9-

laurylether were obtained from Sigma Chemical (St. Louis, MO). Phospholipids were assayed using enzymatic kits from Wako chemicals GmbH (Neuss, Germany). Thromboxane B<sub>2</sub> (TXB<sub>2</sub>) concentrations were determined using the enzyme immunoassay Biotrak system from Amersham Biosciences (Orsay, France). Esculetin, methyl arachidonyl fluorophosphonate (MAFP), 1-[6-[[17β-3-Methoxyestra-1,3,5(10)-trien-17-yl]-amino]hexyl]-1H-pyrrole-2,5-dione (U73122) and bromoenol lactone (BEL) were from Biomol (Plymouth Meeting, PA).

### **Isolation and labelling of lipoproteins**

VLDL ( $d < 1.006$  g/mL) and lipoprotein-deprived plasma ( $d > 1.21$  g/mL) were isolated from human plasma by preparative ultracentrifugation (15). Depending upon the volume of plasma, the ultracentrifugation was performed either in a Beckman LE 80K using a 50.2 fixed-angle rotor or in a Beckman TL-100 tabletop ultracentrifuge using a TLA 100.3 fixed-angle rotor. The resulting preparations were then extensively dialyzed against a buffer containing 150 mM NaCl, 1 mM EDTA, 10 mM Tris-HCl, and 10 mM NaN<sub>3</sub>, pH 7.4. VLDL were labelled with [<sup>14</sup>C] PAPC (1 μCi/10 μmol lipoprotein PL) as reported previously (16). The desired amount of radioactive label was dried under nitrogen, solubilized in ethanol, and added to VLDL under vortexing. To avoid destruction of the lipoprotein structure, the final proportion of ethanol in the samples was maintained at < 1% (v/v). The samples were then incubated for 3 h at 37°C, and finally the labelled VLDL were reisolated by ultracentrifugation.

### **Platelet isolation**

Fresh blood was collected at the local blood bank (Etablissement Français du Sang) from healthy volunteers. Blood was drawn into a one-seventh volume of a solution containing 19.6 mM citric acid, 89.4 mM sodium citrate, 16.1 mM NaH<sub>2</sub>PO<sub>4</sub>, and 128.7 mM dextrose, pH 5.6. The platelet isolation procedure was essentially based on that described previously (17). Briefly, platelet-rich plasma was obtained after blood centrifugation at 200 g for 17 min at 20°C and acidified to pH 6.4 with 0.15 M citric acid. Platelets were immediately pelleted by centrifugation at 900 g for 12 min and washed in acidified lipoprotein-deprived plasma. After repelleting, the platelets were finally washed

and resuspended in a Tyrode-HEPES buffer solution containing 137 mM NaCl, 2.7 mM KCl, 0.41 mM NaH<sub>2</sub>PO<sub>4</sub>, 11.9 mM NaHCO<sub>3</sub>, 1 mM MgCl<sub>2</sub>, 5.5 mM glucose, and 5 mM HEPES, pH 7.35. Platelet suspensions were left for 1 h at room temperature before experiments were started.

### **Platelet aggregation**

To avoid abnormal experimental data that could result from unknown medical treatment taken by blood donors, each platelet preparation was controlled for its functional ability to aggregate before being used in our studies. Aggregations were induced by arachidonic acid and performed in a Chronolog dual-channel aggregometer (Coulter, Margency, France) according to the turbidimetric method of Born (18).

### **Transfers of labelled phospholipids from VLDL to platelets**

Labelled VLDL (200 nmol/ml PL) were incubated with platelets ( $3 \times 10^8$  cells) in a final volume of 1.5 ml at 37°C for 1 h (9). For each given experiment the VLDL and platelets were each isolated from a single donor. At the end of the incubations, platelets were separated from the medium by centrifugation. The pelleted platelets were first washed in plasma to remove non-specifically adsorbed labelled VLDL and then in Tyrode-HEPES buffer. The final pellets were dissolved by overnight incubation in 0.25 ml of 0.4% polyoxyethylen-9-laurylether and counted for radioactivity. The results were finally corrected for non-specifically adsorbed radioactivity at time zero (non-incubated platelets).

When platelet activation was desired, LPL (500 ng/ml) or thrombin (0.1 U/ml) were added at the beginning of the incubations. In contrast, to inhibit specific metabolic pathways various inhibitors were added to platelets during pre-incubation periods as follows: 2 min for U73122 (20 μM), a PLC inhibitor (19), 5 min for aspirin (300 μM), a cyclooxygenase inhibitor (20) and esculetin (20 μM), a lipoxygenase inhibitor (21), 10 min for MAFP (100 μM), a cPLA<sub>2</sub> and iPLA<sub>2</sub> inhibitor (22-23) and BEL (100 nM), a Ca<sup>2+</sup> independent-PLA<sub>2</sub> (iPLA<sub>2</sub>) inhibitor (22,24), and 20 min for BAPTA-AM (20 μM), an intracellular Ca<sup>2+</sup> chelator (25-26).

### **Determination of thromboxane B<sub>2</sub> production**

The production of TXB<sub>2</sub> by platelets was determined at the beginning and at the end of incubations carried out under various conditions as described above. However, platelets were not separated from their media before the TXB<sub>2</sub> assay, thereby permitting the measurement of total TXB<sub>2</sub>.

### **Determination of the [1-<sup>14</sup>C] arachidonic acid content of platelet phospholipids**

Phospholipid hydrolysis was assayed in platelets prelabelled with [1-<sup>14</sup>C] arachidonic acid (20 nCi/mL) for 1 h at 37°C (27). After separation from the medium by centrifugation, platelets were washed in Tyrode-HEPES buffer and incubated for 5 min in the absence or presence of thrombin or cPLA2 inhibitors. For these particular experiments, no VLDL was added to the incubation medium to avoid phospholipid transfers. At the end of the incubations, platelets were washed and resuspended in one ml of buffer. Lipids were then extracted and separated by thin layer chromatography in a solvent composed of hexane : diethyl ether : acetic acid (80 : 20 : 1). The phospholipid spots were finally scrapped off and counted for radioactivity.

### **Determination of plasma phospholipid transfer protein activity**

Phospholipid transfer protein (PLTP) activity was determined using an in vitro assay in which we measured the transfer of radiolabelled phospholipid from VLDL to HDL using delipidated plasma as the source of PLTP, as previously described (28). Basal as well as PLTP-facilitated phospholipid transfers were measured in the absence or presence of cPLA2 inhibitors at the same concentrations as those used with platelets. The results were expressed as percent of phospholipid transferred during two hours incubations.

## **RESULTS**

To explore the dependence of VLDL- associated PL transfer to platelets upon platelet activation, we performed a first series of experiments in which various inhibitors were used to block



different metabolic pathways involved in platelet activation. The transfers of [ $^{14}\text{C}$ ]PAPC from VLDL to platelets were measured after incubations of one hour at 37°C. They were stimulated by about 2.5- and 3.5- fold when platelets were activated by either thrombin (0.1 U/mL) or LPL (500 ng/mL), respectively (**Fig. 1**). When the experiments were performed in the presence of aspirin, a cyclooxygenase inhibitor, at a concentration (300  $\mu\text{M}$ ) known to block the formation of thromboxane, no change was observed in the transfers of PL whether or not stimulated by either thrombin or LPL. Similarly, the transfers of PL remained unchanged when the 12-lipoxygenase-dependent metabolic pathway was inhibited by esculetin (20  $\mu\text{M}$ ). In contrast, when the incubations were carried out in the presence of MAFP, a PLA<sub>2</sub> inhibitor, both the stimulating effects of thrombin and LPL on PL transfers were abolished, resulting in values comparable to that of controls.

Three different PLA<sub>2</sub> enzymes are present in human platelets, including cPLA<sub>2</sub>, iPLA<sub>2</sub> and secretory PLA<sub>2</sub>. Only the two former may be affected by MAFP. However, their activities can be discriminated on the basis of their Ca<sup>2+</sup> dependence since cPLA<sub>2</sub> but not iPLA<sub>2</sub> is a Ca<sup>2+</sup> dependent enzyme. This prompted us to study the thrombin- or LPL- stimulated transfers of [ $^{14}\text{C}$ ]PAPC from VLDL to platelets in the presence of either BAPTA-AM, a Ca<sup>2+</sup> chelator, or U73122, a PLC inhibitor (**Fig. 2**). The results showed that both BAPTA-AM and U73122 inhibited the stimulating effects of thrombin and LPL, clearly suggesting that PL import into platelets was controlled by a cPLA<sub>2</sub> rather than iPLA<sub>2</sub>- dependent process. This was confirmed by the results obtained when the incubations were performed in the presence of 100 nM BEL. The latter, which at this concentration inhibits iPLA<sub>2</sub> but not cPLA<sub>2</sub> (29), had no effect on PL transfers whether or not stimulated by either thrombin or LPL. To control that metabolic inhibitors did not directly modify the ability of VLDL to transfer phospholipids, in particular by inhibiting the activity of PLTP that could be present at the lipoprotein surface, we performed in vitro assays in which we measured the transfers of radiolabelled phospholipids from VLDL to HDL (**Table 1**). None of MAFP, BAPTA-AM or U73122 had any effect on basal or PLTP-facilitated phospholipid transfers.

Finally, we determined the metabolic effects of BAPTA-AM and U73122 on platelet phospholipid hydrolysis and TXB<sub>2</sub> production. In the absence of inhibitors, we observed a decrease of the arachidonic content of phospholipids that was stimulated by thrombin, clearly showing a major

hydrolysis (**Table 2**). In the presence of U73122 which inhibits both cPLA<sub>2</sub> and PLC, phospholipid hydrolysis was strongly inhibited, while it was only partially decreased in the presence of BAPTA-AM which inhibits only cPLA<sub>2</sub>. In addition, clear stimulating effects of thrombin and LPL emerged when platelet production of TXB<sub>2</sub> was studied (**Fig.3**). The thrombin-stimulated production of TXB<sub>2</sub> was totally inhibited by BAPTA-AM or U73122, while they only partially decreased the stimulating effect of LPL. As expected, BEL, that specifically inhibits iPLA<sub>2</sub>, had no effect on TXB<sub>2</sub> production.

## DISCUSSION

We have recently shown that platelets are able to import PL from VLDL and that this process is stimulated when platelets are activated through the action of thrombin or LPL (9). The present work was intended to characterize the relationship between the transfer of PL from VLDL to platelets and platelet activation. The latter results from the combination of complex metabolic pathways that can be summarized as follows. Plasma membrane PL may be hydrolyzed by cPLA<sub>2</sub>, resulting in the release of fatty acids. Among these, arachidonic acid can be metabolized through two different pathways (30-32). The first, that depends upon cyclooxygenase activity, leads to the formation of prostaglandins and thromboxane A<sub>2</sub>, a very potent platelet activator. The second pathway is governed by the action of 12-lipoxygenase that transforms arachidonic acid in 12-hydroperoxy-eicosatetraenoic acid, a metabolite also able to favor platelet activation. In addition, phosphatidylinositoldiphosphate (PIP<sub>2</sub>) may also be hydrolyzed by PLC, leading to the formation of diacylglycerols and inositoltriphosphate (IP<sub>3</sub>). While diacylglycerols stimulate metabolic cascades resulting in protein phosphorylation, granule secretion and release of fatty acids by di- and monoacylglycerol lipases, IP<sub>3</sub> favors the increase of Ca<sup>2+</sup> intracellular concentration (3-4).

To study the putative involvement of these metabolic pathways, we measured the transfer of PL from VLDL to platelets in the presence of aspirin or esculetin that inhibit cyclooxygenase and 12-lipoxygenase, respectively (20-21). We observed no changes in the transfers of PL to platelets whether or not stimulated by thrombin or LPL, indicating that neither cyclooxygenase- nor 12-lipoxygenase-

dependent processes were directly responsible for the magnitude of PL transfers. In contrast, these PL transfers were decreased when the incubations were performed in the presence of MAFP, an inhibitor of PLA<sub>2</sub>. PLA<sub>2</sub> is a superfamily of enzymes consisting of secretory and intracellular species. The latter comprise cPLA<sub>2</sub> and iPLA<sub>2</sub> which are both inhibited by MAFP (22-23). To discriminate between these two enzymes, we took advantage of the differences in their mechanism of action. cPLA<sub>2</sub> but not iPLA<sub>2</sub> is a Ca<sup>2+</sup> dependent- enzyme (29). The activation of cPLA<sub>2</sub> depends upon two synergistic processes: the catalytic activity of the enzyme requires a previous phosphorylation while its translocation to the membrane necessitates the mobilization of cytosolic Ca<sup>2+</sup> (33-34). Since the latter may be regulated by the PLC pathway, we measured the thrombin- and LPL- stimulated transfers of PL from VLDL to platelets in the presence of U73122, a PLC inhibitor (19). The PL transfers were clearly inhibited. However, as mentioned above, in addition to IP<sub>3</sub>- stimulated Ca<sup>2+</sup> mobilization, the action of PLC generates diacylglycerols that stimulate different metabolic cascades (3-4). Thus, to distinguish between these various effects, we performed experiments where U73122 was substituted for BAPTA-AM, a Ca<sup>2+</sup> chelator (25-26). The PL transfers were similarly decreased, showing that the effect of PLC was due to its ability to mobilize cytosolic Ca<sup>2+</sup>. Overall consideration of these results clearly suggests that the transfer of PL from VLDL to platelets depends upon a cPLA<sub>2</sub> rather than an iPLA<sub>2</sub>- dependent process. This concept was finally confirmed by our observation that PL transfers remained unaffected when iPLA<sub>2</sub> was specifically inhibited in the presence of BEL (24).

In addition to phospholipid transfers, the importance of cPLA<sub>2</sub> in platelet metabolism was assessed by measuring the arachidonic acid content of phospholipids in platelets and their TXB<sub>2</sub> production. A strong phospholipid hydrolysis was observed during platelet incubations, that was in part dependent upon cPLA<sub>2</sub> activity. As previously shown (9), both thrombin and LPL stimulated the platelet TXB<sub>2</sub> production. The stimulating effect of thrombin was totally inhibited by BAPTA-AM or U73122 while that of LPL was only partially decreased. This could be expected since LPL-stimulated platelet activation is initially due to the uptake by platelets of the fatty acids released during LPL-mediated lipolysis of VLDL (9).

Although the question as to how the transfers of PL are precisely affected by cPLA<sub>2</sub> activity has not been directly addressed in this work, a likely mechanism may be considered. Since the

cPLA<sub>2</sub>- stimulated hydrolysis of PL occurs at the inner leaflet of the plasma membrane, the enzyme activity necessarily results in disequilibrium of the PL concentrations between the membrane inner and outer leaflets. Several physiological processes are known to lead to comparable disequilibrium which is compensated by flip-flop mechanisms (35-36). Thus, on the basis of a similar mechanism, we might speculate that in our case, the net translocation of PL from the outer to the inner leaflet of the membrane would cause a PL deficit in the outer leaflet that might be compensated by the import of PL from VLDL.

## **ACKNOWLEDGEMENTS**

This work was supported by the Institut National de la Santé et de la Recherche Médicale. Salam Ibrahim was a recipient for a grant from the French Ministry of Education and Research.

## REFERENCES

1. Samuelsson, B., M. Goldyne, E. Granström, M. Hamberg, S. Hammarström, and C. Malmsten. 1978. Prostaglandins and thromboxanes. *Ann. Rev. Biochem.* **47**: 997-1029.
2. Clark, J.D., A.R. Schievella, E.A. Nalefski, and L.L. Lin. 1995. Cytosolic phospholipase A2. *J. Lipid Mediat. Cell Signal.* **12**: 83-117.
3. Ha, K.S., and J.H. Exton. 1993. Differential translocation of protein kinase C isozymes by thrombin and platelet-derived growth factor. A possible function for phosphatidylcholine-derived diacylglycerol. *J. Biol. Chem.* **268**: 10534-10539.
4. Hug, H., and T.F. Sarre. 1993. Protein kinase C isoenzymes: divergence in signal transduction ? *Biochem. J.* **291**: 329-343.
5. Kent, C. 1995. Eukariotic phospholipid biosynthesis. *Ann. Rev. Biochem.* **64**: 315-343.
6. Engelmann, B., C. Kögl, R. Kulschar, and B. Schaipp. 1996. Transfer of phosphatidylcholine, phosphatidylethanolamine and sphingomyelin from low- and high-density lipoprotein to human platelets. *Biochem. J.* **315**: 781-789.
7. Dobner, P., E. Koller, and B. Engelmann. 1999. Platelet high affinity low density lipoprotein binding and import of lipoprotein derived phospholipids. *FEBS Lett.* **444**: 270-274.
8. Urban, S., S. Zieseniss, M. Werder, H. Hauser, R. Budzinski, and B. Engelmann. 2000. Scavenger receptor B1 transfers major lipoprotein-associated phospholipids into the cells. *J. Biol. Chem.* **275**: 33409-33415.
9. Ibrahim, S., A. Djimet-Baboun, V. Pruneta-Deloche, C. Calzada, M. Lagarde, and G. Ponsin. 2006. Transfer of very low density lipoprotein-associated phospholipids to activated human platelets. *J. Lipid Res.* **47**: 341-348.
10. Jackson, R.L., L.R. McLean, and R.A. Demel. 1987. Mechanism of action of lipoprotein lipase and hepatic triglyceride lipase. *Am. Heart J.* **113**: 551-554.

11. Deckelbaum, R.J., S. Eisenberg, M. Fainaru, Y. Barenholz, and T. Olivecrona. 1979. In vitro production of human low density lipoprotein-like particles: a model for very low density lipoprotein catabolism. *J. Biol. Chem.* **254**: 6079-6087.
12. Glomset, J.A. 1968. The plasma lecithin: cholesterol acyl transferase reaction. *J. Lipid Res.* **9**: 155-167.
13. Patsch, J.M., A.M. Gotto Jr., T. Olivecrona, and S. Eisenberg. 1978. Formation of high density lipoprotein 2- like particles during lipolysis of very low density lipoproteins in vivo. *Proc. Natl. Acad. Sci. USA.* **75**: 4519-4523.
14. Engelmann, B., B. Schaipp, P. Dobner, M. Stoeckelhuber, C. Kögl, W. Siess, and A. Hermetter. 1998. Platelet agonists enhance the import of phosphatidylethanolamine into human platelets. *J. Biol. Chem.* **273**: 27800-27808.
15. Havel, R.J., H.A. Eder, and J.H. Bragdon. 1955. The distribution and chemical composition of ultracentrifugally separated lipoproteins in human serum. *J. Clin. Invest.* **34**: 1345-1354.
16. Elchebly, M., T. Pulcini, B. Porokhov, F. Berthezène, and G. Ponsin. 1996. Multiple abnormalities in the transfer of phospholipids from VLDL and LDL to HDL in non-insulin-dependent diabetes. *Eur. J. Clin. Invest.* **26**: 216-223.
17. Lagarde, M., P.A. Bryon, M. Guichardant, and M. Dechavanne. 1980. A simple and efficient method for platelet isolation from their plasma. *Thromb. Res.* **17**: 581-588.
18. Born, G.V.R. 1962. Aggregation of blood platelets by adenosine diphosphate and its reversal. *Nature.* **194**: 927-929.
19. Bleasdale, J.E., N.R. Thakur, R.S. Gremban, G.L. Bundy, F.A. Fitzpatrick, R.J. Smith, and S. Bunting. 1990. Selective inhibition of receptor-coupled phospholipase C-dependent processes in human platelets and polymorphonuclear neutrophils. *J. Pharmacol. Exp. Ther.* **255**:756–768.
20. Vane, J.R., R.M. Botting. 2003. The mechanism of action of aspirin. *Thromb. Res.* **110**: 255-258.
21. Sekiya, K., H. Okuda, S. Arichi. 1982. Selective inhibition of platelet lipoxygenase by esculetin. *Biochim Biophys Acta.* **713**: 68-72.
22. Balsinde, J., M.A. Balboa, P.A. Insel, and E.A. Dennis. 1999. Regulation and inhibition of phospholipase A<sub>2</sub>. *Annu. Rev. Pharmacol. Toxicol.* **39**:175–89.

23. Lio, Y.C., L.J. Reynolds, J. Balsinde, E.A. Dennis. 1996. Irreversible inhibition of  $\text{Ca}^{2+}$  independent phospholipase  $\text{A}_2$  by methyl arachidonyl fluorophosphonate. *Biochim Biophys Acta*. **1302**: 55-60.
24. Ackermann, A.J., K. Conde-Frieboes, and E.A. Dennis. 1995. Inhibition of Macrophage  $\text{Ca}^{2+}$ -independent Phospholipase  $\text{A}_2$  by Bromoenol Lactone and Trifluoromethyl Ketones. *J. Biol. Chem.* **270**: 445-450.
25. Börsch-Haubold, A.G., R.M. Kramer, and S.P. Watson. 1995. Cytosolic phospholipase  $\text{A}_2$  is phosphorylated in collagen- and thrombin-stimulated human platelets independent of protein kinase C and Mitogen-activated Protein Kinase. *J. Biol. Chem.* **270**: 25885–25892.
26. Roberts, D.E., A. McNicol, and R. Bose. 2004. Mechanism of collagen activation in human platelets. *J. Biol. Chem.* **279**: 19421–19430.
27. Bills, T.K., J.B. Smith, and M.J. Silver. 1977. Selective release of arachidonic acid from the phospholipids of human platelets in response to thrombin. *J. Clin. Invest.* **60**: 1-6.
28. Elchebly, M., T. Pulcini, B. Porokhov, F. Berthezène, and G. Ponsin. 1996. Multiple abnormalities in the transfer of phospholipids from VLDL and LDL to HDL in non-insulin-dependent diabetes. *Eur. J. Clin. Invest.* **26**:216-223.
29. HaZen, S.L., L.A. Zupan, R.H. Weissg, D.P. Getmang, and R.W. Gross. 1991. Suicide inhibition of canine myocardial cytosolic calcium-independent phospholipase  $\text{A}_2$ : Mechanism-based discrimination between calcium-dependent and –independent phospholipases  $\text{A}_2$ . *J. Biol. Chem.* **266**: 7227-7232.
30. Blockmans, D., H. Deckmyn, and J. Vermynen. 1995. Platelet activation. *Blood Reviews.* **9**: 143-156.
31. Balsinde, J., M.V. Winstead, and E.A. Dennis. 2002. Phospholipase  $\text{A}_2$  regulation of arachidonic acid mobilization of arachidonic acid. *FEBS Lett.* **531**: 2-6.
32. Coulon, L., C. Calzada, P. Moulin, E. Vericel, and M. Lagarde. 2003. Activation of p38 mitogen-activated protein kinase/cytosolic phospholipase  $\text{A}_2$  cascade in hydroperoxide-stressed platelets. *Free Radic. Biol. Med.* **35**:616-25.



33. Yedgar, S., D. Lichtenberg, and E. Schnitzer. 2000. Inhibition of phospholipase A<sub>2</sub> as a therapeutic target. *Biochim Biophys Acta*. **1488**: 182-187.
34. McNicol, A., and Shibou, T.S. 1998. Translocation and phosphorylation of cytosolic phospholipase A<sub>2</sub> in activated platelets. *Thromb. Res.* **92**: 19–26.
35. Bevers, E.M., P. Comfurius, D.W. Dekkers, and R.F. Zwaal. 1999. Lipid translocation across the plasma membrane of mammalian cells. *Biochim. Biophys. Acta*. **1439**: 317–330.
36. Sims, P.J., and Wiedmer, T. 2001. Unraveling the mysteries of phospholipids scrambling. *Thromb. Haemost.* **86**: 266–275.

## FIGURE LEGENDS

**Fig.1.** Effect of metabolic inhibitors on the transfers of [ $^{14}\text{C}$ ] PAPC from VLDL to platelets. Platelets were incubated with [ $^{14}\text{C}$ ]PAPC- labelled VLDL (200 nmoles of PL / mL) for 1 h at 37°C in a total volume of 1.5 mL, in the presence or absence of thrombin (0.1 U/mL) or LPL (500 ng / mL). To study the putative dependence of PL transfers upon various metabolic pathways, the results obtained in basal conditions (open bars) were compared to those resulting from the effects of aspirin (300  $\mu\text{M}$ ), a cyclooxygenase inhibitor (closed bars), esculetin (20  $\mu\text{M}$ ), a 12-lipoxygenase inhibitor (hatched bars) or MAFP (100  $\mu\text{M}$ ), a phospholipase A<sub>2</sub> inhibitor (grey bars). Transfers were expressed as the percentage of radioactivity incorporated by  $3 \times 10^8$  platelets. Values shown are means  $\pm$  SEM from four to six independent experiments.

**Fig.2.** Comparison of the effects of cytosolic Ca<sup>2+</sup>- dependent (cPLA<sub>2</sub>) and Ca<sup>2+</sup>- independent phospholipase A<sub>2</sub> (iPLA<sub>2</sub>) on the transfer of [ $^{14}\text{C}$ ] PAPC from VLDL to platelets. Platelets were incubated with [ $^{14}\text{C}$ ]PAPC- labelled VLDL (200 nmoles of PL / mL) for 1 h at 37°C in a total volume of 1.5 mL, with or without thrombin (0.1 U/mL) or LPL (500 ng / mL). To discriminate between the putative effects of cPLA<sub>2</sub> and iPLA<sub>2</sub>, the results obtained in basal conditions (open bars) were compared to those observed in the presence of U73122 (20  $\mu\text{M}$ ), a phospholipase C inhibitor (closed bars), BAPTA-AM (20  $\mu\text{M}$ ), a Ca<sup>2+</sup> chelator (hatched bars) or BEL (100 nM), an iPLA<sub>2</sub> inhibitor (grey bars). Transfers were expressed as the percentage of radioactivity incorporated by  $3 \times 10^8$  platelets. Values shown are means  $\pm$  SEM from four independent experiments.

**Fig.3.** Effects of cytosolic Ca<sup>2+</sup>- dependent (cPLA<sub>2</sub>) and Ca<sup>2+</sup>- independent phospholipase A<sub>2</sub> (iPLA<sub>2</sub>) inhibitors on the production of thromboxane B<sub>2</sub> (TXB<sub>2</sub>) by platelets. Platelets were incubated with VLDL (200 nmoles of PL / mL) for 1 h at 37°C in a total volume of 1.5 mL, in the absence (open bars) or presence of U73122 (20  $\mu\text{M}$ ), a phospholipase C inhibitor (closed bars), BAPTA-AM (20  $\mu\text{M}$ ), a Ca<sup>2+</sup> chelator (hatched bars) or BEL (100 nM), an iPLA<sub>2</sub> inhibitor (grey bars). The thrombin (0.1 U/mL)- or LPL (500 ng / mL)- stimulated productions of TXB<sub>2</sub> were expressed as the percentage of those obtained in basal conditions. TXB<sub>2</sub> concentrations were measured using a commercial kit, as described in Materials and Methods. Values shown are means  $\pm$  SEM from three independent experiments.