



HAL
open science

IL-18 Does not Increase Allergic Airway Disease in Mice When Produced by BCG.

Latiffa Amniai, Franck Biet, Philippe Marquillies, Camille Locht, Joël Pestel, André-Bernard Tonnel, Catherine Duez

► **To cite this version:**

Latiffa Amniai, Franck Biet, Philippe Marquillies, Camille Locht, Joël Pestel, et al.. IL-18 Does not Increase Allergic Airway Disease in Mice When Produced by BCG.. *Journal of Biomedicine and Biotechnology*, 2007, 2007, pp.67276. 10.1155/2007/67276 . inserm-00305279

HAL Id: inserm-00305279

<https://inserm.hal.science/inserm-00305279>

Submitted on 24 Jul 2008

HAL is a multi-disciplinary open access archive for the deposit and dissemination of scientific research documents, whether they are published or not. The documents may come from teaching and research institutions in France or abroad, or from public or private research centers.

L'archive ouverte pluridisciplinaire **HAL**, est destinée au dépôt et à la diffusion de documents scientifiques de niveau recherche, publiés ou non, émanant des établissements d'enseignement et de recherche français ou étrangers, des laboratoires publics ou privés.



Distributed under a Creative Commons Attribution 4.0 International License

IL-18 does not increase allergic airway disease in mice when produced by BCG

Latiffa Amniai ¹, Franck Biet ², Philippe Marquillies ¹, Camille Locht ³, Joël Pestel ⁴, André-Bernard Tonnel ¹, Catherine Duez ¹.

¹ INSERM, Institut National de la Santé et de la Recherche Médicale U774, 59019 Lille, France ; Institut Pasteur de Lille, 59019 Lille, France ; Université de Lille 2, 59800 Lille, France

² UR 1282, Infectiologie animale, Santé publique (IASP-311), INRA centre de Tours, 37380 Nouzilly, France

³ INSERM, Institut National de la Santé et de la Recherche Médicale U629, 59019 Lille, France ; Institut Pasteur de Lille, 59019 Lille, France

⁴ UMR CNRS 8017, 59655 Villeneuve d'Ascq, France ; Université de Lille I 59655 Villeneuve d'Ascq, France

Corresponding author:

Catherine Duez

INSERM U774, Institut Pasteur de Lille

1 rue du Pr Calmette, BP 245

59019, Lille Cedex, France

Phone: (33) 3 20 87 71 83

Fax: (33) 3 20 87 73 45

e-mail: catherine.duez@pasteur-lille.fr

Abstract

Whilst BCG inhibits allergic airway responses in murine models, IL-18 has adversary effects depending on its environment. We therefore constructed a BCG strain producing murine IL-18 (BCG-IL-18) and evaluated its efficiency to prevent an asthma-like reaction in mice.

BALB/cByJ mice were sensitized (day (D) 1 and D10) by intraperitoneal injection of ovalbumin (OVA)-alum and primary (D20-22) and secondary (D62, 63) challenged with OVA aerosols. BCG or BCG-IL-18 were intraperitoneally administered 1hr before each immunization (D1 and D10). BCG-IL-18 and BCG were shown to similarly inhibit the development of AHR, mucus production, eosinophil influx and local Th2 cytokine production in BAL, both after the primary and secondary challenge.

These data show that IL-18 did not increase allergic airway responses in the context of the mycobacterial infection, and suggest that BCG-IL-18 and BCG are able to prevent the development of local Th2 responses and therefore inhibit allergen-induced airway responses even after restimulation.

Key words: Asthma, mouse model, BCG, IL-18, airway hyperresponsiveness

Abbreviations: AHR = Airway hyperresponsiveness, BAL: Bronchoalveolar lavage, BCG:

Mycobacterium bovis Bacillus Calmette-Guerin

Introduction

Airway inflammation and hyperresponsiveness, increased Th2 cytokine production in bronchoalveolar lavage and enhanced IgE secretion in serum are the hallmarks of allergic asthma. Increased prevalence of asthma has been related to decreased microbial infection or immunization during early infancy. In particular, *Mycobacterium bovis* Bacillus Calmette-Guerin (BCG) vaccination have been suggested to decrease the risk of asthma in a population with family history of rhinitis or eczema [1, 2]. Studies in mouse models have shown that BCG immunization inhibits the development of asthma features in ovalbumin (OVA)-sensitized mice [3-8]. Several authors reported a switch toward a Th1 pattern of response [3, 5, 9], with little evidence of increased regulatory cell activity, since production of the regulatory cytokine IL-10 as well as production of IL-4 was suppressed [3, 9].

Originally named IFN- γ -inducing factor, IL-18 is important for the generation of protective immunity to mycobacteria. IL-18-deficient mice infected with *Mycobacterium tuberculosis* and *Mycobacterium bovis* BCG developed increased granulomatous lesions while IFN- γ production decreased [10]. IL-18 role on allergic inflammation is complex, as it can promote both Th1 and Th2 responses depending on its environment, the time of its administration and/or the response assessment. IL-18 deficiency selectively enhances OVA-induced eosinophilia in mice, whereas IL-18 gene transfer by adenovirus into the respiratory tract at the time of OVA airway challenge prevents the development of AHR and decreases allergen-specific IL-4 production, airway eosinophilia, mucus production and increases IFN- γ production [11, 12]. In contrast, high doses of IL-18 administered alone or concomitantly with the antigen have been shown to increase allergic sensitization including serum IgE, Th2 cytokines and airway eosinophilia, especially 3 weeks after IL-18 injection [13]. Intraperitoneal coadministration of IL-18 and IL-12 or IL-18 and IL-2 during the airway challenge period inhibits airway hyperresponsiveness (AHR), eosinophilia and serum IgE

levels [14, 15], whilst intranasal of IL-18 plus IL-2 induced allergic airway disease in naïve mice [16].

A BCG producing IL-18 was shown to exhibit enhanced antimicrobial immunity compared to non recombinant BCG due to increased production of IFN- γ [17]. Therefore we constructed a recombinant BCG producing IL-18 (BCG-IL-18) in order to increase the efficiency of intraperitoneally injected BCG in preventing allergic reactions. This strain produces low doses of IL-18 and increases Th1 response [18]. We previously showed that i.p. BCG-IL-18 immunization at the time of OVA-sensitization decreases bronchoalveolar lavage (BAL) eosinophilia, IL-5 production by lymph node cells, but increases IFN- γ production by lymph node cells [19]. The aim of the present work was firstly to evaluate the effect of non-recombinant BCG and BCG-IL-18 immunization on AHR, and secondly to analyze BCG and BCG-IL-18 effects in the long term, i.e. after a secondary challenge occurring 6 weeks after the primary OVA challenge. We showed that both BCG and BCG-IL-18 were able to similarly prevent AHR and airway inflammation (BAL eosinophilia, Th2 cytokine production in BAL and mucus production) after primary and secondary challenges.

Materials and Methods

Sensitization and airway challenge for acute and chronic inflammatory reaction

10-12 week old female BALB/c mice (*H-2^d*) (Iffa Credo, l'Arbresle, France) were sensitized by intraperitoneal (i.p.) injection of 20 µg ovalbumin (OVA) (Grade V, Sigma Chemical Co., St. Louis, MO) emulsified in 2.25 mg aluminum hydroxide (AlumImject: Pierce, Rockford, IL) in a total volume of 100 µl on days (D)1 and 10. Mice were challenged (20 min) via the airways with OVA (1% in saline) for 3 days (D20, 21 and 22) using ultrasonic nebulization. Mice were divided into 2 groups: the groups of OVA-sensitized mice were i.p. injected with OVA and received OVA challenges aerosols. The groups of non-sensitized mice only received OVA-challenges by aerosols.

In the primary challenge protocol, 48 hours after the last OVA challenge (D24), AHR and tissues were assessed. For the secondary challenge protocol, 6 weeks after the first challenge, mice were exposed to OVA challenges (1% in saline on D62, 5% in saline on D63), and airway reactivity and tissues were assessed on D64. Non-sensitized control groups only received OVA-challenges by aerosols.

Immunization with BCG or IL-18-producing recombinant BCG

Non-sensitized and sensitized mice were or were not treated with BCG or BCG-IL-18. *Mycobacterium bovis* BCG (Pasteur strain 1173P2; WHO Stockholm, Sweden) and the recombinant BCG producing IL-18 (BCG-IL-18) [18] were grown, using stationary flasks, at 37°C in Sauton medium containing 10 µg/ml HgCl₂ when required and frozen until use. Non-recombinant BCG or rBCG producing IL-18 were administered intraperitoneally on D0 and D10 (5 X 10⁶ CFU in a final volume of 100 µl).

Airway Hyperresponsiveness

Airway responsiveness was assessed using single-chamber whole body plethysmography (Buxco Electronics, Sharon, CT), as described previously [20]. Enhanced pause (Penh) was used as the measure of airway responsiveness in this study. In the plethysmography, mice were exposed for 2 min to nebulized PBS and followed by increasing concentrations of nebulized methacholine (MCh) (1,5–12 mg/ml in PBS) (Sigma-Aldrich) using an AeroSonic ultrasonic nebulizer (Systam). After each nebulization, recordings were taken for 3 min. The Penh values measured during each 3-min sequence were averaged and were expressed for each MCh concentration as the percentage of baseline Penh values measured after PBS exposure. There were no significant differences in any of the treatment groups for baseline (PBS) Penh.

Bronchoalveolar Lavage

Immediately after assessment of AHR, lungs were lavaged via the tracheal tube with PBS (1x 1ml). Total leukocyte numbers were counted with a hemocytometer. Differential cell counts were performed on bronchoalveolar lavages by counting at least 200 cells on cytocentrifuged preparations, stained with Diff Quick (Dade Behring, Deerfield, IL) and differentiated by standard hematological procedures.

Histochemistry

Lungs were fixed by inflation (1 ml) and immersion in Immunohistofix® (Aphase, Gosselies, Belgium) and embedded in Immunohistowax® (Aphase). For detection of mucus containing cells in fixed airway tissue, 5µm sections were stained with Periodic Acid Schiff (PAS) (Sigma Aldrich, Missouri, USA) and hematoxylin (Labonord, Templemars, France) as previously described [21].

Measurement of cytokines

IFN- γ , IL-5, IL-12 and IL-13 levels in the bronchoalveolar lavage fluid (BALF) were measured by ELISA according to the manufacturer direction (Pharmingen, San Diego, CA, for IFN γ , IL-5, IL-12; R&D systems, Mineapolis, MN, for IL-13). The limits of detection were 4 pg/ml for IL-5, 10 pg/ml for IL-12 and IFN- γ , and 1.5 pg/ml for IL-13.

Measurement of total and OVA-specific serum IgE

Serum levels of total IgE, and OVA-specific IgE were measured by ELISA as previously described [19]. Briefly, 96-well plates (Immulon 2, Dynatech, Chantilly, VA) were coated with either OVA (5 μ g/ml) or purified anti-IgE (clone 02111D, Pharmingen). After addition of serum samples, a biotinylated anti-IgE antibody (clone 02122D, Pharmingen) was used as detecting antibody, and the reaction amplified with avidin-horseradish peroxidase (Sigma). The OVA-specific antibody titers of the samples were related to pooled standards that were generated in the laboratory and expressed as ELISA units per ml (EU/ml). Total IgE levels were calculated by comparison with known mouse IgE standards (Pharmingen). The limit of detection was 100 pg/ml for total IgE.

Statistical analysis

Values for all measurements were expressed as the mean \pm standard error of the mean (SEM). Statistical analyses were done using STATVIEW 5 software. Differences within all groups were first evaluated for each parameter using the Kruskal Wallis test. When statistical significance was observed, differences were subsequently analyzed by using the Mann Whitney test. p values for significance were set to 0.05.

Results

BCG and BCG-IL-18 prevented airway hyperresponsiveness following primary and secondary challenge

OVA-sensitized mice developed airway hyperresponsiveness compared to the non-sensitized mice after primary (figure 1A) and secondary (figure 1B) challenge. Mouse treatment by intraperitoneal injections with BCG or IL-18-producing-BCG (BCG-IL-18) before and during sensitization significantly inhibited airway hyperresponsiveness (figure 1A). In order to evaluate the long term effect of BCG and BCG-IL-18 on AHR, we waited 6 weeks after the primary challenge and then rechallenged the animals. It has previously been shown that at this time point, the inflammatory reaction and AHR developing after primary challenge are resolved but that a secondary airway challenge induces a strong inflammatory reaction with concomitant development of AHR [22]. Under these conditions, BCG and BCG-IL-18 treatment inhibited the increase in airway reactivity (Figure 1B). Baseline airway reactivity was not affected by OVA or BCG or BCG-IL-18 treatment (data not shown).

BCG and BCG-IL-18 decreased BAL inflammation and goblet cell hyperplasia following primary and secondary challenge

In OVA-sensitized mice, inflammatory cell recruitment into the airways was increased after primary as well as secondary airway challenge (Figure 2). Increased total cell numbers in BAL were largely due to increased numbers of eosinophils. There was also a small but significant ($p=0.0005$) increase in the number of lymphocytes compared to the non-sensitized mice (Figure 2). Administration of BCG or BCG-IL-18, led to a significant ($p<0.05$) decrease in total cell numbers, macrophage, lymphocyte and eosinophil numbers in the primary challenge protocol compared to OVA-sensitized mice (Figure 2A). In the secondary challenge

protocol, administration of BCG or BCG-IL-18 significantly decreased eosinophil numbers only (Figure 2B).

Lung tissue was obtained and processed 48 hr after allergen provocation. To assess the degree of goblet cell hyperplasia, tissue sections were stained with Periodic Acid Schiff (PAS). After primary (Figure 3A-D) and secondary (Figure 3E-H) challenge, non-sensitized mice showed no PAS-positive cells, whereas non-treated OVA-sensitized mice showed many PAS-positive cells (92% and 81% of PAS positive cells/mm basement membrane for the primary and secondary challenge protocol respectively). In contrast, sensitized and BCG or BCG-IL-18 treated mice showed only scattered PAS-positive cells (71% and 66% of PAS positive cells/mm basement membrane for BCG treated mice in the primary and secondary challenge protocol respectively and 48% and 40% of PAS positive cells/mm basement membrane for BCG-IL-18 treated mice in the primary and secondary challenge protocol respectively), with a stronger inhibition for BCG-IL-18 treatment after the secondary challenge (Figure 3H).

Effect of BCG and BCG-IL-18 on cytokine production following primary and secondary challenges

BAL fluid was obtained in order to assess Th1 (IFN- γ), pro-Th1 (IL-12), and Th2 (IL-5, IL-13) cytokine levels, 48 or 24 hours after primary or secondary allergen challenge, respectively. Th1 (IFN- γ) and pro-Th1 (IL-12) cytokines were decreased in OVA-sensitized mice compared to non-sensitized mice, significantly after the primary challenge, and non-significantly after the secondary challenge (Figure 4). IL-5 and IL-13 production was significantly increased in OVA-sensitized mice after the primary and the secondary challenge. Treatment of OVA-sensitized mice with BCG or BCG-IL-18 did not significantly affect IFN- γ , IL-12 (Figure 4) or IL-10 (data not shown) production, but significantly inhibited IL-5 and IL-13 production both in primary and secondary challenge protocols.

BCG and BCG-IL-18 treatment did not modify total IgE production but inhibited OVA-specific IgE

Serum from OVA-sensitized mice showed elevated total IgE levels and OVA-specific IgE antibodies compared to non-sensitized mice following the primary and the secondary challenge protocol (Figure 5). Treatment with BCG and BCG-IL-18 did not affect total IgE measured after the primary and the secondary challenge (Figures 5A, 5B). OVA-specific IgE measured after the primary or the secondary challenge were not modified after BCG treatment (Figures 5C, 5D). In contrast, treatment with BCG-IL-18 significantly increased levels of OVA-specific IgE after the primary and decreased them after the secondary challenge compared to OVA-sensitized mice (Figures 5C, 5D).

Discussion

In this study we showed that BCG and BCG-IL-18 given at the time of OVA-sensitization were efficient to prevent AHR even after a secondary challenge performed 6 weeks after the initial allergen challenge. This effect is related to the decrease in the local Th2 immune response, as seen with the decreased BAL IL-5 and IL-13 production. In human as well as in mouse models, Th2 cells are indeed considered as the central cells involved in the development of asthma [23, 24]. Others have shown that BCG was effective in attenuating allergic airway inflammation and associated changes in pulmonary function in animal models. Administered before OVA sensitization, BCG decreases BAL eosinophilia and IL-5 production in the draining lymph nodes in an IFN- γ dependent manner [4]. In our hands, in a previous study, similar effects were found both with intraperitoneal and intranasal administration at the dose of 5×10^6 CFU, when BCG was given at the time of OVA-sensitization [19]. Systemically injected (by intravenous injection) 14 days before the challenge, BCG also suppressed OVA-induced airway responses [9]. In the present study, BCG was injected i.p. at the time of sensitization, and inhibition of airway responses was still observed 10 weeks after the first injection. Only few studies investigated BCG long-term effects: similar results to ours were obtained although BCG was always given long before OVA-sensitization. Given from 4 to 16 weeks before OVA-sensitization by i.n. immunization, BCG prevented airway allergic inflammation [7, 8].

After primary and secondary challenge, BCG and BCG-IL-18 treatment locally decreased the production of IL-5 and IL-13, in parallel to decreased mucus hyperplasia, arguing for a role, in this process, of IL-13, the main cytokine involved in mucus production [25, 26]. However, BCG treatment did not modify IFN- γ and IL-12 production in BAL, in contrast to decreased IFN- γ production seen in the spleen [9] or the draining lymph nodes [19]. This absence of effect on IFN- γ production in BAL may be due to the way of injection (i.p. instead i.n. route

of administration) and the analyzed tissue. Another hypothesis, pointed out by Trujillo and colleagues [27] suggests that the treatment before sensitization may inhibit the onset and consequently the recruitment of Th2 cells thus preventing from allergic reaction. Our previous studies are in favor of this hypothesis as we showed that i.p. administration of BCG or BCG-IL-18 increased *in vitro* production of IFN- γ by splenocytes or lymph node cells, whilst decreasing IL-5 production [18, 19].

In the present work, total IgE production was unchanged after BCG treatment either following primary or secondary challenge, suggesting that BCG was not able to totally prevent Th2 responses. Although *Mycobacterium tuberculosis* has been shown to induce IgE production [28], BCG was shown to prevent OVA-specific IgE production after systemic (intravenous), but not local (i.n.) administration [4, 7, 9]. In our study, BCG does not seem to induce total IgE either as non-sensitized mice treated with BCG-IL-18 did not develop total IgE response, either in the primary or the secondary challenge protocol. However BCG and BCG-IL-18 treatment decreased OVA-specific IgE after the secondary challenge, as well as OVA-specific IgG1 (data not shown). Only BCG-IL-18 treatment effect was significant suggesting a potentiating effect of IL-18 produced by BCG and a long term effect of BCG-IL-18.

BCG-IL-18 did not demonstrate a stronger ability to prevent airway inflammation and hyperresponsiveness than BCG alone, in contrast to what we previously showed [19]. The protocol used to sensitize mice in this first study was much weaker (sensitizing doses were twice lesser and OVA aerosol concentration were 0.1% OVA instead of 1% in the present study) and did not allow the measurement of significant AHR. The absence of a differential effect between BCG and BCG-IL-18 may therefore be related to the strength of the sensitization protocol and the already maximal effect of the non-recombinant BCG on AHR. We previously showed that BCG and BCG-IL-18 were still viable and detected after 94 days. Moreover, although undetectable, IL-18 was efficiently produced *in vivo* by the recombinant

BCG, as suggested by its potentiating effect over non recombinant BCG [18, 19]. In the present study, BCG-IL-18 significantly modified OVA-specific IgE production in contrast to non recombinant BCG, suggesting that IL-18 is really produced *in vivo*, probably in low concentration.

Finally, after the primary and secondary OVA-challenge, BCG or BCG-IL-18 treatment inhibits BAL eosinophilia. BCG was already known to inhibit this feature of asthma when given before sensitization [7]. Interestingly BCG-IL-18 increased the number of BAL lymphocytes compared to BCG-treated mice and non-treated mice. This was not accompanied with decreased or increased airway responses, in contrast to the increased Th2 response induced by IL-18 as previously shown [13]. The only parameter increased after BCG-IL-18 treatment compared to BCG treatment or no treatment was the OVA-specific IgE after the primary challenge. These data suggest a systemic transient effect of IL-18 produced by BCG. IL-18 has been shown previously to increase Th2 responses and more particularly IgE production [13, 29, 30]. The modulation of the production of Th1/Th2 cytokines by IL-18 was suggested to be time dependent [31], which might relate to our observation that OVA-specific IgE transiently increased due to IL-18 produced by i.p. injected BCG. This increase in IgE production was not accompanied by effect on airway inflammation and function. Dissociation of serum IgE and airway inflammation and hyperresponsiveness was previously demonstrated in the same model [32]. Conflicting data have been reported on the role of IL-18 in airway inflammation. In the presence of IL-12, IL-18 administered i.p. before the allergen challenge has been reported to attenuate eosinophil accumulation [14]. OVA-sensitized IL-18 deficient mice exhibit greater BAL eosinophilia than wild-type mice [11]. In contrast, local instillation of recombinant IL-18 was reported to favor the eosinophil recruitment through the release of eotaxin from bronchial epithelial cells without any effect on AHR [33]. Moreover i.p. injection of large quantities of IL-18, in the absence of IL-12, also increases the recruitment of

eosinophils in the airways [34]. Finally, lower doses of IL-18 regularly administered intraperitoneally increase eosinophil recruitment in the airways 3 weeks after the last injection [13]. BCG-IL-18 administered in our study produces only low doses of IL-18 as previously demonstrated [18] and no IL-18 could be detected in BAL (data not shown). Moreover, BCG has been shown to favor IL-12 production by macrophages [6]. The low dose of IL-18 released and the cytokine environment may explain the absence of an increased AHR and eosinophilia in BCG-IL-18 treated mice even after OVA restimulation.

In conclusion, we showed that BCG as well as BCG producing IL-18 administered at the time of sensitization prevents airway responses after primary as well as secondary OVA-challenge, suggesting their effect after systemic injection, which had not been demonstrated yet. Moreover, our study suggests that, thank to the environment induced by BCG, low doses of IL-18 do not increase Th2 responses, even after secondary allergen challenge.

Figure legends

Figure 1: BCG and BCG-IL-18 prevent the development of AHR

OVA-sensitized mice were OVA-sensitized and challenged, whereas non-sensitized mice were challenged only. At the time of each OVA sensitization, treated mice were injected either with BCG or BCG-IL-18. Airway reactivity was evaluated after inhalation of increasing doses of metacholine by whole body plethysmography and penh (enhanced pause) measurement after primary (A) or secondary (B) challenge. Results are expressed as mean \pm sem of penh percentage increase above PBS inhalation. Statistical analysis were performed and showed $p < 0.04$ for differences between OVA-challenged and OVA-sensitized and challenged mice (*), between OVA-sensitized and challenged mice and OVA-sensitized and challenged mice + BCG treatment (#), and between OVA-sensitized and challenged mice and OVA-sensitized and challenged mice + BCG-IL-18 treatment (&).

Figure 2: BCG and BCG-IL-18 effect on BAL composition after primary and secondary challenge.

BAL infiltrate was analyzed after primary (A) or secondary (B) challenge. Results are expressed as mean \pm sem of cell number. *: $p < 0.05$.

Figure 3: BCG and BCG-IL-18 effect on mucus hyperplasia after primary and secondary challenge.

Mucus production was detected using Periodic Acid Schiff staining on lung section from non-sensitized mice (A, E), sensitized mice (B, F), sensitized mice treated with BCG (C, G) or BCG-IL-18 (D, H), after primary (A-D) or secondary (E-H) OVA-challenge.

Figure 4: BCG and BCG-IL-18 effect on BAL cytokine production after primary and secondary challenge.

BAL cytokines were measured after primary (A) or secondary (B) challenge. Results are expressed as mean \pm sem pg/ml. *: $p < 0.03$.

Figure 5: BCG and BCG-IL-18 decrease OVA-specific IgE, but do not modify serum total IgE.

IgE were measured after primary (A, C) or secondary (B, D) challenge. Results are expressed as mean \pm sem ng/ml for total IgE (A, B), and mean \pm sem ELISA units/ml (EU/ml) for OVA-specific IgE (C, D). *: $p < 0.01$.

Aknowledgments

This work was supported by INSERM. L. Amniai is supported by a grant from Ministère de l'Education et de la Recherche. C. Duez was supported by a grant from the Société de Pneumologie de Langue Française and a grant from the Société Française d'Allergologie et d'Immunologie Clinique. BCG and BCG-IL-18 construction work was supported by Institut Pasteur de Lille, INSERM, Région Nord-Pas de Calais and a EC grant (QLRT-PL 1999-01093). The authors wish to thank the Dr A. Tsicopoulos and Dr P. Gosset for critical reviewing of the manuscript.

References

- [1] G. B. Marks, K. Ng, J. Zhou, et al., "The effect of neonatal BCG vaccination on atopy and asthma at age 7 to 14 years: an historical cohort study in a community with a very low prevalence of tuberculosis infection and a high prevalence of atopic disease," *J Allergy Clin Immunol*, vol. 111, 3, pp. 541-549, 2003.
- [2] S. S. da Cunha, A. A. Cruz, I. Dourado, et al., "Lower prevalence of reported asthma in adolescents with symptoms of rhinitis that received neonatal BCG," *Allergy*, vol. 59, 8, pp. 857-862, 2004.
- [3] I. S. Choi, X. H. Lin, Y. A. Koh, et al., "Strain-dependent suppressive effects of BCG vaccination on asthmatic reactions in BALB/c mice," *Ann Allergy Asthma Immunol*, vol. 95, 6, pp. 571-8, 2005.
- [4] K. J. Erb, J. W. Holloway, A. Sobeck, et al., "Infection of mice with *Mycobacterium bovis*-Bacillus Calmette-Guerin (BCG) suppresses allergen-induced airway eosinophilia," *J Exp Med*, vol. 187, 4, pp. 561-569, 1998.
- [5] M. T. Hopfenspirger and D. K. Agrawal, "Airway hyperresponsiveness, late allergic response, and eosinophilia are reversed with mycobacterial antigens in ovalbumin-prensensitized mice," *J Immunol*, vol. 168, 5, pp. 2516-2522, 2002.
- [6] C. Hubeau, M. Singer, M. Lagranderie, et al., "Extended freeze-dried *Mycobacterium bovis* Bacillus Calmette-Guerin induces the release of interleukin-12 but not tumour necrosis factor-alpha by alveolar macrophages, both in vitro and in vivo," *Clin Exp Allergy*, vol. 33, 3, pp. 386-393, 2003.
- [7] T. Major, G. Wohlleben, B. Reibetanz, et al., "Application of heat killed *Mycobacterium bovis*-BCG into the lung inhibits the development of allergen-induced Th2 responses," *Vaccine*, vol. 20, 11-12, pp. 1532-1540, 2002.

- [8] M. A. Nahori, M. Lagranderie, J. Lefort, et al., "Effects of Mycobacterium bovis BCG on the development of allergic inflammation and bronchial hyperresponsiveness in hyper-IgE BP2 mice vaccinated as newborns," *Vaccine*, vol. 19, 11-12, pp. 1484-1495, 2001.
- [9] U. Herz, K. Gerhold, C. Gruber, et al., "BCG infection suppresses allergic sensitization and development of increased airway reactivity in an animal model," *J Allergy Clin Immunol*, vol. 102, 5, pp. 867-874, 1998.
- [10] I. Sugawara, H. Yamada, H. Kaneko, et al., "Role of interleukin-18 (IL-18) in mycobacterial infection in IL-18-gene-disrupted mice," *Infect Immun*, vol. 67, 5, pp. 2585, 1999.
- [11] T. Kodama, T. Matsuyama, K. Kuribayashi, et al., "IL-18 deficiency selectively enhances allergen-induced eosinophilia in mice," *J Allergy Clin Immunol*, vol. 105, 1 Pt 1, pp. 45-53, 2000.
- [12] D. M. Walter, C. P. Wong, R. H. DeKruyff, et al., "IL-18 gene transfer by adenovirus prevents the development of and reverses established allergen-induced airway hyperreactivity," *J Immunol*, vol. 166, 10, pp. 6392-6398, 2001.
- [13] J. S. Wild, A. Sigounas, N. Sur, et al., "IFN-gamma-inducing factor (IL-18) increases allergic sensitization, serum IgE, Th2 cytokines, and airway eosinophilia in a mouse model of allergic asthma," *J Immunol*, vol. 164, 5, pp. 2701-2710, 2000.
- [14] C. L. Hofstra, I. Van Ark, G. Hofman, et al., "Prevention of Th2-like cell responses by coadministration of IL-12 and IL-18 is associated with inhibition of antigen-induced airway hyperresponsiveness, eosinophilia, and serum IgE levels," *J Immunol*, vol. 161, 9, pp. 5054-5060, 1998.
- [15] S. Matsubara, K. Takeda, T. Kodama, et al., "IL-2 and IL-18 attenuation of airway hyperresponsiveness requires STAT4, IFN-gamma, and natural killer cells," *Am J Respir Cell Mol Biol*, vol. 36, 3, pp. 324-332, 2007.

- [16] Y. Ishikawa, T. Yoshimoto and K. Nakanishi, "Contribution of IL-18-induced innate T cell activation to airway inflammation with mucus hypersecretion and airway hyperresponsiveness," *Int Immunol*, vol. 18, 6, pp. 847-55, 2006.
- [17] Y. Luo, H. Yamada, X. Chen, et al., "Recombinant *Mycobacterium bovis* bacillus Calmette-Guerin (BCG) expressing mouse IL-18 augments Th1 immunity and macrophage cytotoxicity," *Clin Exp Immunol*, vol. 137, 1, pp. 24-34, 2004.
- [18] F. Biet, L. Kremer, I. Wolowczuk, et al., "Mycobacterium bovis BCG producing IL-18 increases antigen-specific IFN- γ production in mice," *Infect. Immun*, vol. 70, 12, pp. 6549-6557, 2002.
- [19] F. Biet, C. Duez, L. Kremer, et al., "Recombinant *Mycobacterium bovis* BCG producing IL-18 reduces IL-5 production and bronchoalveolar eosinophilia induced by an allergic reaction," *Allergy*, vol. 60, 8, pp. 1065-72, 2005.
- [20] E. Hamelmann, J. Schwarze, K. Takeda, et al., "Noninvasive measurement of airway responsiveness in allergic mice using barometric plethysmography," *Am J Respir Crit Care Med*, vol. 156, pp. 766-775, 1997.
- [21] A. Tomkinson, G. Cieslewicz, C. Duez, et al., "Temporal association between airway hyperresponsiveness and airway eosinophilia in ovalbumin-sensitized mice," *Am J Respir Crit Care Med*, vol. 163, 3 Pt 1, pp. 721-30, 2001.
- [22] A. Kanehiro, T. Ikemura, M. J. Makela, et al., "Inhibition of phosphodiesterase 4 attenuates airway hyperresponsiveness and airway inflammation in a model of secondary allergen challenge," *Am J Respir Crit Care Med*, vol. 163, 1, pp. 173-184, 2001.
- [23] P. J. Barnes, "Cytokine modulators for allergic diseases," *Curr Opin Allergy Clin Immunol*, vol. 1, 6, pp. 555-560, 2001.
- [24] M. Larche, D. S. Robinson and A. B. Kay, "The role of T lymphocytes in the pathogenesis of asthma," *J Allergy Clin Immunol*, vol. 111, 3, pp. 450-463, 2003.

- [25] G. Grunig, M. Warnock, A. E. Wakil, et al., "Requirement for IL-13 independently of IL-4 in experimental asthma," *Science*, vol. 282, 5397, pp. 2261-2263, 1998.
- [26] M. Wills-Karp, J. Luyimbazi, X. Xu, et al., "Interleukin-13: central mediator of allergic asthma," *Science*, vol. 282, 2258-2261, pp. 1998.
- [27] Trujillo-Vargas CM, Mayer KD, Bickert T, et al., "Vaccinations with T-helper type 1 directing adjuvants have different suppressive effects on the development of allergen-induced T-helper type 2 responses," *Clin Exp Allergy*, vol. 35, 8, pp. 1003-1013, 2005.
- [28] A. J. Yong, J. M. Grange, R. D. Tee, et al., "Total and anti-mycobacterial IgE levels in serum from patients with tuberculosis and leprosy," *Tubercle*, vol. 70, 4, pp. 273-279, 1989.
- [29] T. Hoshino, H. Yagita, J. R. Ortaldo, et al., "In vivo administration of IL-18 can induce IgE production through Th2 cytokine induction and up-regulation of CD40 ligand (CD154) expression on CD4+ T cells," *Eur J Immunol*, vol. 30, 7, pp. 1998-2006, 2000.
- [30] T. Yoshimoto, B. Min, T. Sugimoto, et al., "Nonredundant roles for CD1d-restricted natural killer T cells and conventional CD4+ T cells in the induction of immunoglobulin E antibodies in response to interleukin 18 treatment of mice," *J Exp Med*, vol. 197, 8, pp. 997-1005, 2003.
- [31] T. Uchida, M. Kinoshita, M. Fukasawa, et al., "IL-18 time-dependently modulates Th1/Th2 cytokine production by ligand-activated NKT cells," *Eur J Immunol*, vol. 37, 4, pp. 966-77, 2007.
- [32] E. Hamelmann, K. Takeda, J. Schwarze, et al., "Development of eosinophilic airway inflammation and airway hyperresponsiveness requires interleukin-5 but not immunoglobulin E or B lymphocytes," *Am J Respir Cell Mol Biol*, vol. 21, 4, pp. 480-9, 1999.
- [33] E. Campbell, S. L. Kunkel, R. M. Strieter, et al., "Differential roles of IL-18 in allergic airway disease: induction of eotaxin by resident cell populations exacerbates eosinophil accumulation," *J Immunol*, vol. 164, 2, pp. 1096-1102, 2000.

[34] K. Kumano, A. Nakao, H. Nakajima, et al., "Interleukin-18 enhances antigen-induced eosinophil recruitment into the mouse airways," *Am J Respir Crit Care Med*, vol. 160, 3, pp. 873-878, 1999.

Figure 1




Figure 2




Figure 3




Figure 4




Figure 5

