

Sickness absence for upper limb disorders in a French company.

Kayigan Wilson d'Almeida, Catherine Godard, Annette Leclerc, Gérard Lahon

► **To cite this version:**

Kayigan Wilson d'Almeida, Catherine Godard, Annette Leclerc, Gérard Lahon. Sickness absence for upper limb disorders in a French company.. Occupational Medicine, Oxford University Press (OUP), 2008, 58 (7), pp.506-8. 10.1093/occmed/kqn084 . inserm-00309077

HAL Id: inserm-00309077

<https://www.hal.inserm.fr/inserm-00309077>

Submitted on 5 Aug 2009

HAL is a multi-disciplinary open access archive for the deposit and dissemination of scientific research documents, whether they are published or not. The documents may come from teaching and research institutions in France or abroad, or from public or private research centers.

L'archive ouverte pluridisciplinaire **HAL**, est destinée au dépôt et à la diffusion de documents scientifiques de niveau recherche, publiés ou non, émanant des établissements d'enseignement et de recherche français ou étrangers, des laboratoires publics ou privés.

Sickness absence for upper limb disorders in a French company

D'Almeida Kayigan Wilson ^{1,2*}, Godard Catherine ², Leclerc Annette ¹, Lahon Gérard ²

¹ Santé publique et épidémiologie des déterminants professionnels et sociaux de la santé INSERM : U687, IFR69, Université Paris Sud - Paris XI, Université de Versailles-Saint Quentin en Yvelines, Hôpital Paul Brousse 16, av Paul Vaillant Couturier 94807 VILLEJUIF,FR

² Service Général de Médecine de Contrôle EDF, 22-28 rue de Joubert, 75009 Paris,FR

* Correspondence should be addressed to: Kayigan Wilson D'Almeida <kayialmeida@hotmail.com>

MESH Keywords Adult ; Female ; France ; epidemiology ; Humans ; Male ; Middle Aged ; Musculoskeletal Diseases ; epidemiology ; Occupational Diseases ; epidemiology ; Shoulder Impingement Syndrome ; epidemiology ; Sick Leave ; statistics & numerical data ; Socioeconomic Factors ; Upper Extremity ; Young Adult

INTRODUCTION

Musculoskeletal disorders (MSDs) are an important public health issue because of their socio-economic consequences.[1–4]. In recent decades, research have helped to improve epidemiological knowledge and management of MSDs, particularly of the upper limb (MSDUL).

Less is known however about their impact on sick leave in industry. Studies focusing on workers compensation include only a portion of workers [5], because sick leave without claims for compensation for work-related injuries is frequent in countries such as France, where workers are compensated for disability leave regardless of whether the health problem is work-related.

Our objectives were to describe the sickness absence episodes for MSDUL in the French national power and gas company -EDF and Gaz de France (EDF-GDF)- and to study the associations with age, gender, and socio-economic status.

METHODS

Study population

The study population consisted of about 140 000 workers from EDF-GDF. EDF-GDF has its own medical insurance system handled by a department called Service Général de Médecine de Contrôle (SGMC). A specificity of this medical system is compulsory verification of any sick time off. All employees with sick leave must see a physician of SGMC's medical department, who codes diagnoses according to a simplified version of the International Classification of Diseases of the World Health Organization. The information is then recorded in an epidemiology registry. The items available in that database include absences with diagnosis and duration, and the worker's profile (age, gender, occupation, industrial sector).

Data collection and methods

The database of sick leave from January 2000 to December 2004 was reviewed to identify the absences that might be due to MSDUL. Long-term sick leave (one year or more) were excluded as available data suggested that an absence for MSDUL would rarely be so long. We anticipated that MSDULs would be coded as disease of the locomotor system, aponeuroses and tendons; arthroses or diseases of the nervous system not due to alcohol. A preliminary selection of these categories found 21 890 sickness absence episodes possibly due to MSDUL. For each, a questionnaire was sent to the company physician, who checked whether it was a MSDUL and coded the diagnoses according to a list derived from a European consensus [6]. Finally, 5543 sickness absence episodes were classified as MSDUL.

Annual incidence rates for sick leave due to MSDUL were calculated by gender, age, diagnosis, and socio-economic status (SES) coded according to the French classification [7]. The number of workers with at least one absence due to MSDUL was divided by the number of full-time equivalent workers aged 20–59 years in the corresponding category. Incidence rates according to SES were calculated only for 2004 since this information was not available for the other years.

The number of work days lost per diagnosis was calculated.

Differences in rates were tested using a chi square test. P-values below 0.05 were considered statistically significant throughout the analysis.

This study was made in accordance with ethical requirements for epidemiologic research in France.

RESULTS

Among the 5543 sickness absence episodes, rotator cuff syndrome was diagnosed most frequently, accounting for 31% of absences (Table 1). Other common diagnoses were carpal tunnel syndrome (16%), epicondylitis (14%), and neck MSD (11%). Less frequent

diagnoses included extensor/flexor tendonitis and tenosynovitis (8%), elbow, wrist and fingers arthritis (3%), cubital tunnel syndrome (2%), De Quervain's disease (1%), Guyon's canal syndrome (0.6%), radial tunnel syndrome (0.4%) and neuropathy caused by vibrations (0.2%). "Other" MSDUL, coded as such because they did not match any of the diagnoses proposed in the questionnaire, accounted for 13%.

Among the frequent diagnoses, carpal tunnel syndrome and rotator cuff syndrome were associated with the longest sick leave durations (36.7 and 30.7 days respectively). Longer absences were found, however, for less frequent diagnoses, such as Guyon's canal syndrome (55.3 days).

Most of the workers (77%) had a single sickness absence episode for MSDUL in the study period. For the others, the absences were most often several episodes with the same diagnosis.

The incidence rate of sickness absence spells due to MSDUL increased from 5.3 in 2000 to 6.3 per 1000 person-years in 2004.

Sick leave for MSDUL increased with age and incidence was higher for women than for men (6.5 and 5.7 per 1000 person-years respectively, $P < 0.01$).

Sick leave due to MSDUL was associated with SES ($P < 0.001$). Office and blue-collar workers had a higher risk than managers and professionals or intermediate white-collar occupations (Figure 1).

DISCUSSION

The study's weaknesses are its retrospective design and the absence of a standardized clinical examination. Moreover, the shortest absence episodes were not always seen by the physicians (in 2006, 28% of episodes ≤ 7 days were not checked). Nonetheless, probably few sick leaves due to MSDUL were missed, since their duration was usually longer (Table 1).

The higher incidence of sickness absence episodes for MSDUL among women, the increase with age and the association with SES are consistent with the literature.[8,9].

The annual incidence of sick leave due to MSDUL was about 6 per 1000 person-years. This result cannot be compared to another one since there is no publication reporting incidence of sick leave for MSDUL [10]. Absenteeism cannot be a surrogate for disease burden or incidence as it depends on attitudes to sick leave, which has many determinants; suffering from MSDUL does not necessarily induce sick leave. Nevertheless, absenteeism is one dimension of the severity of the disorders, a useful indicator for the company and an indicator of the societal cost.

CONCLUSION

The most common diagnoses of MSDUL in sick leaves were rotator cuff syndrome, carpal tunnel syndrome, epicondylitis and neck MSD. Sick leave due to MSDUL was associated with socio-economic status, women had a higher risk and the incidence increased with age. Absenteeism could be a useful indicator in the prevention of MSDUL, because it identifies the most disabling diagnoses as shoulder disorders and shows that older workers, especially manual and office workers, are those most at risk.

References:

1. Morse TF, Dillon C, Warren N, Levenstein C, Warren A The economic and social consequences of work-related musculoskeletal disorders: the Connecticut Upper-Limb Surveillance Project (CUSP). *Int J Occup Environ Health*. 1998; 4: 209- 216
2. Thiehoff R Economic significance of work disability caused by musculoskeletal disorders. *Orthopade*. 2002; 31: 949- 956
3. Roux CH, Guillemin F, Boini S Impact of musculoskeletal disorders on quality of life: an inception cohort study. *Ann Rheum Dis*. 2005; 64: 606- 611
4. Morken T, Riise T, Moen B Frequent musculoskeletal symptoms and reduced health-related quality of life among industrial workers. *Occup Med*. 2002; 52: (2) 91- 98
5. Hashemi L, Webster BS, Clancy EA, Courtney TK Length of disability and cost of work-related musculoskeletal disorders of the upper limb. *J Occup Environ Med*. 1998; 40: 261- 269
6. Sluiter JK, Rest KM, Frings-Dresen MH Criteria document for evaluating the work-relatedness of upper-limb musculoskeletal disorders. *Scand J Work Environ Health*. 2001; 27: (Suppl 1) 1- 102
7. Institut National de la Statistique et des Etudes Economiques (INSEE).
8. Wijnhoven HA, de Vet HC, Picavet HS Prevalence of musculoskeletal disorders is systematically higher in women than in men. *Clin J Pain*. 2006; 22: 717- 724
9. Melchior M, Krieger N, Kawachi I Work factors and occupational class disparities in sickness absence: findings from the GAZEL cohort study. *Am J Public Health*. 2005; 95: 1206- 1212
10. Huisstede BM, Bierma-Zeinstra SM, Koes BW, Verhaar JA Incidence and prevalence of upper-extremity musculoskeletal disorders. A systematic appraisal of the literature. *BMC Musculoskelet Disord*. 2006; Jan 31 7: 7-

Figure 1

Annual incidence of sickness absence episodes due to MSDUL according to the socio-economic status (incidence per 1000 person-years, year 2004)




Table 1Diagnoses associated with sickness absence for MSDUL (¹) and average duration of the sickness absences in the period [2000–2004]

Diagnosis	Number of sickness absences	Average duration of a sickness absence (days)
Rotator cuff syndrome	1722	30.77
Carpal tunnel syndrome	866	36.74
Epicondylitis	795	20.83
Neck musculoskeletal disorders	598	16.37
Extensor/tensor tendonitis and tenosynovitis	452	23.44
Elbow, wrist and fingers arthrosis	173	31.86
Cubital tunnel syndrome	94	42.68
De Quervain's disease	61	19.82
Guyon's canal syndrome	31	55.30
Radial tunnel syndrome	22	27.81
Neuropathy caused by vibrations	9	39.67

¹ limited to the diagnoses proposed in the questionnaire