

# Studying Receptor-mediated cell adhesion at the single molecule level

Anne Pierres, Anne-Marie Benoliel, Pierre Bongrand

► **To cite this version:**

Anne Pierres, Anne-Marie Benoliel, Pierre Bongrand. Studying Receptor-mediated cell adhesion at the single molecule level: Surface-bound adhesion receptors. *Cell Communication and Adhesion*, Taylor & Francis, 1998, 5, pp.375. inserm-00358477

**HAL Id: inserm-00358477**

**<https://www.hal.inserm.fr/inserm-00358477>**

Submitted on 3 Feb 2009

**HAL** is a multi-disciplinary open access archive for the deposit and dissemination of scientific research documents, whether they are published or not. The documents may come from teaching and research institutions in France or abroad, or from public or private research centers.

L'archive ouverte pluridisciplinaire **HAL**, est destinée au dépôt et à la diffusion de documents scientifiques de niveau recherche, publiés ou non, émanant des établissements d'enseignement et de recherche français ou étrangers, des laboratoires publics ou privés.

# ***Studying receptor-mediated cell adhesion at the single molecule level***

Pierres Anne , Benoliel Anne-Marie , Bongrand Pierre \*

*Adhésion cellulaire INSERM : U387, FR*

\* Correspondence should be addressed to: Pierre Bongrand < pierre.bongrand@inserm.fr >

## **Abstract**

### **Summary**

Cell adhesion is essentially mediated by specific interactions between membrane receptors and ligands. It is now apparent that the mere knowledge of the on- and off-rate of association of soluble forms of these receptors and ligands is not sufficient to yield accurate prediction of cell adhesive behavior. During the last few years, a variety of complementary techniques relying on the use of hydrodynamic flow, atomic force microscopy, surface forces apparatus or soft vesicles yielded accurate information on i) the dependence of the lifetime of individual bonds on applied forces and ii) the distance dependence of the association rate of bound receptors and ligands. The purpose of this review is, first to recall the physical significance of these parameters, and second to describe newly obtained results. It is emphasized that molecular size and flexibility may be a major determinant of the efficiency of receptor mediated adhesion, and this cannot be studied by conventional methods dealing with soluble molecules.

**MESH Keywords** Animals ; Cell Adhesion ; physiology ; Cell Adhesion Molecules ; chemistry ; Cells ; chemistry ; cytology ; Humans ; Microscopy, Atomic Force

**Author Keywords** Cell Adhesion ; Intermolecular forces ; Flow chamber ; Surface forces apparatus ; atomic force microscopy ; biomembrane probe

## **INTRODUCTION**

Cell adhesion is usually mediated by specific interactions between dedicated membrane receptors and ligands. The study of numerous biological systems revealed that the outcome of intercellular contacts is not entirely accounted for by the affinity between interacting adhesion molecules. Here are a few representative examples :

Inflammation is a ubiquitous process of prominent pathological significance. An early step is the initial interaction between flowing leukocytes and activated endothelial cells expressing adhesion molecules called selectins. These molecules have a unique capacity to bind cells passing by with a velocity of several hundreds of micrometers per second and make them roll on the vessel wall with a hundredfold slower pace. An attractive hypothesis would be that the functional properties of selectins be linked to i) particularly high rates of association and dissociation (Lawrence and Springer, 1991), ii) high tensile strength (Alon et al., 1995), and iii) capacity of forming adhesion while cell membranes are maintained at relatively high distance by repulsive surface molecules (Patel et al., 1995). Clearly, it is of interest to test this hypothesis by measuring aforementioned parameters (Kaplanski et al., 1993 ; Alon et al., 1995).

Migration on a receptor-bearing surface is another important cellular process : although different cell populations may display varying locomotory behavior, displacement often involves the forward emission of protrusions such as lamellipodia, with subsequent adhesion allowing the cell to contract and detach its rear part from the substratum (Stossel, 1993). Understanding these phenomena at the molecular level requires accurate knowledge of the behavior of adhesion molecules : efficient anchoring of the anterior protrusion requires that a sufficient number of bonds be formed to stand tensile forces of several hundreds of piconewtons (Oliver et al., 1994). Detachment of the posterior part is also highly dependent on the mechanical properties of adhesion molecules (Crowley and Horwitz, 1995).

Many adhesion receptors such as selectins (Ushiyama et al., 1993), integrins (Lollo et al., 1993) or members of the immunoglobulin superfamily (van der Merwe et al., 1993) were produced in soluble form and assayed for estimation of the kinetic or equilibrium constants of association with their ligands. However, while these parameters usually provided a satisfactory description of the interaction between soluble molecules, or between a soluble molecular species and surface-bound receptors, it was rapidly clear that more sophisticated tools were required to account for the association between surface-bound molecules. Indeed, as emphasized by Pierres et al. (1996a), the rate of bond formation between membrane-bound molecules is a function of the distance  $d$  between the anchoring points of these molecules. Further, the rate of bond dissociation between surface-bound receptors is usually dependent on the distractive force  $F$  applied on these molecules. Also, the experimental rate of separation of surfaces bound by a ligand-receptor couple is dependent on both the applied force and the motion of surfaces following bond rupture.

Until recently, no experimental approach allowed a direct determination of aforementioned functions. Indeed, standard techniques such as equilibrium dialysis (e.g. Kabat, 1968) or more refined methods such as plasmon resonance based technology (van der Merwe et al., 1993)


could not be applied to bound molecules. Also, although some theoretical models of cell adhesion provided a link between association rates and measurable aspects of cell behavior such as contact areas (Bell et al., 1984), adhesion efficiency (Hammer and Lauffenburger, 1987) or binding strength (Dembo et al., 1988), they relied on too many unknown parameters or untested assumptions to allow a safe derivation of molecular properties. This situation emphasizes the remarkable interest of simple theoretical models elaborated by Bell (1978) who provided a fairly simple link between the behavior of soluble and bound adhesion molecules, and suggested some testable predictions concerning the mechanical strength of molecular bonds. Thus, he estimated at a few tens of piconewtons the force required to rapidly break a single interaction comparable to an antigen-antibody bond.

Remarkably, within a few years, a variety of experimental approaches allowed fairly direct test of Bell's model. Indeed, experiments based on hydrodynamic forces (Tha et al., 1986), soft vesicles used as tunable transducers (Evans et al., 1991), or atomic force microscopy (Florin et al., 1994) demonstrated that the strength of many ligand-receptor interactions was of the order of several tens of piconewtons (or slightly higher values when the high affinity interaction between avidin and biotin was studied). A common feature of these techniques was to study a few or even individual bonds, which alleviated many difficulties hampering the interpretation of previous experiments. Other authors reported on the distance dependence of the energy of interaction between receptors and ligands (Helm et al., 1991), the natural lifetime of weakly stressed bonds (Kaplanski et al., 1993), the precise dependence of the bond lifetime on applied force (Alon et al., 1995) or the distance dependence of association rate (Pierres et al., 1997).

The aim of the present review is therefore twofold : first, we shall discuss the significance of the physical parameters we need define to achieve a satisfactory description of bonding behavior. Second, we shall review recent methodological advances allowing direct measurement of these parameters. For the sake of clarity, we shall sequentially discuss the processes of bond formation and dissociation.

## RATE OF BOND FORMATION

We shall start by following the simple approach used by George Bell (1978). In a solution, the formation of a bond between two molecules A and B is conceptually separated into sequential steps (Figure 1) :



The first step is the formation of a so-called encounter complex between A and B : these molecules are brought into binding distance through mere diffusion. This distance will be denoted as  $R_{AB}$ . It is expected to be close to the sum of the radii of molecules A and B if these are modeled as spheres.

The second step is bond formation. This is much more complicated, since it requires suitable rotation in order that reactive sites come in close contact ; then the energy barrier separating free and bound states must be overcome by thermal fluctuations.

Although the rate of formation of the bound species C can be calculated analytically with respect to the parameters defined in (1), the exact formula is somewhat awkward. Thus, Bell took advantage of a widely used procedure consisting of considering a steady state phase, where the concentrations of molecules A and B as well as the intermediate complex AB may be considered as fairly constant. He obtained :

$$\frac{d[C]}{dt} = k_+ [A] [B]$$

$$k_+ = \frac{d_+ r_+}{d_- + r_+}$$

(where the square brackets represent the concentration of any molecular species). There remains to understand the difference between reactions involving free and surface-bound molecules. Bell suggested that the second step might proceed with similar rate between free and bound molecules. Simple estimates based on diffusion equations might thus allow a crude derivation of the rate of association between membrane molecules from the properties of soluble species. This simple view played a major role as a starting point in understanding molecular interactions at the cell surface. However, as will be now discussed, some points must be clarified in order to discuss recent experimental data within the framework of Bell's theory. We shall discuss sequentially the two steps described in equation (1).

### Diffusion phase

First, let us look at an intuitive representation of the steady state phase : the basic assumption is that the average concentration of species B at distance  $R_{AB}$  from molecules A is some constant  $\kappa[B]$ . If the reaction rate  $r_+$  is very high,  $\kappa$  is expected to be close to zero, and the rate

limiting step is the arrival of molecules B in contact with A. The overall reaction is said to be diffusion-limited. On the contrary, if  $r_+$  is much lower than  $d_+$ , constant  $\kappa$  is close to 1. The steadystate reaction rate can be calculated with a theory first elaborated by Smoluchowski (1917). As briefly described in Appendix I, in a three-dimensional medium :

$$d_+ = 4\pi(D_A + D_B)(1 - \kappa)R_{AB}$$

where  $D_A$  and  $D_B$  are the diffusion constants of molecules A and B respectively. Now, if molecules are embedded in a membrane, the surface rate of formation of the encounter complex may be calculated in a similar way, yielding (Appendix I) :

$$d_{+m} = 2\pi(D_{Am} + D_{Bm})(1 - \kappa) / \ln(R_{cell} / R_{AB})$$

where  $R_{cell}$  is the cell radius. Note that this formula may seem at variance with Bell's equation. However, the logarithm is not very different from unity and there is no order-of-magnitude discrepancy between this and Bell's equation.

### Reaction step

The reaction step is conceptually much more complicated than the diffusion phase. It seems reasonable to split this process into a rotation step that is required to bring binding sites in contact, and the intermolecular association that is driven by intermolecular potential (Figure 1).

### Molecular rotation

As pointed out by Bell (1978), the rate constant of some antibody-hapten reactions may be fairly close to the diffusion limit. This may seem somewhat surprising, since it might be thought that only a low proportion of molecular encounters should happen with an orientation compatible with binding (Figure 1). It is therefore of interest to note that there is some evidence supporting the concept that hydrodynamic (Brune and Kim, 1994) or electrodynamic (Helm et al., 1992) forces between "well designed" molecules might induce couples generating rotation conducive to the acquisition of correct orientation during intermolecular approach.

The problem is to know what happens when molecules are bound to a surface. The following two predictions may be safely suggested :

- If molecules are rigid, only a minimal fraction of orientations may be compatible with binding (Figure 2), resulting in a dramatic decrease of the binding rate as compared to free structures.
- If adhesion molecules are flexible, the diffusion of binding sites should not be substantially decreased on a membrane. Indeed, if we model these sites as points located on the surface of a rigid spherical molecule of radius  $a$ , the diffusion component due to translation is expected to be equal to  $kT/6\pi\mu a$ , where  $k$  is Boltzmann's constant,  $T$  is the absolute temperature and  $\mu$  is the medium viscosity, whereas the diffusion component due to rotation (i.e. sphere radius times rotational diffusion coefficient) should be  $kT/8\pi\mu a$  (see e. g. Hill, 1960).

The qualitative conclusion suggested by these remarks is that rigid molecules are not expected to interact if they are bound to surfaces. If they are flexible, binding sites are expected to move with comparable velocity in solution and on a membrane. Therefore, the kinetics of bond formation should be proportional to the fraction of available molecular conformations corresponding to a molecular orientation compatible with binding. As shown on Figure 2, this probability is expected to depend on the distance between the anchoring points of considered molecules.

### Molecular association

A detailed discussion of the general mechanisms of the kinetics of molecular interactions would not fall into the scope of this review. However, molecular flexibility (as well as the flexibility of anchoring to the membranes) is likely to play a dominant role at this stage. Indeed, according to Eyring's celebrated theory of reaction rates (Eyring, 1935 ; Hill, 1960), the absolute rate of molecular association should be proportional to the rate of passage through an "activation state". Both the probability of formation of this state and the transition rate per activated complex between molecules of comparable size should be inversely proportional to a power of at least 1/2 of the molecular mass. Thus, molecules of about 5 nm radius rigidly bound to cells of 5  $\mu$ m radius should display at least  $10^9$  fold (i.e.  $[5\mu\text{m}/5\text{nm}]^3$ ) lower chemical reaction rates than freely moving molecules. On the contrary, if molecules are flexible enough, reactive sites might behave as molecules with an effective mass comparable to the masses of free receptors and ligands.

It is therefore concluded that surface-bound molecules will be able to interact efficiently only if they are fairly flexible. In this case, the diffusive rate of binding sites is comparable to that of free molecules, but this diffusion is constrained within a very small volume. The binding rate should therefore be exquisitely sensitive to the surface distance, with a characteristic range comparable to molecular size. These conclusions were, in fact, intuitively obvious, but they must be borne in mind in order to interpret results obtained with the methods that will be described in the second part of this review. A semi-quantitative model for the distance dependence of bond association is presented in appendix II. Theoretical curves for the distance-dependence of association rate between surface-bound molecules are presented on Figure 3.

## BOND DISSOCIATION

As emphasized by Bell (1978), the spontaneous rupture of a molecular bond requires that i) a sufficient amount of thermal energy be accumulated to reach the transition state and ii) unbound molecules get separated by diffusion.

When at least one of interacting molecular species is in soluble form, it is reasonable to consider that the debonding process is correctly described by a single numerical constant, that is the off-rate ( $k_-$ ), or probability of rupture per unit of time. When both ligands and receptors are bound to surfaces, two important differences must be considered :

- First, the bond may be subjected to some distractive force  $F$ . The off-rate  $k_-$  must therefore be considered as a function of  $F$  (i.e.  $k_-(F)$ ). Bell (1978) suggested to apply to individual bonds an empirical formula that was obtained by studying the tensile strength of macroscopic bodies (Zurkhov, 1956) :

$$k_-(F) = k_-(0) \exp(\gamma F / kT)$$

Bell argued that the constant  $\gamma$  must represent the molecular range of interaction, and he suggested a tentative value of a few tens of nanometers for the antigen-antibody bond. Some theoretical justification for equation (6) may be found within the framework of Eyring's theory (see appendix III and Evans and Ritchie, 1997). As shown below, the conclusion that the tensile strength of molecular bonds should be of order of several tens of piconewtons was fully supported by further experimental studies. More recently Evans et al.. (1991) suggested the following empirical formula :

$$k_-(F) = v_0(F / F_0)^a$$

where  $a$  is an empirical parameter describing the mechanical behavior of the considered bond. The important conclusion is that the off-rate is dependent on applied force, leaving as a challenge to experimentalists an accurate determination of this dependence in several representative models.

- Second, free and bound molecules are expected to display quite different behavior. The problem is to know whether rebinding is likely to occur after bond rupture. Indeed, even if a surface-bound molecule is flexible enough to remove its binding site from a receptor with a time scale comparable to that found with a free molecule, the local concentration of the dissociated binding site will remain quite high (the concentration of a molecule constrained in a volume of 10 nm size is of order of 1.7 mM). Thus, the apparent lifetime of a surface-bound molecule subjected to a tensile strength  $F$  may be dependent on the dynamic properties of the particle, i.e. on its effective mass. Indeed, this mass determines the velocity with which the molecule will go away after detachment. This problem may be important if the reciprocal value of the on-rate (i.e. the binding time) is smaller than the time required for the detached surface to move by a molecular length when it is subjected to a distractive force  $F$ .

## EXPERIMENTAL STUDY OF THE INTERACTION BETWEEN SURFACEBOUND MOLECULES

We shall now review some recent experimental studies on the interactions between individual surface bound molecules. The basic methodologies that yielded most published information during the last years were reviewed in a recent book (Bongrand, Claesson & Curtis, 1994)

### The surface forces apparatus

The surface forces apparatus (Israelachvili and Tabor, 1972 ; Israelachvili and Adams, 1978) has been used for more than twenty years to study short-distance interactions between surfaces. The basic principle was to bring into close distance two crossed cylinders coated with silvered mica surfaces, with simultaneous measurement of intersurface distance and interaction force. Three points may be emphasized ( Israelachvili, 1991):

- The distance between surfaces is measured by means of an interferometric technique allowing better than angström accuracy, and precise determination of surface deformation (Helm et al., 1989). Note that the interest of using mica is due to the smoothness of this material at the atomic scale.

- The force is measured with an accuracy of order of 10 nanonewtons : this is achieved by combining accurate distance measurement, mounting of a cylinder on a soft spring (cantilever), and control of additional displacement with a piezoelectric device.

• When the radius of curvature  $R$  of the cylinders ( $\sim 1\text{cm}$ ) is much higher than the range of the interaction, it may be shown that the measured force  $F$  is related to the interaction energy  $W$  per unit area through the simple formula :

$$W = F / 2\pi R$$

(This is the so-called Derjaguin approximation). The surface forces apparatus (SFA) allowed direct measurement of van der Waals forces in water (Israelachvili and Adams, 1978). It was then applied to surfaces coated with lipid films or macromolecules (see e. g. Luckham and Klein, 1985 ; Claesson, 1994) and yielded very sensitive energy/distance curves. Recently, (Helm et al., 1991 ; Israelachvili et al., 1994), this apparatus was applied to specific interactions between biological molecules. The first model studied was the avidin-biotin association : this was chosen in view of its extremely high affinity. In order to obtain very regular arrays of avidin and biotin binding sites, the authors incorporated these molecules in lipid layers that were deposited on mica surfaces. Other models such as antigen-antibody bonds (Leckband et al., 1995) or nucleic acid interactions (Pincet et al., 1994) were also studied with this technique. The major conclusion was that specific interactions resulted in very strong attraction within a very narrow range of distances (Figure 4 : the peak width was of order of one angström). The peak value of this attraction could in principle allow an absolute determination of the interaction energy, provided the surface density of binding sites was determined. The estimate yielded by equation (8) for the avidinbiotin interaction energy is about 17 kT (Helm et al., 1991 ;  $k$  is Boltzmann's constant and  $T$  is the absolute temperature). The binding constant for the avidin-biotin association in solution is about  $10^{15} \text{ M}^{-1}$ . In a study made on the interaction between fluorescyl groups and a specific antibody (with an affinity of  $5 \times 10^9 \text{ M}^{-1}$ ), assuming a surface area of  $45 \text{ nm}^2$  per antibody site, with 75 % of active sites, the binding energy measured with the surface forces apparatus is about 6 kT (Leckband et al., 1995). Note that the relationship between measured binding energy and affinity constant is not at all straightforward. This point is discussed in appendix IV.

Another important finding (Leckband et al., 1992 & 1995) was that a weak attraction between protein sites was detectable when they were separated by a distance as high as 8.5 nm (Leckband et al., 1992). Thus, as emphasized by the authors, when two adhesion molecules approach each other through diffusion, electrodynamic steering forces may ensure suitable orientation in order that binding sites are facing each other when molecules come in contact. Interestingly, these forces are due to a particular spatial organisation of the molecule rather than the existence of average opposite charges on ligand and receptor molecules, since this force disappeared on protein denaturation (Leckband et al., 1995).

Another interesting finding was that the specific interaction between avidin and biotin molecules could be detected only when they were deposited on fluid, not solid, monolayers, thus emphasizing the importance of a molecular match in short range interactions (see also McGuiggan and Israelachvili, 1990). The importance of molecular length and flexibility is indeed obvious in many biological models.

A final advantage of the surface forces apparatus is that the accurate control of surface distance may yield very safe information on molecular conformation on interfaces. However, there are some basic limitations with this technique : first, since many molecular interactions are averaged, it has not been possible at the present time to derive kinetic parameters of bond formation and dissociation. Second, avidin-biotin interaction was found to be stronger than the cohesive strength of molecular layers, thus hampering precise study of the detachment process. Third, quantitative discrimination between nonspecific van der Waals forces and specific effects may be somewhat complicated. Fourth, it seems difficult to apply this methodology to irregular surfaces such as are found in biological systems. It is therefore interesting to compare the data obtained with this approach to results provided by other techniques. During the last few years, several authors obtained very accurate information on individual ligand-receptor bonds with atomic force microscopy.

### **Atomic force microscopy**

As briefly sketched on Fig. 5, the basic principle of an atomic force microscope (Binnig et al., 1982) is deceptively simple (see e. g. Erlandsson & Olsson, 1994, for a brief description). A very sharp needle (the tip width may be on the nanometer scale) is mounted on a soft cantilever (typically of order of 0.1 N/m stiffness) whose position may be detected with angström accuracy. The tip is moved near a suitable material sample. A piezoelectric device is used to control the tip-to-sample distance. Now, the usual way of using an atomic force microscope (AFM) is to scan the sample by lateral displacement of the tip while the interaction force is maintained constant by a feedback mechanism. The actual tip-to-surface distance is easily calculated since imposed displacement and cantilever deformation are known. It is thus possible to obtain an image with nanometer resolution. However, the AFM was recently applied to the study of intermolecular forces by derivatizing the tip and the surface with complementary ligand and receptor sites, and varying only the distance between these structures. It was thus possible to derive force/distance curves. The rupture of specific bonds resulted in sharp jumps of the tip (Figure 5), whose amplitude might be translated in a force that was defined as the bond mechanical strength.

In contrast with the surface forces apparatus, the AFM does not allow an absolute determination of the distance between surfaces. However, there are several advantages with this technique : first, there is no need for perfectly smooth surfaces, and it might well be possible to

study bond formation on the surface of actual cells (Hinterdorfer et al., 1996). Second, since the interaction area is much lower than with the SFA, the relative importance of van der Waals forces is much less, and no complicated calculation is required to subtract the effect of these interactions. Third, hydrodynamic interactions are also much less important, making it possible to impose fairly rapid relative displacement of the tip and the sample. As a consequence, this technique might in principle yield kinetic information on individual bonds. Indeed, in a typical experiment, the AFM tip velocity may be of order of 100 nm/s, which is fiftyfold higher than was achieved with the SFA (Leckband et al., 1995).

Further, while the theoretical force and displacement sensitivity of the AFM might be of order of 0.01 pN and 0.01 nm respectively (Lee et al., 1994), several authors measured adhesion forces with a sensitivity of a few piconewtons. Indeed, Hoh et al. (1992) may have detected the rupture of individual hydrogen bonds in aqueous medium. Recently, much interesting information was reported on ligand-receptor interactions :

The streptavidin-biotin bond was studied by several authors (Florin et al., 1994 ; Lee et al., 1994, Moy et al., 1994). When interacting surfaces were coated with limited densities of avidin and biotin groups, the rupture of specific bonds appeared as sharp jumps of the tip. Two arguments were used to support the view that individual bonds were indeed detected. First, it was argued that when a limited proportion of approaches between the tip and sample resulted in binding, the bond number must be low (Lee et al., 1994). Second, when histograms of the distribution of rupture forces were drawn, there appeared a series of quantized peaks that were multiples of a single value (Florin et al., 1994). The reported value of the rupture strength of the avidin-biotin bond varied between 160 pN (Florin et al., 1994), 257 pN (Moy et al., 1994) and 300–400 pN (Lee et al., 1994). The force required to separate paired adenine and thymine groups was estimated at 54 pN (Boland and Ratner, 1995). Other authors estimated at 400 pN the rupture force of association between two cell adhesion proteoglycans (Dammer et al., 1995). The rupture force of interaction between biotin and polyclonal anti-biotin antibodies was estimated at 111.5 pN (Dammer et al., 1996), and the rupture force of the interaction between human albumin and specific antibodies was about 240 pN (Hinterdorfer et al., 1996). Interestingly, the latter authors reported that this force of 240 pN reduced the bond lifetime by a factor of  $8 \times 10^5$  as compared to the value measured on soluble molecules. Here are some points of practical or theoretical interest :

First, it was noted that the measured bond strength was dependent on the cantilever stiffness (Lee et al., 1994). This point was indeed consistent with a previous report by Evans et al., (1991) who emphasized that bond rupture was a stochastic event whose frequency was dependent on applied force. Further, Evans et al. (1994) suggested an empirical relationship between the level of force "f" for most frequent rupture depends and the rate of loading. It is therefore not surprising that when a very low loading rate was used to probe the streptavidin-biotin interaction with so-called biomembrane-force probe (BFP : Evans and Ritchie, 1997), the detachment force was about 50 pN (Merkel et al., 1995).

Second, it was reported that, in contrast with studies made with the SFA on lipid-incorporated adhesion molecules, adhesion receptors were not damaged with AFM studies, which allowed to perform hundreds or thousands of attachment/detachment cycles without any need for a lateral displacement of the tip (Florin et al., 1994 ; Hinterdorfer et al., 1996).

Third, the importance of using adhesion molecules with sufficient flexibility (Hinterdorfer et al., 1996) or a soft substratum (Florin et al., 1994) was emphasized, thus supporting the concept that geometrical constraints may play an essential role in the efficiency of interaction between surface-bound molecules.

Fourth, the significance of measured forces was studied by comparing the binding strength of interaction between avidin and a series of biotin derivatives : a linear relationship was found between the force and the dissociation enthalpy, not the free energy (Moy et al., 1994 ; Chilkoti et al., 1995). This yielded an estimate of 0.95 nm for the rupture length (Moy et al., 1994). Also, these results suggested that detachment acted as an adiabatic process (i.e. this process did not involve any energy exchange between the ligand-receptor couple and the solvent). This point was by no means obvious, and it had to be clarified in order to allow significant comparison between experimental data and results of computer simulation (Grubmüller et al., 1996).

Fifth, whereas initial studies were focussed on bond rupture, Hinterdorfer et al. (1996) recently attempted to estimate the rate of bond formation between an AFM tip coated with a few antibody molecules and antigen-coated surfaces. Thus, they obtained an estimate of  $5 \times 10^4 \text{ M}^{-1} \text{ s}^{-1}$  for the association rate, which fell within the range of antigen-antibody association constants.

### **Use of lipid vesicles as transducers**

The use of lipid vesicles as transducers was pioneered by Evans et al. (1991). They used micropipettes to approach red blood cells coated with limiting densities of various adhesion molecules. One of these cells was stiffened by chemical cross-linking or aspiration with high

pressure. The other one was maintained with moderate pressure. After a contact of several tens of seconds, a pipette was pulled, resulting in the application of an increasing disruptive force. This force could be calculated by analysis of the cell deformation (Figure 6). The rupture time, and the corresponding force were then recorded. An important finding was that the force required for rapid (i. e. a few second) bond rupture was similar when adhesion was mediated by different antigen-antibody couples or lectin-sugar interactions. This rupture force was close to 10–20 pN.

The authors later improved this method by biochemically glueing a microscopic bead to the surface of a lipid vesicle : accurate determination of the bead position could thus be achieved with interference reflection microscopy (Figure 6). Further, the bead was approached to a test surface that could be moved with a piezoelectric device.

The interest of this technology is i) to take advantage of a tunable transducer with a wide range of forces and ii) to allow some visual control of surface position.

### **Use of hydrodynamic flow to study bond formation and dissociation**

The first experimental check of Bell's theory was provided by Tha et al. (1986). They made use of a device called the "traveling microtube" that had been developed by H. Goldsmith for some years. the basic principle was to drive doublets of antibody-agglutinated erythrocytes into a glass capillary tube with very low pressure. A microscope was used to monitor the motion of individual doublets. The trick was to settle the tube on a moving stage with a velocity opposite to the flow. The observed doublet was thus immobile with respect to the microscope and could be monitored during a prolonged period of time. Also, erythrocytes were made spherical by incubation in hypotonic media, which allowed precise calculation of forces with theoretical results from fluid mechanics. Recording the motion gave direct information on the hydrodynamic distractive force and doublet lifetime. Another important point was to use limiting concentrations of agglutinating antibodies (in order that cells be held together by a few and even a single bond). It was thus possible to observe doublet rupture in presence of weak hydrodynamic forces. The strength of an antigen-antibody bond was estimated at a few tens of piconewtons. These results were essentially confirmed in a later study based on improved methodology allowing rapid variation of the applied force, with a cone-and plate rheoscope (Tees et al., 1993 ; Goldsmith et al., 1994), and replacement of polyclonal with monoclonal antibodies. Finally, since there remained the possibility that cell separation might involve the uprooting of membrane bound antigens rather than rupture of the antigen-antibody bond (Bell, 1978 ; Evans et al., 1991), experiments were resumed in the same laboratory with ligand-derivatized latex spheres. The motion of flowing doublets was followed with a range of shear rates. Computer simulation was performed to fit experimental values of bond lifetime with predictions from Bell's theory. The natural lifetime (i.e. at zero force) of the interaction between a polysaccharide antigen and antibody bonds was found to be 25 seconds, with an empirical interaction range  $\gamma$  (equation 6) of 0.12 nm (Tees and Goldsmith, 1996). The interaction between immunoglobulin G and protein G, a natural IgG receptor of bacterial origin, was also studied : the natural lifetime of the interaction was 175 seconds and the interaction range  $\gamma$  was 0.39 nm (Kwong et al., 1996).

A major limitation of the above experiments is that it was difficult to obtain accurate information on short-lived bonds. This difficulty could be overcome with a parallel-plate flow chamber : the basic idea was to observe the motion of receptor-coated cells or particles along ligand-coated surfaces with a very low hydrodynamic force (say a few piconewtons, i.e. less than the rupture strength of a single bond). It was thus reasoned that single molecular bonds should result in the formation of detectable arrests. When blood polymorphonuclear neutrophils were driven along activated endothelial cells (Kaplanski et al., 1993), they displayed transient arrests with an average lifetime of about 2 seconds, and these binding events were inhibited when endothelial cells were treated with anti-E-selectin antibodies. More extensive data were obtained on a similar system by Alon et al. (1995) who studied the binding of neutrophils with surfaces coated with various densities of P-selectin, with different values of the flow rate : they estimated at 1 second the natural lifetime of P-selectin-ligand bond, and the bond interaction distance (i.e. Bell's parameter  $\gamma$ ) was 0.05 nm.

Since an accurate study of cell motion was made difficult by velocity variations due to cell surface asperities, Pierres et al. (1994 & 1995) used the same technique to study the displacement of spherical particles coated with anti-rabbit immunoglobulins along surfaces derivatized with rabbit Ig. Unexpectedly, they observed short-term arrests (with a lifetime ranging between about one and two seconds), whereas the attachment of soluble antibodies lasted several hours. A quantitative analysis of arrest duration strongly suggested the existence of a biphasic binding process, with formation of an intermediate adhesion state. When the shear rate was increased, the lifetime of the intermediate state decreased with an empirical interaction range  $\gamma$  of about 0.08 nm (unpublished data). It must be emphasized that this finding was in line with previously reported data (Beeson and McConnell, 1994). Note that it is not surprising that these intermediate states were revealed with the flow chamber technology rather than the AFM : indeed, when a sphere moves within binding distance of a surface with a velocity of 10  $\mu\text{m/s}$ , the relative velocity between the receptors and ligands is of order of 5  $\mu\text{m/s}$  (Goldman et al., 1967), as compared to a fiftyfold lower value of the AFM tip velocity.



In a later study, Pierres et al. (1996b) improved the performance of the flow chamber methodology by using image analysis to estimate bead position with 0.1  $\mu\text{m}$  accuracy and a time resolution of 5 milliseconds. It was thus possible to study short term interactions between recombinant CD2 molecules and their ligand CD48. These molecules are expected to play a role in the initial interaction between T lymphocytes and antigen-presenting cells. The natural bond lifetime was of the order of one tenth of a second with a bond range parameter  $\gamma$  of 0.13 nm (Pierres et al., 1996c).

Recently, the flow chamber was used to study the on-rate of bond formation (Pierres et al., 1997) : CD48-coated spheres were driven along CD2-derivatized surfaces. About 2,500 individual trajectories were recorded, yielding 350,000 positions. Results were used to derive a relationship between instantaneous sphere velocity and binding probability. Numerical data provided by Goldman et al. (1967) were then used to plot the binding probability versus the sphere-to-surface distance. The binding site density was then determined, which allowed an absolute estimate of the variations of the rate of bond formation between CD2 and CD48 molecules versus the distance between the anchoring points of these molecules. The association rate was  $0.03 \text{ second}^{-1}$  at 10 nm separation, and it was inversely proportional to the cube of the distance. This finding is not inconsistent with the crude model described in Appendix II.

In conclusion, the flow chamber allowed quantitative determination of both bond rupture rate as a function of applied force, and bond formation kinetics as a function of distance. Also, the occurrence of intermediate binding states was readily demonstrated.

### Other methods

Although they were less extensively used than aforementioned approaches, some techniques were demonstrated to yield valuable information on the association between bound adhesion molecules.

A rapidly developing methodology consists of exerting a force of several piconewtons on a microscopic object by focussing a laser beam on this object on the stage of a microscope (these devices are called laser traps or "optical tweezers"). The force is generated by the deflection of photons when light rays are diffracted (see e. g. Ashkin, 1992). This technique was recently used by Miyata et al. (1996) to study the strength and lifetime of actin- $\alpha$ -actinin interaction. The authors reported very heterogeneous bond lifetime, with a mean duration of 0.52 second. The force dependence was complex, with wide variations of the dissociation constant, which made very difficult a quantitative check of Bell's model. The binding range  $\gamma$  was estimated between 0.05 nm and 0.3 nm.

A recently described method consisted of observing the thermal motion of immunoglobulin-coated spheres near protein A-coated surfaces (protein A is a natural immunoglobulin receptor of microbial origin). Total internal reflection microscopy allowed to monitor elevations as small as 1 nm. The authors took advantage of Boltzmann's distribution to derive an energy/distance relationship between beads and the surface (Liebert and Prieve, 1995).

## CONCLUSION

Experimental studies made on cell adhesion have long suggested that a single parameter such as the affinity constant could not give a complete account of the behavior of cell surface receptors. Recent methodological advances recently yielded an impressive amount of information on ligand receptor interactions, with fairly direct determination of the relationship between i) dissociation rate and applied force, and ii) association rate and distance of interacting molecules. It may thus be expected that accurate comparison between experimental data and theoretical predictions, including results from computer simulation experiments, will dramatically improve our understanding of the mechanisms of macromolecule interactions within the next few years. This will allow in-depth understanding of the relationship between the structure and function of the multiple adhesion molecules that have recently been characterized.

### References:

- Alon R , Hammer DA , Springer TA 1995; Lifetime of the P-selectin-carbohydrate bond and its response to tensile force in hydrodynamic flow. *Nature*. 374: 539- 542
- Ashkin A 1992; Forces of a single-beam gradient laser trap on a dielectric sphere in the ray optics regime. *Biophys J*. 61: 569- 582
- Beeson C , McConnell HM 1994; Kinetic intermediates in the reactions between peptides and proteins of major histocompatibility complex class II. *Proc Natl Acad Sci USA*. 91: 8842- 8845
- Bell GI 1978; Models for the specific adhesion of cells to cells. *Science*. 200: 618- 627
- Bell GI , Dembo M , Bongrand P 1984; Cell adhesion - Competition between nonspecific repulsion and specific bonding. *Biophys J*. 45: 1051- 1064
- Binnig G , Rohrer H , Gerber C , Weibel E 1982; Surface studies by scanning tunneling microscopy. *Phys Rev Lett*. 49: 57- 61
- Boland T , Ratner BD 1995; Direct measurement of hydrogen bonding in DNA nucleotide bases by atomic force microscopy. *Proc Natl Acad Sci USA*. 92: 5297- 5301
- Bongrand P , Capo C , Depieds R 1982; Physics of Cell Adhesion. *Progress Surface Sci*. 12: 217- 286
- Brune D , Kim S 1994; Hydrodynamic steering effects in protein association. *Proc Natl Acad Sci(USA)*. 91: 2930- 2934
- Chilkoti A , Boland T , Ratner BD , Stayton P 1995; The relationship between ligand-binding thermodynamics and protein-ligand interaction forces measured by atomic force microscopy. *Biophys J*. 69: 2125- 2130
- Claesson PM Editor: Bongrand P , Claesson P , Curtis A 1994; The surface force technique : interactions between surfactant bilayers and between protein layers. *Studying Cell Adhesion*. Springer; Heidelberg 15- 35


- Crowley E , Horwitz AF 1995; Tyrosine phosphorylation and cytoskeletal tension regulate the release of fibroblast adhesion. *J Cell Biol.* 131: 525- 537
- Dammer U , Popescu O , Wagner P , Anselmetti D , Güntherodt HJ , Misevic GN 1995; Binding strength between cell adhesion proteoglycans measured by atomic force microscopy. *Science.* 267: 1173- 1175
- Dammer U , Hegner M , Anselmetti D , Wagner P , Dreier M , Huber W , Güntherodt HJ 1996; Specific antigen/antibody interactions measured by force microscopy. *Biophys J.* 70 : 2437- 2441
- Dembo M , Torney DC , Saxman K , Hammer D 1988; The reaction-limited kinetics of membrane-to-surface adhesion and detachment. *Proc Roy Soc London B.* 234: 55- 83
- Erlandsson R , Olsson L Editor: Bongrand P , Claesson P , Curtis A 1994; The study of biomolecules on surfaces using the scanning force microscope. *Studying Cell Adhesion.* Springer; Heidelberg 51- 63
- Evans E , Ritchie K 1997; Dynamic strength of molecular adhesion bonds. *Biophys J.*
- Evans E , Berk D , Leung A 1991; Detachment of agglutinin-bonded red blood cells. I - Forces to rupture molecular point attachments. *Biophys J.* 59: 838- 848
- Evans E , Merkel R , Ritchie K , Tha S , Zilker A Editor: Bongrand P , Claesson P , Curtis A 1994; Picoforce method to probe submicroscopic actions in biomembrane adhesion. *Studying Cell Adhesion.* Springer; Heidelberg 125- 139
- Eyring H 1935; The activated complex in chemical reactions. *J Chem Phys.* 3: 107- 115
- Eyring H , Walter J , Kimball GE 1944; *Quantum Chemistry.* J. Wiley & Sons, Inc; New York 299- 331
- Florin EL , Moy VT , Gaub HE 1994; Adhesion forces between individual ligand-receptor pairs. *Science.* 264: 415- 417
- Goldman AJ , Cox RG , Brenner H 1967; Slow viscous motion of a sphere parallel to a plane wall.II - Couette flow. *Chem Engr Sci.* 22: 653- 660
- Goldsmith HL , Takamura K , Tha S , Tees D Editor: Bongrand P , Claesson P , Curtis A 1994; Study of cell-cell interactions with the travelling microtube. *Studying Cell adhesion.* Springer; Heidelberg 141- 156
- Grubmüller H , Heymann B , Tavan P 1996; Ligand binding : molecular mechanics calculation of the streptavidinbiotin rupture force. *Science.* 271: 997- 999
- Hammer DA , Lauffenburger DA 1987; A dynamical model for receptormediated adhesion to surfaces. *Biophys J.* 52: 475- 487
- Helm CA , Israelachvili JN , McGuiggan PM 1989; Molecular mechanisms and forces involved in the adhesion and fusion of amphiphilic bilayers. *Science.* 246: 919- 922
- Helm C , Knoll W , Israelachvili JN 1991; Measurement of ligand-receptor interactions. *Proc Natl Acad Sci USA.* 88: 8169- 8173
- Hill TL 1960; *An introduction to Statistical Thermodynamics.* Addison-Wesley; Reading, MA 189- 200
- Hinterdorfer P , Baumgartner W , Gruber HJ , Schilcher K , Schindler H 1996; Detection and localization of individual antibody-antigen recognition events by atomic force microscopy. *Proc natl Acad Sci USA.* 93: 3477- 3481
- Hoh JH , Cleveland JP , Prater CB , Revel JP , Hansma PK 1992; Quantized adhesion detected with the atomic force microscope. *J Am Chem Soc.* 114: 4917- 4918
- Israelachvili JN 1991; *Intermolecular & Surface Forces.* Academic Press; London
- Israelachvili JN , Adams GE 1978; Measurement of forces between two mica surfaces in aqueous electrolyte solutions in the range of 0–100 nm. *J Chem Soc Faraday Trans.* 98: 500 - 514
- Israelachvili JN , Tabor D 1972; The measurement of van der Waals dispersion forces in the range 1.5 to 130 nm. *Proc Roy Soc.* A331: 19- 38
- Israelachvili JN , Leckband DE , Schmitt FJ , Zasadzinski J , Walker S , Chirovolu S Editor: Bongrand P , Claesson P , Curtis A 1994; *Studying Cell Adhesion.* Springer; Heidelberg 37- 49
- Kabat EA 1968; *Structural concepts in immunology and immunochemistry.* Holt, Rinehart and Winston Inc; New York 66- 81
- Kaplanski G , Farnarier C , Tissot O , Pierres A , Benoliel AM , Alessi MC , Kaplanski S , Bongrand P 1993; Granulocyte-endothelium initial adhesion : analysis of transient binding events mediated by E-selectin in a laminar shear flow. *Biophys J.* 64: 1922- 1933
- Lawrence MB , Springer TA 1991; Leukocytes roll on a selectin at physiological flow rates : distinction from and prerequisite for adhesion through integrins. *Cell.* 65: 859- 873
- Leckband DE , Israelachvili JN , Schmitt FJ , Knoll W 1992; Long range attraction and molecular rearrangements in receptor-ligand interactions. *Science.* 255: 1419- 1421
- Leckband DE , Kuhl T , Wang HK , Herron J , Müller W , Ringsdorf H 1995; 4-4-20 anti-fluoresceyl IgG recognition of membrane bound hapten : direct evidence for the role of protein and interfacial structure. *Biochemistry.* 34: 11467- 11478
- Lee GU , Kidwell DA , Colton RJ 1994; Sensing discrete streptavidin-biotin interactions with atomic force microscopy. *Langmuir.* 10: 354- 357
- Liebert RB , Prieve DC 1995; Speciec-specific long range interactions between receptor/ligand pairs. *Biophys J.* 69: 66- 73
- Lollo BA , Chan KWH , Hanson EM , Moy VT , Brian AA 1993; Direct evidence for two affinity states for lymphocyte function-associated antigen 1 on activated T cells. *J Biol Chem.* 268: 21693- 21700
- Luckham PF , Klein J 1985; Interactions between smooth solid surfaces in solutions of adsorbing and nonadsorbing polymers in good solvent conditions. *Macromolecules.* 18: 721- 728
- Marra J , Israelachvili J 1985; Direct measurements of forces between phosphatidylcholine and phosphatidylethanolamine bilayers in aqueous electrolyte solutions. *Biochemistry.* 24 : 4608- 4618
- McGuiggan PM , Israelachvili JN 1990; Adhesion and short-range forces between surfaces. Part II : effects of surface lattice mismatch. *J Mater Res.* 5: 2232- 2243
- Merkel R , Ritchie K , Evans E 1995; Slow loading of biotin-streptavidin bonds yields unexpectedly low detachment forces. *Biophys J.* 68: A404-
- Miyata H , Yasuda R , Kinoshita K Jr 1996; Strength and lifetime of the bond between actin and skeletal muscle  $\alpha$ -actinin studied with an optical trapping technique. *Biochim Biophys Acta.* 1290: 83- 88
- Moy VT , Florin EL , Gaub HE 1994; Intermolecular forces and energies between ligands and receptors. *Science.* 266: 257- 259
- Oliver T , Lee J , Jacobson K 1994; Forces exerted by locomoting cells. *Semin Cell Biol.* 5: 139- 147
- Page MI , Jencks WP 1971; Entropic contributions to rate accelerations in enzymic and intramolecular reactions and the chelate effect. *Proc Natl Acad Sci USA.* 68: 1678- 1683
- Patel KD , Nollert U , McEver RP 1995; P-selectin must extend a sufficient length from the plasma membrane to mediate rolling of neutrophils. *J Cell Biol.* 129: 1155- 1164
- Pierres A , Benoliel AM , Bongrand P Editor: Mow VC , Guilak F , Tran-Son-Tay R , Hochmuth RM 1994; Initial steps of cell-substrate adhesion. *Cell mechanics and cellular engineering.* Springer; New York 145- 159
- Pierres A , Benoliel AM , Bongrand P 1995; Measuring the lifetime of bonds made between surface-linked molecules. *J Biol Chem.* 270: 26586- 26592
- Pierres A , Benoliel AM , Bongrand P 1996a; Measuring bonds between surfaceassociated molecules. *J Immunological Methods.* 196: 105- 120
- Pierres A , Benoliel AM , Bongrand P 1996b; Experimental study of the rate of bond formation between individual receptor-coated spheres and ligand-bearing surfaces. *J Physique III.* 6: 807- 824
- Pierres A , Benoliel AM , Bongrand P 1996c; Determination of the lifetime and force dependence of interactions of single bonds between surface-attached CD2 and CD48 adhesion molecules. *Proc Natl Acad Sci USA.* 93: 15114- 15118
- Pierres A , Benoliel AM , Bongrand P , van der Merwe PA 1997; The dependence of the association-rate of surface-attached adhesion molecules CD2 and CD48 on separation distance. *FEBS Lett.*
- Pincet F , Perez E , Bryant G , Lebeau L , Mioskowski C 1994; Long-range attraction between nucleosides with shortrange specificity : direct measurements. *Phys Rev Lett.* 73: 2780- 2784
- Smoluchowski MV 1917; Versuch einen mathematischen theorie der koagulationskinetik kolloider lösungen. *Z Physik Chem.* 92: 129- 168
- Stossel TP 1993; On the crawling of animal cells. *Science.* 260: 1086- 1094
- Tees DF , Coenen O , Goldsmith HL 1993; Interaction forces between red cells agglutinated by antibody. IV Time and force dependence of break-up. *Biophys J.* 65: 1318- 1334

- Tha SP , Shuster J , Goldsmith HL 1986; Interaction forces between red cells agglutinated by antibody. II - measurement of hydrodynamic force of breakup. Biophys J. 50: 1117-1126
- Ushiyama S , Laue TM , Moore KL , Erickson HP , McEver RP 1993; Structural and functional characterization of monomeric soluble P-selectin and comparison with membrane P-selectin. J Biol Chem. 268: 15229- 15237
- van der Merwe PA , Brown MH , Davis SJ , Barclay AN 1993; Affinity and kinetic analysis of the interaction of the cell adhesion molecules rat CD2 and CD48. EMBO J. 12: 4945-4954
- Zurkhov SN 1965; Kinetic concept of the strength of solids. Int J Fract Mech. 1: 311- 323

**Figure 1**

Association between soluble molecules


The formation of a complex between molecules A and B may be divided into i) a diffusive step (1) leading to the formation of an encounter complex (EC) and ii) a reaction step, involving molecular reorientation (2) and association of complementary sites (3).



**Figure 2**

Dependence of bond formation on molecular shape and flexibility


The association between two surface-bound molecules cannot occur if binding sites are not in contact (A) or if molecular rigidity prevents the acquisition of a suitable orientation of binding sites (B). Therefore, the binding efficiency is expected to be much higher when interacting molecules are sufficiently flexible (C).



**Figure 3**

Expected distance-dependence of the association rate of surface-bound adhesion molecules


The crude model described in appendix II was used to estimate the distance dependence of the association rate between a surface coated with freely rotating rod-like molecules and i) a surface coated with rigidly bound ligands with parallel orientation, located in a thin layer (**thick line**), and ii) a surface coated with freely rotating ligands (**thin line**).



**Figure 4**

Study of molecular interactions with the surface forces apparatus


The measured force  $F$  is proportional to the interaction energy  $W$  per unit area. The thin line represents a typical interaction curve between two charged surfaces. When these surfaces are approached, electrostatic repulsion (1) is overcome by van der Waals attraction (2), followed by short range repulsion (see e. g. Marra and Israelachvili, 1985). Typical ligand receptor interactions (thick line) sometimes displayed long-range "steering" attraction (Leckband et al., 1992 & 1995) (1') followed by a very sharp attraction (2'). Note that the force and distance scales are arbitrary.



**Figure 5**

Study of molecular interactions with atomic force microscopy


**A:** The tip of nanometer width is glued to a soft cantilever (c) whose deflection is continuously monitored by a suitable sensor during approach to the sample S. It is thus possible to obtain force/distance curves (**B**). If the sample and tip are coated with specific ligands and receptors, the repulsion that is observed during approach (steps (1) and (2)) is replaced by attraction (3) when the distance is increased. The bond rupture (4) appears as a single (figure) or multiple (not shown) jump whose amplitude represents the bond strength.



**Figure 6**

Use of soft vesicles as tunable transducers to monitor molecular interactions

**(A)** The soft vesicle (1) is manoeuvred into contact with a rigid sphere (2) with two micropipettes ( $P_1$  and  $P_2$ ). After bond formation, the distance between pipettes is increased under continuous video recording. The force exerted on bond rupture is derived by analyzing the deformation of the soft vesicle (Evans et al., 1991). **(B)** a small rigid sphere S is chemically glued to a soft vesicle. The distance between the sphere and the surface is measured with interference reflection microscopy (Evans et al., 1994).



**Figure 7**

theoretical model for the association between surface-bound adhesion receptors

**(A)** Surface  $S_1$  is coated with rod-like molecules of length  $L$ , freely rotating at their anchoring point  $O$ . Ligand sites are considered as smeared in a region of thickness  $\delta$  on the surface  $S_2$  located at distance  $d$  from  $S_1$ . Two different cases were considered quantitatively (as plotted on Figure 3), with free orientation of binding sites in  $S_2$  (**B**) or parallel orientation of these sites (**C**).

