

1 **OLIVE OIL AND COGNITION: RESULTS FROM THE THREE-CITY STUDY**

2

3 *Dem Ger Cogn Dis accepted September 2009-09-07*

4 Claudine Berr<sup>1,2</sup>, Florence Portet<sup>1,2</sup>, Isabelle Carriere<sup>1</sup>, Tasnime N Akbaraly<sup>1,3</sup>, Catherine  
5 Feart<sup>4</sup>, Véronique Gourlet<sup>5</sup>, Nicole Combe<sup>6</sup>, Pascale Barberger-Gateau<sup>4</sup>, Karen Ritchie<sup>1</sup>

6

7 <sup>1</sup>INSERM U888, Montpellier, F-34093 France ; Université Montpellier1, Montpellier, F-  
8 34000 France.

9 <sup>2</sup> CHU Montpellier, CMRR Languedoc Roussillon, F-34000 Montpellier

10 <sup>3</sup> Department of Epidemiology and Public Health. University College London. 1-19  
11 Torrington Place. WC1E6BT London, UK

12 <sup>4</sup>INSERM, U897, Bordeaux, F-33076 France ; Université Bordeaux2, Bordeaux, F-33076  
13 France

14 <sup>5</sup>INSERM, U708, Paris, F- 75000 France; Université Paris6, Paris, F-75000, France

15 <sup>6</sup>ITERG, Département de Nutrition & Santé, Université Bordeaux1, Talence, F-33405 France.

16

17 *Corresponding Author : Claudine Berr, MD, PhD*

18 INSERM U888, Pathologies du système nerveux: recherche épidémiologique et clinique,  
19 Hôpital La Colombière, 39 Avenue Charles Flahault, BP 34493, 34093 Montpellier, Cedex 5,  
20 France

21 Tel: 33 (0)4 99 614 566, Fax : 33 (0)4 99 614 579

22 Mail: [claudine.berr@inserm.fr](mailto:claudine.berr@inserm.fr)

23

24 Short title: Olive oil and cognition

1 **Abstract**

2 **Background:** Olive oil is a major component of the Mediterranean diet suggested to be  
3 beneficial to counteract Alzheimer's disease.

4 **Aim :** Our objective is to examine the association between olive oil use, cognitive deficit and  
5 decline in a large elderly population

6 **Methods:** We followed 6947 subjects with a brief baseline food frequency questionnaire and  
7 repeated cognitive tests. Olive oil intake was categorized as none (22.7%), moderate (use for  
8 cooking or dressing, 39.9%) and intensive (for both cooking and dressing, 37.4%).  
9 Associations between olive oil and cognitive outcomes were examined taking into account  
10 socio-economic factors, health behaviors, health measures and other dietary intakes.

11 **Results:** Participants with moderate or intensive use of olive oil compared to those who never  
12 used olive oil showed lower odds of cognitive deficit for verbal fluency and visual memory.  
13 For cognitive decline, the association with intensive use was significant for visual memory  
14 (adjusted OR=0.83, IC 95% 0.69 to 0.99) but not for verbal fluency (OR=0.85, 0.70 to 1.03).

15 **Conclusions:** This olive oil-cognition association needs to be confirmed. However, our  
16 findings already shed light on the potential importance of olive oil in the Mediterranean diet  
17 and on its beneficial effects on health.

18

19 **Key words:** olive oil- fatty acids- cohort study- cognition-dementia

1  
2  
3  
4  
5  
6  
7  
8  
9  
10  
11  
12  
13  
14  
15  
16  
17  
18  
19  
20  
21  
22  
23  
24

**Introduction**

A recent meta-analyses showed that a greater adherence to the Mediterranean diet was associated with lower overall mortality, mortality from cancer and cardiovascular diseases [1]. In 2006, in a northern Manhattan cohort, findings on overall diet and cognition suggest that high adherence to the Mediterranean diet (MeDi) decreases the risk of cognitive decline and Alzheimer’s disease (AD) in a non demented elderly population [2] and decreases the risk of mortality in AD patients [3]. More recently in the same cohort, higher adherence to the MeDi was associated with a trend for reduced risk of developing mild cognitive impairment (MCI) and with reduced risk of MCI conversion to AD[4]. Furthermore, when investigating the combined association of diet and physical activity[5], it showed that both factors are independently associated with a reduced risk for AD. In a subsample of 1410 subjects included in the E3C Bordeaux cohort [6] with 5 year follow-up after a 24-hour dietary recall, higher adherence to MeDi was not associated with the risk for incident dementia but was associated with a slower MMSE cognitive decline. In an associated editorial, D Knopman [7] suggests that the MeDi may act on cognition through cerebrovascular mechanisms or by affecting the metabolism of the  $\beta$ -amyloid or the tau protein and that effects may be brought about by some components of the MeDi diet.

The so-called Mediterranean diet is characterized by a high intake of vegetables, legumes, fruits and cereals, a moderate to high intake of fish, a high monounsaturated-to-saturated fatty acids ratio and a low intake of dairy products and meat [8]. While several studies have focused on the association between intake of fruits and vegetables or fish and cognitive outcomes other dietary habits which have been previously associated with dementia or cognitive decline (fish or omega 3 oil, fruits and vegetables) [9-20], little is known about

## Olive oil and cognition

1 the association between olive oil, a major component of the Mediterranean diet, and cognitive  
2 functioning.

3  
4 Thus, our objective is to examine this association in a large population based study of  
5 persons aged over 65 years with a four-year follow-up of cognitive functioning.

1 **Methods**

2 *Participants*

3 The 3C study is a multi-center cohort study, conducted in 3 French cities- Bordeaux (South-  
4 West), Montpellier (South-East), and Dijon (North-East) - and designed to estimate the risk of  
5 dementia and cognitive impairment attributable to vascular factors. Persons eligible for  
6 recruitment into the study had to be i) living in these cities or their suburbs and registered on  
7 the electoral rolls, ii) aged 65 years and over, iii) not institutionalized. Between March 1999  
8 and March 2001, 9,294 individuals, randomly drawn from electoral rolls, agreed to participate  
9 in this project. The practical organization of the different components of the data collection  
10 varied slightly between centers and during follow-up. Participants were invited to the study  
11 examination center or in a mobile center in a specially equipped vehicle or examined and  
12 interviewed at home. The detailed description of the study protocol has been published  
13 elsewhere[21]. The study protocol was approved by the Ethics Committee of the University  
14 Hospital of Kremlin-Bicêtre. Each participant signed an informed consent.

15       Following the screening procedure described below, we excluded 217 participants  
16 with prevalent dementia at baseline. The initial study sample was thus composed of 9077  
17 participants. At the second round of the 3C study (2001-2002), 7919 (87.2 %) were examined  
18 again, 217 were deceased, and 941 refused or were lost-to follow-up. At the third round  
19 (2003-2004), 7053 were examined (79.6 % of the survivors after the second round), the  
20 cumulated number of deaths was 551, and 1473 refused or were lost-to follow-up. A total of  
21 8083 (89.1%) participants had at least one follow-up examination over the 4 years. Of these  
22 8083 subjects, 1136 did not have a dietary questionnaire (n=55) or one of the covariates,  
23 leaving 6947 subjects for statistical analysis (of which 86.0% had made 2 visits).

24

1 *Dietary assessment*

2           A brief food frequency questionnaire was administered at baseline to assess the dietary  
3 habits of the participants for nine broad categories of food: 1) meat and poultry, 2) fish  
4 (including seafood), 3) eggs, 4) milk and dairy products, 5) cereals (including bread and  
5 starches), 6) raw fruits, 7) raw vegetables, 8) cooked fruit or vegetables, 9) pulses. Frequency  
6 of consumption was recorded in six classes: never, less than once a week, once a week, 2-3  
7 times a week, 4-6 times a week, daily. Dietary habits of the sample have been described  
8 elsewhere [22]. Participants were also invited to indicate the dietary fats that they used at least  
9 once a week for dressing, cooking, or spreading among the following list: butter, margarine,  
10 corn oil, peanut oil, sunflower or grape seed oil, olive oil, mixed oil, duck or goose fat, lard,  
11 “Vegetaline”<sup>©</sup> shortening (mainly saturated fat), colza oil, walnut oil, soya oil. Three  
12 categories of olive oil consumption were then defined: “no use”, “moderate use”: using olive  
13 oil for cooking or dressing alone and “intensive” use: using olive oil both for cooking and  
14 dressing.

15

16 *Neuropsychological evaluation*

17           The cognitive tests were administered by trained psychologists at baseline and then  
18 regularly repeated during the 4-years of follow-up. The 30-item Mini-Mental-State-  
19 Examination (MMSE) [23] was used as an index of global cognitive performance. The  
20 recognition form of the Benton Visual Retention Test (BVRT) which evaluates immediate  
21 visual memory, consists in the presentation for 10 seconds of a stimulus card displaying a  
22 geometric figure after which individuals are asked to identify the initial figure among 4  
23 possibilities[24]. Fifteen figures are successively presented and scores range from 0 to 15.  
24 The Isaacs Set Test (IST),[25], consisting of generating words belonging to semantic

1 categories in 30 seconds, measured mostly semantic verbal fluency abilities but also speed of  
2 verbal production .

3  
4 Deficit in global cognitive functioning, visual memory and verbal fluency were  
5 defined respectively by a score below the first quartile of its distribution: 26 for the MMSE,  
6 10 for the BVRT and 40 for the IST. Cognitive decline was defined as being in the lowest  
7 15% of the difference between baseline score and either follow-up visit for the three tests  
8 assessed at each wave, corresponding to a loss of 3 points on the MMSE score, 9 points on  
9 IST and 3 points on BVRT. Diagnosis of prevalent cases of dementia at baseline and incident  
10 cases during the follow-up was based on a three-step procedure with an independent  
11 committee of neurologists reviewing all potential cases of dementia and is fully detailed  
12 elsewhere [26] [27].

13  
14 *Covariates*

15 Socio-demographic informations recorded at baseline include age, gender, marital  
16 status, educational level (no or primary/secondary/high school/university) and income level  
17 per month (<750 €, 750-1500€, 1500-2250 €, >2250 €). Health behaviours were assessed by  
18 smoking status (non/former/current) and diet. Dietary habits include intake of fish/seafood (<  
19 once a week, once a week,  $\geq$  twice a week), fruits and vegetables (<2 portions/day, 2  
20 portions/day,  $\geq$  3 portions/day), intake of wine or other alcoholic beverages per week (0, 1-12,  
21 13-36, >36 g/day), coffee consumption ( $\leq$ 2 cups/day, 2-3 cups/day, >3 cups/day) and regular  
22 consumption (yes/no) of at least one oil rich in omega3 polyunsaturated fatty acids (PUFA)  
23 (colza, walnut or soya oils). Health factors include history of cardiovascular diseases (yes/no),  
24 hypertension (defined as having systolic blood pressure  $\geq$  160 mmHg or diastolic blood  
25 pressure  $\geq$  85 mmHg or use of antihypertensive drugs) [28], diabetes (fasting glycaemia  $\geq$  7.0  
26 mmol/L or having an anti-diabetic treatment), hypercholesterolemia (total plasma cholesterol

1  $\geq 6.20$  mmol/L or use of lipid-lowering drugs), and body mass index (BMI) categorised in  
2 three classes (normal ( $< 25$  kg/m<sup>2</sup>), overweight (25-30 kg/m<sup>2</sup>) and obese ( $\geq 30$  kg/m<sup>2</sup>)).  
3 Depressive symptoms were assessed using the 16 cut-off of the Center for Epidemiological  
4 Studies-Depression scale (CES-D) [29]. Apolipoprotein E (APOE) genotype (at least 1 vs. no  
5  $\epsilon 4$  allele) was also available and was carried out at the Lille Genopole.

6

### 7 *Statistical analyses*

8         Difference in baseline characteristics according to olive oil consumption categories  
9 were tested using Chi<sup>2</sup> tests. Logistic regression analyses were performed to estimate 1) the  
10 cross sectional association between olive oil consumption and cognitive deficit at baseline 2)  
11 the longitudinal association between olive oil consumption and cognitive decline during the  
12 follow-up. In these analyses, the adjustments for covariates were carried out in three steps. A  
13 basic model, model 0, adjusted for sex, age, study center, educational level, income and  
14 baseline cognitive score. Health behaviour and health status variables were added in the  
15 model 1. Smoking status and dietary habits were further added in the model 2. Interactions  
16 between olive oil consumption and covariates included in model 0 were tested and found to  
17 be non significant.

18         Finally, a sensitivity analysis was carried out., the association between olive oil intake  
19 and cognitive decline was re-examined after excluding the 4-year incident dementia cases, as  
20 dietary changes or inaccurate reporting among demented participants are common. Results of  
21 logistic regression were expressed by odds ratios (OR) with their 95% confidence intervals  
22 (CI). Analyses were carried out using SAS software (version 9.1).



1 Results

2 Participants with at least one follow-up, with no diagnosis of dementia at baseline and  
3 for whom complete data on all covariate at baseline were available were included in this  
4 analysis (n=6947). Compared with those excluded (n=2130), participants included in the  
5 analyses were more likely to be younger, with higher educational level and higher income.  
6 Controlling for these factors, a lower proportion of non users of olive oil and of subjects with  
7 poor baseline cognitive functioning was observed in participants included compared to those  
8 excluded (appendix).

9 Baseline characteristics of the 6947 participants as a function of olive oil consumption  
10 are described in Table 1. Among them, 1576 subjects declared no olive oil use (22.7% of total  
11 sample), 2772 declared moderate use (39.9%) only for cooking (5.8%) or dressing (34.1%) ,  
12 and 2599 both for cooking and dressing (intensive use, 37.4 %).

13 INSERT TABLE 1

14 Many factors are associated with olive oil use. The particular place of education is  
15 illustrated in figure 1, more educated participants are more likely to consume olive oil  
16 compared to less educated ones.

17 INSERT FIGURE 1

18 Cross sectional association between olive oil consumption and cognitive deficit at  
19 baseline is shown in Figure 2 which presents odds ratio obtained with logistic regression  
20 (model 2). Compared to those who never used olive oil, participants with high or moderate  
21 consumption of olive oil show significantly lower odds of cognitive deficit in visual memory  
22 and verbal fluency, while no significant association is showed for global cognitive  
23 functioning.

24 INSERT FIGURE 2



1 **Discussion**

2           The present report shows in a large non-demented elderly population that olive oil  
3 consumption habits are significantly associated with selective cognitive deficit and cognitive  
4 decline, independently of other dietary intakes and after adjusting for potential confounders.  
5 Intensive use of olive oil in diet is associated with lower odds of cognitive deficit in visual  
6 memory and verbal fluency and decline in visual memory, however the relationship is not  
7 significant when assessing global cognitive functioning with the MMSE.

8           Olive oil is a major component of the traditional Mediterranean diet. It has been  
9 suggested to have beneficial effect on CHD risk factors [30]. As cardiovascular risk factors  
10 have increasingly been shown to be important for dementia [31] and a greater adherence to  
11 the Mediterranean diet has been shown to reduce the incidence of Alzheimer's disease[2], it is  
12 of particular interest to investigate the association between olive oil consumption and  
13 cognitive decline in an elderly population. To our knowledge, this study is the first to  
14 investigate this relationship with olive oil in a follow-up study, thus making comparisons  
15 difficult.

16           Olive oil contains 70-80 % of monounsaturated fatty acids (MUFA) (oleic acid (18:1  
17 n-9)) and 8–10% of polyunsaturated fatty acids (PUFA) (6-7 % linoleic acid and 1–2% a-  
18 linolenic acid) [32] with great variability depending on production zone, latitude, climate,  
19 variety and stage of maturity of olives when collected [33]. Recent publications on diet in  
20 dementia have specifically focused on PUFA but few examine the association with  
21 monounsaturated fatty acids (MUFA) intakes. A Spanish cross-sectional study [34] showed  
22 that a lower intake of saturated fatty acids (SFA), MUFA and cholesterol was associated with  
23 a better cognitive performance. Cross sectional analyses in an Italian cohort (ILSA) showed  
24 that MUFA intake was significantly associated with better global cognitive functions and  
25 selective attention [35]. More recently, a longitudinal analysis performed in the same cohort

1 [18] showed that high MUFA and PUFA energy intakes were associated with a better  
2 cognitive performance in a 8.5 –year follow-up. In the analyses of 732 participants of the  
3 EPIC Greeks cohort [36], intake of olive oil, MUFA and SFA exhibited a weak but not  
4 significant positive association with cognitive function as assessed by the MMSE, 6 to 13  
5 years after enrolment. In this cohort, intake of PUFA was inversely associated with  
6 performance in the MMSE. Furthermore, four other studies have investigated the association  
7 between intake of omega 3 PUFA and cognition or dementia, giving conflicting results [9,15-  
8 17].

9         However, olive oil is more than a MUFA fat. Olive oil is a functional food which,  
10 besides having a high level of MUFA, contains other minor components with biological  
11 properties [37] such as natural antioxidants including phenolic compounds. Most antioxidant  
12 compounds have been hypothesized to have a protective effect against dementia and cognitive  
13 impairment [38]. Selective antioxidants, including E vitamins, and flavonoids, may reduce  
14 neuronal damage and death from oxidative reactions by inhibiting the generation of reactive  
15 oxygen species, lipid peroxidation, apoptosis, protein oxidation, damage to cell membranes  
16 and/or DNA and beta-amyloid toxicity or deposition [39,40]. Numerous epidemiological  
17 studies have supported these hypotheses in relation to antioxidant nutrients [41-46]. However  
18 at this stage, it is difficult to be more precise as to the mechanism by which olive oil exerts its  
19 beneficial effect on cognitive outcomes, and further studies are needed.

20         Our study has the following limitations. First, tests proposed in the E3C study covered  
21 a limited number of cognitive functions but the battery was prepared in order to screen for  
22 dementia cases and these tests have different metrological properties [47]. Due to ceiling  
23 effects, the BVRT is less appropriate to assess cognitive changes in subjects with very high  
24 cognitive level functioning but is rather sensitive to cognitive change in the low cognitive  
25 levels. The multiple-choice forms used in the E3C study allow assessment of visuospatial

1 working memory - a memory system particularly vulnerable to aging [48]. The selective  
2 association between this test and olive oil consumption could be explained by its metrological  
3 properties, different from those of IST, rather than by a direct biological explanation, which is  
4 more speculative. Furthermore, even if non significant in the final model, we also observed an  
5 association between verbal fluency performance and olive oil. Absence of significant  
6 association with MMSE score may also be related to its skewed distribution.

7 Secondly, our dietary questionnaire is relatively short. We did not collect any  
8 information on portion size, thus excluding any quantitative approach or adjustment on  
9 caloric intake. Nevertheless, this FFQ allowed us to characterize olive oil consumers in this  
10 aged population and to control for other dietary habits which have been previously associated  
11 with dementia and cognitive decline (fish or omega 3 oil, fruits and vegetables) [9-20]. Even  
12 in FFQ more detailed, there is a large degree of reporting error in energy intake (generally  
13 under reporting) which is strongly correlated with BMI and low socio economic status. The  
14 best we could do, according to the type of data collected, was to adjust for consumption of  
15 fish, consumption of vegetables , BMI and socio economic status. Our analyses were also  
16 adjusted for other health behaviour factors such as smoking status and alcohol intake,  
17 however the lack of data on physical activities in the whole population constitutes another  
18 limitation of this study. Moreover, we cannot exclude that the beneficial effect of olive oil on  
19 cognitive decline shown in our cohort could actually be explained by the healthy dietary  
20 habits and healthy life habit behaviors associated with the regular consumption of olive oil.  
21 However the analyses adjusted for dietary intake and for socio-economic status showed that  
22 these factors had little impact on the observed associations between olive oil use and  
23 cognitive deficit or decline. Similarly it could also be argued that use of olive oil may reflect  
24 the physical and cognitive ability of the participants to prepare one's own meals. Our adjusted  
25 analyses for numerous health factors and the results obtained in our sensitivity analyses

## Olive oil and cognition

1 carried out after excluding incident cases of dementia make it less probable as similar  
2 associations were observed while the rates of decline were lower.

3 Finally, the design of our study -an observational epidemiological study- does not  
4 permit us to conclude that there is a causal link between use of olive oil and cognitive decline.  
5 However the longitudinal and multicentric design, the large sample size including more than  
6 8000 elderly subjects from the general population, the high quality of cognitive assessment,  
7 the low rate of attrition of our cohort, and the full adjustment for the potential confounders  
8 constitute the major strengths of this report.

9 In conclusion, to the best of our knowledge this is the first study to show that self-  
10 reported use of olive is associated with lower odds of visual memory decline in a large  
11 general elderly population independently of other dietary intakes and after taking into account  
12 potential confounders. Even if further research is needed to confirm our finding, our results  
13 highlight the importance of olive oil in the Mediterranean diet and in its beneficial effects on  
14 health outcome.

## **Acknowledgments**

- **Funding sources**

The Three-City Study is conducted under a partnership agreement between the Institut National de la Santé et de la Recherche Médicale (INSERM), the Institut de Santé Publique et Développement of the Victor Segalen Bordeaux 2 University, and Sanofi-Aventis. The Fondation pour la Recherche Médicale funded the preparation and initiation of the study. The 3C Study is also supported by the Caisse Nationale Maladie des Travailleurs Salariés, Direction Générale de la Santé, Mutuelle Générale de l'Éducation Nationale, Institut de la Longévité, Regional Councils of Aquitaine and Bourgogne, Fondation de France, and Ministry of Research - INSERM Programme "Cohortes et collections de données biologiques." The Lille Genopole was supported by an unconditional grant from Eisai. This work was carried out with the financial support of the « ANR- Agence Nationale de la Recherche - The French National Research Agency » under the « Programme National de Recherche en Alimentation et nutrition humaine », project COGINUT « ANR-06-PNRA-005".

## References

- 1 Sofi F, Cesari F, Abbate R, Gensini GF, Casini A: Adherence to mediterranean diet and health status: Meta-analysis. *Bmj* 2008;337:a1344.
- 2 Scarmeas N, Stern Y, Tang MX, Mayeux R, Luchsinger JA: Mediterranean diet and risk for alzheimer's disease. *Ann Neurol* 2006;59:912-921.
- 3 Scarmeas N, Luchsinger JA, Mayeux R, Stern Y: Mediterranean diet and alzheimer disease mortality. *Neurology* 2007;69:1084-1093.
- 4 Scarmeas N, Stern Y, Mayeux R, Manly JJ, Schupf N, Luchsinger JA: Mediterranean diet and mild cognitive impairment. *Arch Neurol* 2009;66:216-225.
- 5 Scarmeas N, Luchsinger JA, Schupf N, Brickman AM, Cosentino S, Tang MX, Stern Y: Physical activity, diet, and risk of alzheimer disease. *Jama* 2009;302:627-637.
- 6 Feart C, Samieri C, Rondeau V, Amieva H, Portet F, Dartigues JF, Scarmeas N, Barberger-Gateau P: Adherence to a mediterranean diet, cognitive decline, and risk of dementia. *Jama* 2009;302:638-648.
- 7 Knopman DS: Mediterranean diet and late-life cognitive impairment: A taste of benefit. *Jama* 2009;302:686-687.
- 8 Trichopoulou A, Kouris-Blazos A, Wahlqvist ML, Gnardellis C, Lagiou P, Polychronopoulos E, Vassilakou T, Lipworth L, Trichopoulos D: Diet and overall survival in elderly people. *Bmj* 1995;311:1457-1460.
- 9 Morris MC, Evans DA, Bienias JL, Tangney CC, Bennett DA, Wilson RS, Aggarwal N, Schneider J: Consumption of fish and n-3 fatty acids and risk of incident alzheimer disease. *Arch Neurol* 2003;60:940-946.
- 10 Kalmijn S, Launer LJ, Ott A, Witteman JC, Hofman A, Breteler MM: Dietary fat intake and the risk of incident dementia in the rotterdam study. *Ann Neurol* 1997;42:776-782.
- 11 Barberger-Gateau P, Letenneur L, Deschamps V, Peres K, Dartigues JF, Renaud S: Fish, meat, and risk of dementia: Cohort study. *Bmj* 2002;325:932-933.
- 12 Huang TL, Zandi PP, Tucker KL, Fitzpatrick AL, Kuller LH, Fried LP, Burke GL, Carlson MC: Benefits of fatty fish on dementia risk are stronger for those without apoe epsilon4. *Neurology* 2005;65:1409-1414.
- 13 Barberger-Gateau P, Raffaitin C, Letenneur L, Berr C, Tzourio C, Dartigues JF, Alperovitch A: Dietary patterns and risk of dementia: The three-city cohort study. *Neurology* 2007;69:1921-1930.
- 14 Barberger-gateau P, Commenges D, Gagnon M, Letenneur L, Sauvel C, Dartigues JF: Instrumental activities of daily living as a screening tool for cognitive impairment and dementia in elderly community dwellers. *J Am Geriatr Soc* 1992;40:1129-1134.
- 15 Kalmijn S, Feskens EJ, Launer LJ, Kromhout D: Polyunsaturated fatty acids, antioxidants, and cognitive function in very old men. *Am J Epidemiol* 1997;145:33-41.
- 16 van Gelder BM, Tijhuis M, Kalmijn S, Kromhout D: Fish consumption, n-3 fatty acids, and subsequent 5-y cognitive decline in elderly men: The Zutphen elderly study. *Am J Clin Nutr* 2007;85:1142-1147.
- 17 Engelhart MJ, Geerlings MI, Ruitenberg A, Van Swieten JC, Hofman A, Witteman JC, Breteler MM: Diet and risk of dementia: Does fat matter?: The rotterdam study. *Neurology* 2002;59:1915-1921.
- 18 Solfrizzi V, Colacicco AM, D'Introno A, Capurso C, Torres F, Rizzo C, Capurso A, Panza F: Dietary intake of unsaturated fatty acids and age-related cognitive decline: A 8.5-year follow-up of the italian longitudinal study on aging. *Neurobiol Aging* 2006;27:1694-1704.
- 19 Dai Q, Borenstein AR, Wu Y, Jackson JC, Larson EB: Fruit and vegetable juices and alzheimer's disease: The kame project. *The American journal of medicine* 2006;119:751-759.



- 20 Morris MC, Evans DA, Tangney CC, Bienias JL, Wilson RS: Associations of vegetable and fruit consumption with age-related cognitive change. *Neurology* 2006;67:1370-1376.
- 21 Vascular factors and risk of dementia: Design of the three-city study and baseline characteristics of the study population. *Neuroepidemiology* 2003;22:316-325.
- 22 Larrieu S, Letenneur L, Berr C, Dartigues JF, Ritchie K, Alperovitch A, Tavernier B, Barberger-Gateau P: Sociodemographic differences in dietary habits in a population-based sample of elderly subjects: The 3c study. *J Nutr Health Aging* 2004;8:497-502.
- 23 Folstein MF, Folstein SE, McHugh PR: "Mini-mental state". A practical method for grading the cognitive state of patients for the clinician. *Journal of psychiatric research* 1975;12:189-198.
- 24 Amieva H, Gaestel Y, Dartigues JF: The multiple-choice formats (forms f and g) of the benton visual retention test as a tool to detect age-related memory changes in population-based studies and clinical settings. *Nature protocols* 2006;1:1936-1938.
- 25 Isaacs B, Kennie AT: The set test as an aid to the detection of dementia in old people. *Br J Psychiatry* 1973;123:467-470.
- 26 Three-City Study Group: Vascular factors and risk of dementia: Design of the three-city study and baseline characteristics of the study population. *Neuroepidemiology* 2003;22:316-325.
- 27 American Psychiatric Association: Diagnostic and statistical manual of mental disorders, 4th ed, ed 4th. Washington, DC, APA Press, 1994.
- 28 Brindel P, Hanon O, Dartigues JF, Ritchie K, Lacombe JM, Ducimetiere P, Alperovitch A, Tzourio C: Prevalence, awareness, treatment, and control of hypertension in the elderly: The three city study. *Journal of hypertension* 2006;24:51-58.
- 29 Radloff LS: The ces-d scale : A self-report depression scale for research in the general population. *Appl Psychol Meas* 1977;1:385-401.
- 30 Psaltopoulou T, Naska A, Orfanos P, Trichopoulos D, Mountokalakis T, Trichopoulou A: Olive oil, the mediterranean diet, and arterial blood pressure: The greek european prospective investigation into cancer and nutrition (epic) study. *Am J Clin Nutr* 2004;80:1012-1018.
- 31 Kivipelto M, Helkala EL, Laakso MP, Hanninen T, Hallikainen M, Alhainen K, Soininen H, Tuomilehto J, Nissinen A: Midlife vascular risk factors and alzheimer's disease in later life: Longitudinal, population based study. *Bmj* 2001;322:1447-1451.
- 32 Solfrizzi V, D'Introno A, Colacicco AM, Capurso C, Del Parigi A, Capurso S, Gadaleta A, Capurso A, Panza F: Dietary fatty acids intake: Possible role in cognitive decline and dementia. *Exp Gerontol* 2005;40:257-270.
- 33 Boskou D: Olive oil; in Simopoulos A, Visioli F (eds): *Mediterranean diets*. Basel, Karger, 2000, vol 87, pp 56-77.
- 34 Ortega RM, Requejo AM, Andres P, LopezSobaler AM, Quintas ME, Redondo MR, Navia B, Rivas T: Dietary intake and cognitive function in a group of elderly people. *AmJClinNutr* 1997;66:803-809.
- 35 Solfrizzi V, Panza F, Torres F, Mastroianni F, DelParigi A, Venezia A, Capurso A: High monounsaturated fatty acids intake protects against age-related cognitive decline. *Neurology* 1999;52:1563-1569.
- 36 Psaltopoulou T, Kyrozis A, Stathopoulos P, Trichopoulos D, Vassilopoulos D, Trichopoulou A: Diet, physical activity and cognitive impairment among elders: The epic-greece cohort (european prospective investigation into cancer and nutrition). *Public Health Nutr* 2008;1-9.
- 37 Covas MI: Olive oil and the cardiovascular system. *Pharmacol Res* 2007;55:175-186.

- 38 Gillette Guyonnet S, Abellan Van Kan G, Andrieu S, Barberger Gateau P, Berr C, Bonnefoy M, Dartigues JF, de Groot L, Ferry M, Galan P, Hercberg S, Jeandel C, Morris MC, Nourhashemi F, Payette H, Poulain JP, Portet F, Roussel AM, Ritz P, Rolland Y, Vellas B: Iana task force on nutrition and cognitive decline with aging. *J Nutr Health Aging* 2007;11:132-152.
- 39 Floyd RA, Hensley K: Oxidative stress in brain aging. Implications for therapeutics of neurodegenerative diseases. *Neurobiol Aging* 2002;23:795-807.
- 40 Akwa Y, Allain H, Bentue-Ferrer D, Berr C, Bordet R, Geerts H, Nieoullon A, Onteniente B, Vercelletto M: Neuroprotection and neurodegenerative diseases: From biology to clinical practice. *Alzheimer Dis Assoc Disord* 2005;19:226-239.
- 41 Engelhart MJ, Geerlings MI, Ruitenberg A, vanSwieten JC, Holman A, Witteman JCM, Breteler MMB: Dietary intake of antioxidants and risk of alzheimer disease. *Jama J Am Med Assn* 2002;287:3223-3229.
- 42 Morris MC, Evans DA, Bienias JL, Tangney CC, Bennett DA, Aggarwal N, Wilson RS, Scherr PA: Dietary intake of antioxidant nutrients and the risk of incident alzheimer disease on a biracial community study. *Jama Journal of the American Medical Association* 2002;287:3230-3237.
- 43 Zandi PP, Anthony JC, Khachaturian AS, Stone SV, Gustafson D, Tschanz JT, Norton MC, Welsh Bohmer KA, Breitner JCS: Reduced risk of alzheimer disease in users of antioxidant vitamin supplements - the cache county study. *Archives of Neurology* 2004;61:82-88.
- 44 Akbaraly NT, Hininger-Favier I, Carrière I, Arnaud J, Gourlet V, Roussel AM, Berr C: Plasma selenium over time and cognitive decline in the elderly: Results from the eva study *Epidemiology* 2007;18:52-58.
- 45 Akbaraly NT, Faure H, Gourlet V, Favier A, Berr C: Plasma carotenoid levels and cognitive performance in an elderly population: Results of the eva study. *J Gerontol A Biol Sci Med Sci* 2007;62:308-316.
- 46 Letenneur L, Proust-Lima C, Le Gouge A, Dartigues J, Barberger-Gateau P: Flavonoid intake and cognitive decline over a 10-year period. *Am J Epidemiol* 2007
- 47 Proust-Lima C, Amieva H, Dartigues JF, Jacqmin-Gadda H: Sensitivity of four psychometric tests to measure cognitive changes in brain aging-population-based studies. *Am J Epidemiol* 2007;165:344-350.
- 48 Amieva H, Jacqmin-Gadda H, Orgogozo JM, Le Carret N, Helmer C, Letenneur L, Barberger-Gateau P, Fabrigoule C, Dartigues JF: The 9 year cognitive decline before dementia of the alzheimer type: A prospective population-based study. *Brain* 2005;128:1093-1101.

Table 1: Baseline characteristics of the 6947 participants as a function of olive oil use.

		Non use	Moderate use (cooking or dressing)	Intensive use (both cooking and dressing)	Chi2 test
	n	%	%	%	<i>P</i> *
<b>Sex</b>					
Men	2755	23.2	39.9	36.9	0.69
Women	4192	22.4	39.9	37.7	
<b>Age</b>					
65-69	1778	17.5	41.7	40.8	<0.0001
70-74	2334	22.0	39.3	38.7	
75-80	1860	25.2	39.3	35.5	
80+	975	29.2	39.1	31.7	
<b>Education</b>					
No or primary	1654	28.9	38.2	32.9	<0.0001
Secondary	2510	24.5	40.2	35.3	
High school	1419	18.5	40.8	40.7	
University	1364	16.1	40.5	43.4	
<b>City</b>					
Bordeaux	1416	33.0	38.1	28.9	<0.0001
Dijon	3716	25.7	36.2	38.1	
Montpellier	1815	8.4	48.9	42.7	
<b>Marital status</b>					
Married	4128	20.4	40.6	39.0	<0.0001
Divorced	515	20.8	37.9	41.3	
Widowed	1774	27.2	39.0	33.8	
Single	495	28.7	38.2	33.1	
Other	30	10.0	50.0	40.0	
<b>Monthly income</b>					
< 750 €	336	39.0	34.8	26.2	<0.0001
750-1500 €	2122	28.6	39.0	32.4	
1500-2250 €	1976	21.9	40.2	37.9	
>2250 €	2513	16.1	41.1	42.8	
<b>Depressive symptoms</b>					
< 16	5369	22.2	40.3	37.5	0.18
≥ 16	1578	24.3	38.6	37.1	
<b>Cardiovascular</b>					
Disease	6336	22.5	40.0	37.5	0.45
No	611	24.7	39.1	36.2	
Yes					
<b>BodyMass Index</b>					
Normal	3276	19.4	41.4	39.2	

Overweight	2758	24.4	38.5	37.1	<0.0001
Obese	913	29.1	38.9	32.0	
Diabetes					
No	6302	22.0	40.3	37.7	
Yes	645	29.0	36.1	34.9	0.0003
Hypertension					
No	3044	19.9	40.7	39.4	
Yes	3903	24.9	39.3	35.8	<0.0001
Hypercholesterolemia					
No	1794	22.1	40.0	37.9	
Yes	5153	22.9	39.9	37.2	0.79
Smoking					
Never	4230	22.7	39.9	37.4	
Former	2339	22.0	39.6	38.5	0.08
Current	378	26.7	42.1	31.2	
APOE ε4 carrier					
No	5582	22.8	40.1	37.1	0.59
Yes	1365	22.3	39.1	38.6	
Alcohol					
0	1375	29.6	39.4	31.0	
1-12 g/day	3141	20.8	39.9	39.3	<0.0001
13-36 g/day	1843	20.0	41.1	38.9	
> 36g/day	588	25.0	37.4	37.6	
Coffee					
≤ 2 cups/day	4784	23.6	39.8	36.5	
2-3 cups/day	1449	21.5	40.9	37.6	0.005
> 3 cups/day	714	18.8	38.4	42.9	
Fish					
Less than once/week	782	33.6	36.7	29.7	
Once /week	2630	28.4	39.8	31.8	<0.0001
Twice or more/week	3535	16.0	40.7	43.3	
Fruits & vegetables					
Less than twice/day	2000	29.6	37.4	33.0	
Twice /day	2549	22.3	39.6	38.1	<0.0001
3 times or more/day	2398	17.3	42.4	40.3	
Omega 3 oil					
No	6586	23.0	39.8	37.2	
Yes	361	17.5	42.1	40.4	0.05
Butter					
No	2633	21.6	40.3	38.1	
Yes	4314	23.3	39.6	37.0	0.24
Goose and duck fat					
No	6727	22.7	39.9	37.4	
Yes	220	20.0	41.4	38.6	0.63

Corn oil						
No	6762	22.7	39.6	37.7		
Yes	185	21.6	50.3	28.1	0.008	
Peanuts oil						
No	5803	22.1	38.5	39.4		
Yes	1144	25.9	46.9	27.2	<0.0001	
Sunflower oil						
No	3665	16.5	35.8	47.7		
Yes	3282	29.5	44.5	26.0	<0.0001	
MMSE baseline						
≥ 26	5951	21.8	40.2	38.0	0.0002	
< 26	970	27.6	38.5	33.9		
IST 30 baseline**						
> 40	5246	20.4	39.9	39.7	<0.0001	
≤ 40	1624	29.9	39.9	30.2		
BVRT baseline**						
> 10	4973	20.2	41.1	38.7	<0.0001	
≤ 10	1893	28.8	37.3	33.9		

\* P value are given for the global chi2 test of heterogeneity. We have examined all comparisons between moderate use vs. no use and intensive use vs. no use.

For variables for which heterogeneity test was significant at  $p < 0.05$ , all 2 by 2 comparisons were significant except when comparing moderate use vs. no use for: coffee consumption ( $p=0.11$ ), corn oil use ( $p=0.13$ ), peanuts oil ( $p=0.63$ ).

\*\* Cut-off corresponding to the first quartile of the sample distribution

MMSE= Mini Mental State Examination (global cognitive functioning)

IST= Isaacs Set Test (verbal fluency)

BVRT= Benton Visual Retention Test (visual memory)

Table 2: Longitudinal association between olive oil consumption and cognitive decline

	Cognitive decline		Model 0		Model 1		Model 2	
	%	<i>p</i>	OR(95% CI)	<i>p</i>	OR(95% CI)	<i>p</i>	OR(95% CI)	<i>p</i>
Global cognitive functioning		0.37						
No use	14.9		1		1		1	
Moderate use	14.1		0.90 (0.74-1.08)	0.24	0.91 (0.76-1.10)	0.32	0.94 (0.78-1.13)	0.50
Intensive Use	13.3		0.90(0.74-1.09)	0.27	0.91 (0.75-1.10)	0.34	0.95 (0.78-1.15)	0.58
Verbal fluency		0.30						
No use	16.6		1		1		1	
Moderate use	15.7		0.93 (0.78-1.12)	0.47	0.95 (0.79-1.14)	0.55	0.96 (0.80-1.16)	0.69
Intensive Use	14.8		0.83 (0.68-0.99)	0.04	0.83 (0.69-1.00)	0.05	0.85 (0.70-1.03)	0.10
Visual memory		0.01						
No use	21.4		1		1		1	
Moderate use	19.1		0.88 (0.75-1.05)	0.15	0.90 (0.76-1.07)	0.25	0.91 (0.77-1.09)	0.30
Intensive Use	17.8		0.80 (0.68-0.96)	0.01	0.82 (0.69-0.97)	0.02	0.83 (0.69-0.99)	0.04

Model 0: adjusted on age, sex, centre, education, income and baseline cognitive score

Model 1: Model0 +, depressive symptoms, APOE4, cardiovascular disease, hypertension, diabetes, hypercholesterolemia, BMI

Model 2: Model 1 + smoking status and dietary intake of fruit/vegetable, omega3 oil, fish, coffee, and alcohol

Figure 1: Distribution of olive oil use according to level of education




Figure 2: Cross sectional association between olive oil use and cognitive deficit at baseline: risk of cognitive deficit in olive oil users expressed by the odds-ratio value with its 95% confidence interval calculated with logistic regression model (controlling for age, sex, centre, education, income, depressive symptoms, APOE4, cardiovascular disease, hypertension, diabetes, hypercholesterolemia, BMI, smoking status and dietary intake of fruit/vegetable, omega3 oil, fish, coffee, and alcohol)

