

AlphaIIbeta3 integrin: new allelic variants in Glanzmann thrombasthenia, effects on ITGA2B and ITGB3 mRNA splicing, expression, and structure-function.

Vincent Jallu, Mathilde Dusseaux, Simon Panzer, Marie-Françoise Torchet, Nathalie Hezard, Jenny Goudemand, Alexandre De Brevern, Cécile Kaplan

► **To cite this version:**

Vincent Jallu, Mathilde Dusseaux, Simon Panzer, Marie-Françoise Torchet, Nathalie Hezard, et al.. AlphaIIbeta3 integrin: new allelic variants in Glanzmann thrombasthenia, effects on ITGA2B and ITGB3 mRNA splicing, expression, and structure-function.: New mutations in Glanzmann patients and carriers. *Human Mutation*, Wiley, 2010, 31 (3), pp.237-46. <10.1002/humu.21179>. <inserm-00448024>

HAL Id: inserm-00448024

<http://www.hal.inserm.fr/inserm-00448024>

Submitted on 3 Oct 2014

HAL is a multi-disciplinary open access archive for the deposit and dissemination of scientific research documents, whether they are published or not. The documents may come from teaching and research institutions in France or abroad, or from public or private research centers.

L'archive ouverte pluridisciplinaire **HAL**, est destinée au dépôt et à la diffusion de documents scientifiques de niveau recherche, publiés ou non, émanant des établissements d'enseignement et de recherche français ou étrangers, des laboratoires publics ou privés.

**α IIb β 3 integrin : New allelic variants in Glanzmann Thrombasthenia,
effects on mRNA splicing, expression and structure-function**

**V. JALLU¹, M. DUSSEAUX¹, S. PANZER², M.F. TORCHET³, N. HEZARD⁴,
J. GOUDEMANT⁵, A.G. de BREVERN⁶, C. KAPLAN¹**

¹ Platelet Immunology Laboratory, INTS, Paris, France

**² Department of Blood Group Serology and Transfusion Medicine, Division of Blood
Group Serology, Medical University, Vienne, Austria**

³ Centre des Hémophiles, Groupe Hospitalier Necker-Enfant malade, Paris, France

⁴ Laboratoire d'Hématologie, CHU Robert Debré, Reims, France

⁵ Hôpital Cardiologique, Centre Hospitalier Régional Universitaire, Lille, France

**⁶ Dynamique des Structures et Interactions des Macromolécules Biologiques, INSERM
UMR-S 665, Université Paris Diderot - Paris 7, INTS, Paris, France**

Short title :

New mutations in Glanzmann patients and carriers

Correspondance

Dr Cécile KAPLAN

Laboratoire d'Immunologie Plaquettaire

INTS

6, rue Alexandre Cabanel

75015 Paris, France

Email : ckaplan@ints.fr

1
2
3 **Phone : 33 1 44 49 30 66 Fax : 33 1 45 67 19 30**
4

5
6 **Abstract**
7

8
9
10 Glanzmann thrombasthenia (GT) is an autosomal recessive inherited bleeding disorder
11 characterized by an impaired platelet aggregation due to defects in integrin α IIB β 3 (ITGA2B,
12 ITGB3), a fibrinogen receptor. Mutations from 24 GT patients and 2 carriers of various
13 origins, Caucasian, North-African and Asian were characterized. Promoter and exon
14 sequences of α IIB and β 3 genes were amplified and directly sequenced. Among 29 identified
15 mutations, 17 new allelic variants resulting from nonsense, missense and deletion / insertion
16 mutations were described. RNA alterations were evaluated by using web-servers. The α IIB
17 p.S926L, p.V903F, and β 3 p.C38Y, p.M118R, p.G221D substitutions prevented complex
18 expression at the surface of Cos-7 cells by altering the α IIB or the β 3 subunit structure. As
19 shown by free energy analyses applied on the resolved structure of α IIB β 3 and structural
20 modeling of the mutant, the p.K253M substitution of β 3 helped to define a key role of the
21 K253 in the interaction of the α IIB β -propeller and the β 3 β -I domains. Finally, the α IIB
22 p.Q595H substitution allowed cell surface expression of the complex but its corresponding
23 c.2800G>T mutation is predicted to alter normal RNA splicing. Characterization of GT
24 mutations is of interest for prenatal diagnosis, genetic counseling and provides useful tools for
25 structure-function relationship studies of α IIB β 3.
26
27
28
29
30
31
32
33
34
35
36
37
38
39
40
41
42
43
44
45
46
47
48
49
50
51
52
53
54
55
56
57
58
59
60

54 **Keywords**

55 α IIB β 3, ITGA2B, ITGB3, Glanzmann, protein structure modeling
56
57
58
59
60

INTRODUCTION

Glanzmann thrombasthenia (GT) is an autosomal recessive bleeding disorder characterized by a prolonged bleeding time and an impaired platelet aggregation in response to physiological agonists such as ADP, thrombin, collagen and epinephrine [Caen, 1972]. GT results from quantitative and / or qualitative defects of the fibrinogen (Fg) platelet receptor, the glycoprotein α IIb β 3 complex (OMIM accession number #273800). Classically GT has been divided into 3 subtypes defined by the α IIb β 3 content of patient's platelets : Type I, no or only trace amounts of α IIb β 3; Type II and variant, respectively with low and near normal platelet surface content of functionally impaired α IIb β 3 [Nurden, 2005]. To date, more than 100 different mutations involved in GT phenotype have been described in the α IIb gene and 60 in the β 3 gene (<http://sinaicentral.mssm.edu/intranet/research/glanzmann/menu>). The nature of GT mutations is highly variable excepting some that are commonly identified in consanguineous populations such as the α IIb c.409-3_418del (IVS3-3del13) and the β 3 c.2031_2041del (2031del11) respectively detected in Arab and Iraqi-Jewish GT populations [Newman et al., 1991], the α IIb c.1544+1G>A (IVS15+1/G>A) mutation in French Gypsies [Schlegel et al., 1995], or the β 3 c.1053_1058del (1053CCCAGG) deletion in North-Africans [Morel-Kopp et al., 1997].

Characterization of GT mutations in patients is critical for prenatal diagnosis and genetic counseling. We report here the characterization of 29 mutations identified in 24 thrombasthenic patients and 2 asymptomatic carriers. Using site directed mutagenesis, cell expression models, *in silico* tools and the resolved structure of the α IIb β 3 complex, we have analyzed the pathogenic role of 17 newly described allelic variants. Structural modeling studies of an allelic variant of GT has revealed the importance of the Lys253 in the interaction of the α IIb β -propeller and the β 3 β -I domains.

MATERIAL AND METHODS

Patients

Blood samples from patients were sent to our laboratory for confirmation of GT diagnosis and genetic analysis. Propositi and their detailed platelet phenotype are listed in Supp Table S1. Propositi 4 and 15 are GT carriers. Propositus 4 who had presented mild bleeding and healing difficulties during tympanoplasty was addressed to the laboratory for GT investigation after flow cytometry analysis revealed a 50 % reduction of α IIB β 3 platelet surface expression. Propositus 15 was unexpectedly found to be a GT type I carrier during neonatal alloimmune thrombocytopenia diagnosis following in utero fetal death. Her platelets were phenotyped HPA-3a negative but genetically HPA-3a/b as determined by PCR-Sequence Specific Primer testing (data not shown).

As shown in Supp Table SI, 19 propositi of 22 of confirmed platelet phenotype_ (86 %) were found to be of type I (< 5 % of α IIB β 3 and < 25 % of Fg normal platelet contents). Three propositi were found to be of type II as their platelets presented low but significant α IIB β 3 content (\geq 10 %) and a relatively high level of intra-platelet Fg (\geq 50 %). None of these patients had been previously studied for molecular gene defects. Patients described herein gave their informed consent for genetic analysis.

Monoclonal antibodies

Monoclonal antibodies (Moab) SZ1 (anti-GPIX), SZ22 (anti- α IIB), SZ21 (anti- β 3), P2 (anti- α IIB β 3) and E3 (anti-Fg) were purchased from Beckmann Coulter-France (Villepinte, France). Moabs AP2 (anti- α IIB β 3) and AP3 (anti- β 3) were obtained from Lucron Bioproducts

1
2
3 (Lille, France). Moab Y2-51 (anti- β 3) was purchased from DAKO (Trappes, France). Moab
4
5 XIIF9 to β 3 was a generous gift from G.Vezon (CRTS, Bordeaux, France). In some flow
6
7 cytometry experiments, Moabs SZ22, SZ21 and P2 were directly conjugated to FITC.
8
9

10 11 12 **Blood samples**

13
14
15 Blood samples from patients were collected by venipuncture using 5mM ethylene-diamine-
16
17 tetra-acetic acid as an anticoagulant. Platelet were separated by differential centrifugation and
18
19 used or stored in 70% ethanol at -80°C until RNA was extracted. In some cases, 500 μl of
20
21 whole blood were mixed with 1.3 ml of RNA Later (Applied Biosystem, Courtaboeuf,
22
23 France) and were kept frozen at -80°C for subsequent RNA extraction.
24
25
26
27
28
29
30
31
32

33 34 35 **Flow cytometry**

36
37 For platelet glycoprotein and Fg content analysis, 50 μl of EDTA anticoagulated whole
38
39 blood was diluted in 500 μl of phosphate saline buffer (PBS) pH 7.2 containing 1 % (wt/vol)
40
41 BSA-fraction V (Euromedex, Souffelweyersheim, France) and 3 mM EDTA (PBE buffer).
42
43 Cell were fixed by adding 1.5 ml of PBS - 3 mM EDTA, pH 7.2, containing 1.33 % (vol/vol)
44
45 of a methanol-stabilized formaldehyde solution (Fisher Scientific, Illkirch, France) for 15 min
46
47 at room temperature (RT). The reaction was stopped by adding 2 ml of PBE before cells were
48
49 pelleted by centrifugation at 1500 g for 10 min. Pelleted cells were washed in 2 ml PBE
50
51 before to be resuspended in 1 ml PBE containing 0.1 % (wt/vol) sodium azide and store at 4°C
52
53 until use. 40 μl of the fixed cells suspension were incubated with 3 $\mu\text{g}/\text{ml}$ Moab in PBE (100
54
55 μl final volume), for 1 h at RT. After two washes in 2 ml PBE, the bound Moabs were
56
57 detected by using a polyclonal goat anti-mouse IgG conjugated to FITC (Jackson
58
59 Immunoresearch Europe, Newmarket, UK) diluted in PBE (1 μl for 10 ml) for 30 min at RT.
60
Following one wash in 2 ml PBE, mean fluorescence intensity of cells was analyzed on an

1
2
3 XL-MCL Flow cytometer (Beckman Coulter, Villepinte, France). To detect intraplatelet Fg,
4
5 each incubation steps were done in presence of 1% (wt/vol) saponine (Fischer Scientific).
6
7
8 Amounts of proteins are expressed as a percentage of the mean amounts obtained with normal
9
10 platelets from 10 healthy donors.
11
12
13
14

15
16 To analyze α IIb β 3 wild type or mutant complexes expressed at the surface of transfected
17
18 Cos-7 cells, cells were collected by Trypsin-EDTA treatment and washed 2 times in PBS-1%
19
20 (wt/vol) BSA. 5×10^5 cells were incubated in 100 μ l PBS containing 3 μ g/ml Moabs (SZ22,
21
22 SZ21 or P2) conjugated to FITC for 1 h at RT. Cells were washed twice in 2 ml PBS-BSA
23
24 before to be analyzed by flow cytometry. Dead cells were discarded from analysis by using
25
26 the 7-AAD viability dye (Beckmann-Coulter). Results are expressed as a percentage of
27
28 positive cells for the concerned antigen.
29
30
31
32

33 34 35 36 37 38 39 40 41 42 43 44 45 46 47 48 49 50 51 52 53 54 55 56 57 58 59 60

DNA/RNA extractions

Genomic DNA was isolated from peripheral blood according to a published procedure [Jallu et al., 2002] or by using the MagNA Pure Compact Instrument and the Nucleic Acid Isolation Kit I (Roche Diagnostics, Meylan, France). In some cases, DNA samples were obtained by using QIAamp DNA Blood Mini kit (Qiagen, Courtaboeuf, France). RNA extractions were done on platelets by using the Rneasy Protect MiniKit (Qiagen). DNAs and RNAs were extracted according to the manufacturers' instructions.

Sequencing analysis

Sequencing was performed as previously described [Quintanar et al., 1998] or by GenoScreen (Lille, France). Primers defined by Kato and coworkers [Kato et al., 1992] and Jin and co-workers [Jin et al., 1993] were respectively used to amplify coding sequences of

1
2
3
4
5
6
7
8
9
10
11
12
13
14
15
16
17
18
19
20
21
22
23
24
25
26
27
28
29
30
31
32
33
34
35
36
37
38
39
40
41
42
43
44
45
46
47
48
49
50
51
52
53
54
55
56
57
58
59
60

α IIb and β 3 genes from genomic DNA. Primers to amplify promoter sequences of α IIb and β 3 have been respectively defined according to the genes' reference sequences NG_008331.1 and NG_008332. α IIb promoter sequences were studied by using the following primer pairs ("a" forward, "b" reverse) :

IIbP1a CGCCAGAATTCTCAGTCACGAAGCTTGGCTCAAGACGGAG

IIbP1b CCAGCCAGATATCAGCTATGTACTACCACCGTGCTAGTCC

IIbP2a CGCCAGAATTCTCAGTCACGGGTAAAGATTCAAGAGACAT

IIbP2b CCAGCCAGATATCAGCTATGTCAGATTCCTCCACAGGAAG

IIbP3a CGCCAGAATTCTCAGTCACGGAAGGGAAGGAGGAGGAGCT

IIbP3b CCAGCCAGATATCAGCTATGCTTCCCTTACGGCTCACCTC

Underlined sequences are non genomic sequence for sequencing PCR products. Promoter and exon 1 sequences of β 3 gene were amplified with the primer pair

IIIaE1a ATGTGGTCTTGCCCTCAACAGGTAG

IIIaE1b CACGCTCTCACCCAGGAAGTTACAG by using the Advantage GC genomic PCR kit (Clontech, Saint-Germain-en-Laye, France) and then sequencing was done with the forward primer 3aseq-P1a CCGGAAAACCAAATAAGGC and the reverse primer 3aseq-P1b GACCAGTCCTCCGCGTTTGC.

All identified mutations were confirmed by PCR-Restriction Fragment Length Polymorphism (PCR-RFLP) analysis on a second PCR product. Restriction enzymes are listed in tables 1 and 2 with the corresponding mutations. The experimental details of these RFLP assays can be obtained upon request. When the mutation did not introduce restriction modification, sequencing was done on a second PCR-product. For nucleotide numbering the A nucleotide of the ATG start was designated +1 (cDNA ITGA2B and ITGB3 GenBank accession numbers NM_000419.3 and NM_000212.2 respectively). For amino acid numbering used in the text, the first amino acids of the mature α IIb and β 3 proteins were

1
2
3 designated +1. Residue numbering according to the Human Genome Variation Society
4 (HGVS) nomenclature that designed as +1 the translation initiating methionine
5 (http://www.hgvs.org/mutnomen) is also shown in tables 1 and 2. Both numberings differ by
6
7
8
9
10 31 and 26 residues that correspond to the signaling peptides of α IIB and β 3 respectively. All
11
12 identified allelic variations have been deposited on the NCBI dbSNP database
13 (http://www.ncbi.nlm.nih.gov/SNP/) and their corresponding accession numbers are indicated
14
15
16
17 in tables 1 and 2.
18
19

20 21 22 **Site directed mutagenesis and in-vitro expression**

23
24 Site directed mutagenesis and *in vitro* expression in Cos-7 cells were done as previously
25 described [Jallu et al., 2002] or by using the QuikChange® II XL Site-Directed Mutagenesis
26 Kit (Agilent Technologies, Massy, France) according to the manufacturer's instructions. In
27
28 some mutagenesis experiments, plasmid pcDNA3.1-Zeo containing the α IIB cDNA was used
29 instead of pcDNA3.1-Neo (Invitrogen, Cergy Pontoise, France).
30
31
32
33
34
35
36
37
38
39
40
41
42

43 44 45 **Minigene study**

46
47 To study a mutation predicted to affect mRNA splicing, an α IIB gene fragment covering
48 exons 17 to 20 and containing the mutation was synthesized and introduced in the plasmid
49 Exontrap (Mobitech, Goettingen, Germany). The wild type form of the sequence was obtained
50 by site directed mutagenesis. Synthesis, cloning and site directed mutagenesis were done by
51 the Epoch Biolabs, Inc. Company (Sugar Land, TX, USA). Following transient transfection in
52 Cos cells, total RNA was extracted and mRNA amplified for the exon 17 to 20 and directly
53 sequenced.
54
55
56
57
58
59
60

Western-blot analysis

Trypsin-EDTA collected Cos-7 cells were washed 2 times in PBS and 1 time in 50 mM Tris-HCL pH 7.6, 150 mM NaCl. Cells were resuspended (1.5×10^7 cells) in 100 μ l of the same Tris buffer containing 1% (wt/vol) Triton-X 100 and 2 mM para-Methyl Sulfonyl Fluoride (PMSF). They were lysed for 30 min at 4°C and the lysate centrifuged at 16.000 g for 30 min at 4°C. The supernatant was diluted with 4 volumes of a 150 mM NaCl solution containing 2.5 % (wt/vol) Sulfate Dodecyl Sodium (SDS), 31.25 mM n-ethylmaleimide and 2.5 mM PMSF. The SDS lysate was heated to 100°C for 5 min before 35 μ l were used for western-blot analysis as previously described [Jallu et al., 1994] with a minor modification : bound antibodies were detected by using anti-mouse IgG conjugated to peroxidase associated with a chemiluminescence detection technique.

Computational analyses

The effect of mutations on intron splicing has been predicted *in silico* by using the following tools : Genscan (<http://genes.mit.edu/GENSCAN.html> [Burge and Karlin, 1998]) and NNSPLICE 0.9 (http://www.fruitfly.org/seq_tools/splice.html [Reese et al., 1997]). Modifications of the splicing factors binding sites pattern resulting from the mutations were evaluated by using the ESEFinder tool [Cartegni et al., 2003; Smith et al., 2006].

To compare wild type and a mutant headpiece of α IIB β 3, modeling has been done using the structure (PDB code 2VDL) published by Springer et co-workers [Springer et al., 2008]. Residues 1-452 of α IIB and 109-352 of β 3 were taken into account. The interface between the protein chains has a correct resolution (2.75 Å) and the residues implicated at the interface have all good normalized B-factor values. A structural model of the mutant has been built. This model was checked using ProCheck software to analyze the quality of the modification [Laskowski et al., 1993] and the side-chain position was analyzed with SCWRL [Canutescu et

1
2
3 al., 2003]. Finally, the FastContact web server (<http://structure.pitt.edu/servers/fastcontact/>)
4
5 was used to estimate the direct electrostatic and desolvation interaction free energy between
6
7 two proteins [Camacho and Zhang, 2005; Champ and Camacho, 2007]. The computational
8
9 estimation of the binding free energy is based on a statistically determined desolvation contact
10
11 potential and Coulomb electrostatics with a distance-dependent dielectric constant [Camacho
12
13 and Zhang, 2005]. This free energy provides a reasonably good estimate of experimental
14
15 binding affinities from complex crystal structures.
16
17
18
19
20
21

22 RESULTS

23
24
25
26
27
28
29
30
31
32
33
34
35
36
37
38
39
40
41
42
43
44
45
46
47
48
49
50
51
52
53
54
55
56
57
58
59
60
61
62
63
64
65
66
67
68
69
70
71
72
73
74
75
76
77
78
79
80
81
82
83
84
85
86
87
88
89
90
91
92
93
94
95
96
97
98
99
100
101
102
103
104
105
106
107
108
109
110
111
112
113
114
115
116
117
118
119
120
121
122
123
124
125
126
127
128
129
130
131
132
133
134
135
136
137
138
139
140
141
142
143
144
145
146
147
148
149
150
151
152
153
154
155
156
157
158
159
160
161
162
163
164
165
166
167
168
169
170
171
172
173
174
175
176
177
178
179
180
181
182
183
184
185
186
187
188
189
190
191
192
193
194
195
196
197
198
199
200
201
202
203
204
205
206
207
208
209
210
211
212
213
214
215
216
217
218
219
220
221
222
223
224
225
226
227
228
229
230
231
232
233
234
235
236
237
238
239
240
241
242
243
244
245
246
247
248
249
250
251
252
253
254
255
256
257
258
259
260
261
262
263
264
265
266
267
268
269
270
271
272
273
274
275
276
277
278
279
280
281
282
283
284
285
286
287
288
289
290
291
292
293
294
295
296
297
298
299
300
301
302
303
304
305
306
307
308
309
310
311
312
313
314
315
316
317
318
319
320
321
322
323
324
325
326
327
328
329
330
331
332
333
334
335
336
337
338
339
340
341
342
343
344
345
346
347
348
349
350
351
352
353
354
355
356
357
358
359
360
361
362
363
364
365
366
367
368
369
370
371
372
373
374
375
376
377
378
379
380
381
382
383
384
385
386
387
388
389
390
391
392
393
394
395
396
397
398
399
400
401
402
403
404
405
406
407
408
409
410
411
412
413
414
415
416
417
418
419
420
421
422
423
424
425
426
427
428
429
430
431
432
433
434
435
436
437
438
439
440
441
442
443
444
445
446
447
448
449
450
451
452
453
454
455
456
457
458
459
460
461
462
463
464
465
466
467
468
469
470
471
472
473
474
475
476
477
478
479
480
481
482
483
484
485
486
487
488
489
490
491
492
493
494
495
496
497
498
499
500
501
502
503
504
505
506
507
508
509
510
511
512
513
514
515
516
517
518
519
520
521
522
523
524
525
526
527
528
529
530
531
532
533
534
535
536
537
538
539
540
541
542
543
544
545
546
547
548
549
550
551
552
553
554
555
556
557
558
559
560
561
562
563
564
565
566
567
568
569
570
571
572
573
574
575
576
577
578
579
580
581
582
583
584
585
586
587
588
589
590
591
592
593
594
595
596
597
598
599
600
601
602
603
604
605
606
607
608
609
610
611
612
613
614
615
616
617
618
619
620
621
622
623
624
625
626
627
628
629
630
631
632
633
634
635
636
637
638
639
640
641
642
643
644
645
646
647
648
649
650
651
652
653
654
655
656
657
658
659
660
661
662
663
664
665
666
667
668
669
670
671
672
673
674
675
676
677
678
679
680
681
682
683
684
685
686
687
688
689
690
691
692
693
694
695
696
697
698
699
700
701
702
703
704
705
706
707
708
709
710
711
712
713
714
715
716
717
718
719
720
721
722
723
724
725
726
727
728
729
730
731
732
733
734
735
736
737
738
739
740
741
742
743
744
745
746
747
748
749
750
751
752
753
754
755
756
757
758
759
760
761
762
763
764
765
766
767
768
769
770
771
772
773
774
775
776
777
778
779
780
781
782
783
784
785
786
787
788
789
790
791
792
793
794
795
796
797
798
799
800
801
802
803
804
805
806
807
808
809
810
811
812
813
814
815
816
817
818
819
820
821
822
823
824
825
826
827
828
829
830
831
832
833
834
835
836
837
838
839
840
841
842
843
844
845
846
847
848
849
850
851
852
853
854
855
856
857
858
859
860
861
862
863
864
865
866
867
868
869
870
871
872
873
874
875
876
877
878
879
880
881
882
883
884
885
886
887
888
889
890
891
892
893
894
895
896
897
898
899
900
901
902
903
904
905
906
907
908
909
910
911
912
913
914
915
916
917
918
919
920
921
922
923
924
925
926
927
928
929
930
931
932
933
934
935
936
937
938
939
940
941
942
943
944
945
946
947
948
949
950
951
952
953
954
955
956
957
958
959
960
961
962
963
964
965
966
967
968
969
970
971
972
973
974
975
976
977
978
979
980
981
982
983
984
985
986
987
988
989
990
991
992
993
994
995
996
997
998
999
1000
1001
1002
1003
1004
1005
1006
1007
1008
1009
1010
1011
1012
1013
1014
1015
1016
1017
1018
1019
1020
1021
1022
1023
1024
1025
1026
1027
1028
1029
1030
1031
1032
1033
1034
1035
1036
1037
1038
1039
1040
1041
1042
1043
1044
1045
1046
1047
1048
1049
1050
1051
1052
1053
1054
1055
1056
1057
1058
1059
1060
1061
1062
1063
1064
1065
1066
1067
1068
1069
1070
1071
1072
1073
1074
1075
1076
1077
1078
1079
1080
1081
1082
1083
1084
1085
1086
1087
1088
1089
1090
1091
1092
1093
1094
1095
1096
1097
1098
1099
1100
1101
1102
1103
1104
1105
1106
1107
1108
1109
1110
1111
1112
1113
1114
1115
1116
1117
1118
1119
1120
1121
1122
1123
1124
1125
1126
1127
1128
1129
1130
1131
1132
1133
1134
1135
1136
1137
1138
1139
1140
1141
1142
1143
1144
1145
1146
1147
1148
1149
1150
1151
1152
1153
1154
1155
1156
1157
1158
1159
1160
1161
1162
1163
1164
1165
1166
1167
1168
1169
1170
1171
1172
1173
1174
1175
1176
1177
1178
1179
1180
1181
1182
1183
1184
1185
1186
1187
1188
1189
1190
1191
1192
1193
1194
1195
1196
1197
1198
1199
1200
1201
1202
1203
1204
1205
1206
1207
1208
1209
1210
1211
1212
1213
1214
1215
1216
1217
1218
1219
1220
1221
1222
1223
1224
1225
1226
1227
1228
1229
1230
1231
1232
1233
1234
1235
1236
1237
1238
1239
1240
1241
1242
1243
1244
1245
1246
1247
1248
1249
1250
1251
1252
1253
1254
1255
1256
1257
1258
1259
1260
1261
1262
1263
1264
1265
1266
1267
1268
1269
1270
1271
1272
1273
1274
1275
1276
1277
1278
1279
1280
1281
1282
1283
1284
1285
1286
1287
1288
1289
1290
1291
1292
1293
1294
1295
1296
1297
1298
1299
1300
1301
1302
1303
1304
1305
1306
1307
1308
1309
1310
1311
1312
1313
1314
1315
1316
1317
1318
1319
1320
1321
1322
1323
1324
1325
1326
1327
1328
1329
1330
1331
1332
1333
1334
1335
1336
1337
1338
1339
1340
1341
1342
1343
1344
1345
1346
1347
1348
1349
1350
1351
1352
1353
1354
1355
1356
1357
1358
1359
1360
1361
1362
1363
1364
1365
1366
1367
1368
1369
1370
1371
1372
1373
1374
1375
1376
1377
1378
1379
1380
1381
1382
1383
1384
1385
1386
1387
1388
1389
1390
1391
1392
1393
1394
1395
1396
1397
1398
1399
1400
1401
1402
1403
1404
1405
1406
1407
1408
1409
1410
1411
1412
1413
1414
1415
1416
1417
1418
1419
1420
1421
1422
1423
1424
1425
1426
1427
1428
1429
1430
1431
1432
1433
1434
1435
1436
1437
1438
1439
1440
1441
1442
1443
1444
1445
1446
1447
1448
1449
1450
1451
1452
1453
1454
1455
1456
1457
1458
1459
1460
1461
1462
1463
1464
1465
1466
1467
1468
1469
1470
1471
1472
1473
1474
1475
1476
1477
1478
1479
1480
1481
1482
1483
1484
1485
1486
1487
1488
1489
1490
1491
1492
1493
1494
1495
1496
1497
1498
1499
1500
1501
1502
1503
1504
1505
1506
1507
1508
1509
1510
1511
1512
1513
1514
1515
1516
1517
1518
1519
1520
1521
1522
1523
1524
1525
1526
1527
1528
1529
1530
1531
1532
1533
1534
1535
1536
1537
1538
1539
1540
1541
1542
1543
1544
1545
1546
1547
1548
1549
1550
1551
1552
1553
1554
1555
1556
1557
1558
1559
1560
1561
1562
1563
1564
1565
1566
1567
1568
1569
1570
1571
1572
1573
1574
1575
1576
1577
1578
1579
1580
1581
1582
1583
1584
1585
1586
1587
1588
1589
1590
1591
1592
1593
1594
1595
1596
1597
1598
1599
1600
1601
1602
1603
1604
1605
1606
1607
1608
1609
1610
1611
1612
1613
1614
1615
1616
1617
1618
1619
1620
1621
1622
1623
1624
1625
1626
1627
1628
1629
1630
1631
1632
1633
1634
1635
1636
1637
1638
1639
1640
1641
1642
1643
1644
1645
1646
1647
1648
1649
1650
1651
1652
1653
1654
1655
1656
1657
1658
1659
1660
1661
1662
1663
1664
1665
1666
1667
1668
1669
1670
1671
1672
1673
1674
1675
1676
1677
1678
1679
1680
1681
1682
1683
1684
1685
1686
1687
1688
1689
1690
1691
1692
1693
1694
1695
1696
1697
1698
1699
1700
1701
1702
1703
1704
1705
1706
1707
1708
1709
1710
1711
1712
1713
1714
1715
1716
1717
1718
1719
1720
1721
1722
1723
1724
1725
1726
1727
1728
1729
1730
1731
1732
1733
1734
1735
1736
1737
1738
1739
1740
1741
1742
1743
1744
1745
1746
1747
1748
1749
1750
1751
1752
1753
1754
1755
1756
1757
1758
1759
1760
1761
1762
1763
1764
1765
1766
1767
1768
1769
1770
1771
1772
1773
1774
1775
1776
1777
1778
1779
1780
1781
1782
1783
1784
1785
1786
1787
1788
1789
1790
1791
1792
1793
1794
1795
1796
1797
1798
1799
1800
1801
1802
1803
1804
1805
1806
1807
1808
1809
1810
1811
1812
1813
1814
1815
1816
1817
1818
1819
1820
1821
1822
1823
1824
1825
1826
1827
1828
1829
1830
1831
1832
1833
1834
1835
1836
1837
1838
1839
1840
1841
1842
1843
1844
1845
1846
1847
1848
1849
1850
1851
1852
1853
1854
1855
1856
1857
1858
1859
1860
1861
1862
1863
1864
1865
1866
1867
1868
1869
1870
1871
1872
1873
1874
1875
1876
1877
1878
1879
1880
1881
1882
1883
1884
1885
1886
1887
1888
1889
1890
1891
1892
1893
1894
1895
1896
1897
1898
1899
1900
1901
1902
1903
1904
1905
1906
1907
1908
1909
1910
1911
1912
1913
1914
1915
1916
1917
1918
1919
1920
1921
1922
1923
1924
1925
1926
1927
1928
1929
1930
1931
1932
1933
1934
1935
1936
1937
1938
1939
1940
1941
1942
1943
1944
1945
1946
1947
1948
1949
1950
1951
1952
1953
1954
1955
1956
1957
1958
1959
1960
1961
1962
1963
1964
1965
1966
1967
1968
1969
1970
1971
1972
1973
1974
1975
1976
1977
1978
1979
1980
1981
1982
1983
1984
1985
1986
1987
1988
1989
1990
1991
1992
1993
1994
1995
1996
1997
1998
1999
2000
2001
2002
2003
2004
2005
2006
2007
2008
2009
2010
2011
2012
2013
2014
2015
2016
2017
2018
2019
2020
2021
2022
2023
2024
2025
2026
2027
2028
2029
2030
2031
2032
2033
2034
2035
2036
2037
2038
2039
2040
2041
2042
2043
2044
2045
2046
2047
2048
2049
2050
2051
2052
2053
2054
2055
2056
2057
2058
2059
2060
2061
2062
2063
2064
2065
2066
2067
2068
2069
2070
2071
2072
2073
2074
2075
2076
2077
2078
2079
2080
2081
2082
2083
2084
2085
2086
2087
2088
2089
2090
2091
2092
2093
2094
2095
2096
2097
2098
2099
2100
2101
2102
2103
2104
2105
2106
2107
2108
2109
2110
2111
2112
2113
2114
2115
2116
2117
2118
2119
2120
2121
2122
2123
2124
2125
2126
2127
2128
2129
2130
2131
2132
2133
2134
2135
2136
2137
2138
2139
2140
2141
2142
2143
2144
2145
2146
2147
2148
2149
2150
2151
2152
2153
2154
2155
2156
2157
2158
2159
2160
2161
2162
2163
2164
2165
2166
2167
2168
2169
2170
2171
2172
2173
2174
2175
2176
2177
2178
2179
2180
2181
2182
2183
2184
2185
2186
2187
2188
2189
2190
2191
2192
2193
2194
2195
2196
219

1
2
3 patients 3 and 5, and the carrier 4) who present the same 13 base pair deletion c.1442-
4 13_1442-1del. However for these latter propositi, an unknown common ancestry cannot be
5
6 excluded as all are European Caucasians who live in the north-east of France. Two
7
8 heterozygous single base insertions were identified in patient 2. These mutations are predicted
9
10 to result in frame-shift anomalies creating a downstream stop codon and premature
11
12 termination. A 10 base pair deletion identified in patient 14 also leads to a frame shift
13
14 anomaly. Other gene anomalies identified affecting α IIb, were missense mutations. Three of
15
16 them (patients 6, 10, 15 and 16) have not been described so far and 6 (patients 3, 11, 12, 13
17
18 and 18) have already been reported (see references in Table 1). Among the 9 heterozygous
19
20 compounds, 4 second obligatory mutations (patients 5, 7, 17 and 18) remain to be identified
21
22 (NI abbreviation (not identified) in tables).
23
24
25
26
27
28
29
30
31
32

Sequence alterations detected in the β 3 gene

33
34 Ten mutations were identified in the gene coding for the β 3 glycoprotein for 8 propositi
35
36 (see Table 2). One single point mutation (patient 20) is responsible for a nonsense codon and
37
38 a second (patient 19) disrupts an acceptor splice site. An homozygous 4 base pair deletion
39
40 located downstream of the donor splice site (c.1125+3_1125+6del) of intron 8 has been
41
42 identified as the unique mutation in patient 25. In patient 26, two silent homozygous
43
44 mutations, c.1143C>A and c.1260G>A (20507C>A and 20624G>A with genomic
45
46 numbering : accession number J05427) located in exon 9 are observed. These mutations have
47
48 been reported to be responsible for type I GT [Jin et al., 1996]. Though silent, their
49
50 association induces exon 9 skipping and the introduction of five intronic bases in the final
51
52 transcript [Jin et al., 1996]. Analysis of the mRNA from patient 26 has confirmed this deletion
53
54 / insertion mechanism (data not shown). Finally five single point missense mutations were
55
56
57
58
59
60

1
2
3 identified (patients 21, 22, 23 and 24). In two heterozygous patients the second obligatory GT
4
5 mutation remains to be determined.
6
7
8
9

10 **Flow cytometry analysis of mutant complexes expressed in COS-7 cells**

11
12 Seven missense mutations not previously reported in the literature have been further
13 studied by expressing the mutant complexes in COS-7 cells to assess their role in the GT
14 phenotype. Cell surface expression of the mutant complexes was assessed by flow cytometry
15 using Moabs SZ22 to α IIB, SZ 21 to β 3 or P2 to the α IIB β 3 complex (see Figure 1). α IIB
16 p.S926L (patient 16) and p.V903F (patients 6 and 15) and β 3 p.C38Y (patient 21), p.M118R
17 (patient 22), p.G221D and p.K253M (patient 24) substitutions prevented normal complex
18 expression, mimicking the type I GT phenotype observed for the patients. For the α IIB
19 p.S926L (patient 16) and β 3 p.G221D and p.K253M (patient 24) mutants the percentage of
20 positive cells expressing β 3 was slightly higher than in other mutants tested (see Figure 1).
21 This low level expression could be correlated to the 10 % of β 3 expression at the surface of
22 the patients' platelets (see Supp Table S1).
23
24
25
26
27
28
29
30
31
32
33
34
35
36
37
38
39
40
41
42
43
44
45
46
47
48
49
50
51
52
53
54
55
56
57
58
59
60

Surprisingly, the α IIB p.Q595H substitution identified in a GT type I patient did not affect
the expression of the mutant complex which is similar to that obtained with the wild type
form.

As control, untransfected cells did not significantly bind any of the Moabs tested and
irrelevant mouse IgG gave less than 1 % of positive cells whatever the transfectant tested.

In silico analyses of mRNA processing for missense allelic variations and for the intronic deletion c.1125+3_1125+6del

Some exonic single point mutations can alter normal processing of the mRNA and lead to
the synthesis of an aberrant mRNA, associated or not with its degradation. Genscan,

1
2
3
4
5
6
7
8
9
10
11
12
13
14
15
16
17
18
19
20
21
22
23
24
25
26
27
28
29
30
31
32
33
34
35
36
37
38
39
40
41
42
43
44
45
46
47
48
49
50
51
52
53
54
55
56
57
58
59
60

NNSPLICE 0.9 and ESEFinder web servers were used to predict the effect of the missense mutations responsible for the substitutions that have been tested in flow cytometry. From this panel, the c.1878G>C responsible for the p.Q595H substitution was the unique variation predicted to modify normal splicing and to affect ESE binding sites. Splice site prediction by NNSPLICE 0.9 cannot identify the normal donor splice site of intron 18 in the mutant. Genscan analysis did not predict splice anomalies, but scores were lowered for both exons 18 (0.987 to 0.791) and 19 (0.995 to 0,849) implying a probability of correct splicing of 75 % instead of 99 % for the normal sequence. Furthermore ESEFinder shows mutation 1878G>C to increase the score of an SF2/ASF binding site from 2.67 to 3.28 and to create a new one with a score of 2.50 (threshold 1.95). Finally an SRp55 binding site that presents a high score (4.33 - threshold 2.67) encompassing the normal splice site is also created.

In the $\beta 3$ gene, the homozygous 4 bp deletion c.1125+3_1125+6del (patient 25) was the single intronic mutation identified which does not directly affect a splice site. However Genscan analysis predicts the use of a cryptic donor splice site located 70 bp downstream the normal one. If this transcript is not degraded, this would lead to an in frame introduction of 23 amino-acids between $\beta 3$ β -I and hybrid domains. ESEFinder shows no modifications but NNSPLICE 0.9 suggests the contiguous donor splice site would be ignored in the mutant.

Western-blot analysis of the p.C38Y $\beta 3$ mutation effect on the complex expression.

All the $\beta 3$ mutants tested strongly affect complex expression (Figure 1) but the p.C38Y substitution also strongly and specifically affects that of $\beta 3$ when tested with anti- $\beta 3$ Moab SZ21. To understand this specific effect, synthesis of the Y38 $\beta 3$ form of the complex in the transfected cells was analyzed by using a Western-blot technique with Moabs SZ22 to α Ib and SZ21, XIIF9 and AP3 to $\beta 3$ (see Figure 2). SZ22 showed that α Ib is clearly detectable in the lysate of the wild type α Ib $\beta 3$ transfectant (lane c) as in the control platelet lysate (lane a).

1
2
3
4
5
6
7
8
9
10
11
12
13
14
15
16
17
18
19
20
21
22
23
24
25
26
27
28
29
30
31
32
33
34
35
36
37
38
39
40
41
42
43
44
45
46
47
48
49
50
51
52
53
54
55
56
57
58
59
60

For the α IIb-Y38 β 3 mutant transfectant, only a band of slightly greater relative mobility which might correspond to a degradation product of α IIb was detected (lane d). Moab XIIF9 revealed the presence of the Y38 β 3 mutant in the α IIb-Y38 β 3 transfectant (lane d). However, this mutant glycoprotein presents a lower relative mobility when compared to the wild type forms detected in lysates from platelets or wild type α IIb β 3 transfectant (lane c). Both SZ21 and AP3 failed to react with the Y38 β 3 mutant (lanes d) but reacted with normal β 3 expressed in platelet and wild type α IIb β 3 transfectant (lanes a and c respectively). XIIF9 and AP3 but not SZ21 reacted with the endogenous β 3 subunit of the vitronectin receptor expressed in Cos-7 cells as shown by the faint band detected in non transfectant cells (lane b). These results suggested that the p.C38Y mutation should impair the complex formation, or at least its normal processing, as no α IIb of normal mobility could be detected.

Role of the Lys253 of the β -I domain of β 3 in the interaction with the α IIb β -propeller

To understand precisely the impact of the p.K253M mutation, we have analyzed the interface between α IIb (residues 1-452) and β 3 (residues 109-352) subunits using the protein structures obtained by the group of Springer (PDB code 2VDL [Springer et al., 2008]). From a simple visualization, the side chain of the β 3 K253 protrudes from the β -I domain toward the α IIb beta-propeller surface and makes contact with different residues of the α IIb subunit (see Figure 3A). Table 3 summarizes the different results concerning the estimations of the direct electrostatic and desolvation interaction free energy between the α IIb and β 3 chains. Interestingly, K253 has a key role in terms of binding free energy with the most negative contribution (-5.5 kcal/mol) and also in terms of electrostatics (-9.7 kcal/mol). Inversely, desolvation free energy is high (+4.1 kcal/mol). The desolvation of charged and polar amino

1
2
3 acid is unfavorable from an energy point of view. This penalty of desolvation is greatly
4 compensated by Coulomb interactions and by the hydrogen bonds which are formed in the
5
6 complex.
7
8

9
10 Analysis of important interactions of by K253 underlines its weight in the interaction as it
11 has the 1st (-6.7 kcal/mol with D232 of α IIb), the 6th (-2.0 kcal/mol with F231) and the 13th (-
12 0.8 kcal/mol with P238) most negative interactions of the interface. One interaction with
13
14 W262 is slightly unfavorable. Concerning free energy, the same observations are made, with
15
16 two strong interactions with D232 (-5.5 kcal/mol) and F231 (-1.2 kcal/mol). One unfavorable
17
18 interaction is observed again with W262. These results highlight the significance of this
19
20 residue, the most important in terms of electrostatic and free energy computation.
21
22

23
24 A structural model of M253 mutant was constructed. Regarding the whole α IIb and β 3
25
26 interface, the substitution provokes a striking loss of electrostatic energy (+9.1 kcal/mol) but a
27
28 gain in desolvation free energy (-5.72 kcal/mol) that cannot counter-balance the decrease of
29
30 electrostatic energy. As shown in Table 3, the diminution of the length of the side-chains and
31
32 the change from a charged polar amino acid to a hydrophobic amino acid has a deep impact.
33
34 Residue 253 has no more (negative) interaction with either D232 or F231, or P238. Only a
35
36 single interaction with W262 is conserved (see Figure 3B).
37
38

39
40 From a visual point of view, the decrease of electrostatic energy between K253 (see
41
42 Figure 3C) and M253 (see Figure 3D) is striking. Overall, the most important electrostatic
43
44 energy implicated in the interaction becomes negligible. All these contacts are lost in presence
45
46 of the methionine, except for W262 that only preserves an electrostatic link. These results
47
48 indicate that it is rather the loss of K253 lost than the presence of the methionine which is
49
50 responsible for the defect of complex expression.
51
52
53
54
55
56
57
58
59
60

DISCUSSION

1
2
3 The aim of the study was to identify mutations responsible for GT platelet phenotype.
4
5 SDS-PAGE, western-blot and flow cytometry analyses carried out on patients' platelets
6
7 confirmed GT clinical diagnosis. Twenty-nine mutations were identified in both α IIb and β 3
8
9 genes. Most of these mutations (47 %) correspond to missense mutations but nonsense,
10
11 insertion, deletion, and single point mutations affecting splice sites were also found.
12
13

14
15 Mutations previously reported in the literature (see tables 1 and 2) have not been further
16
17 studied in-vitro for their involvement in GT. It can be noted that their presence in the patients
18
19 studied herein and in one or more additional unrelated patients would confirm their role in GT
20
21 phenotype. New mutations directly affecting splice sites, creating nonsense codon or insertion
22
23 / deletion associated with frame shift anomaly were considered as known cause of GT. The
24
25 pathological effects of new allelic variations responsible for missense substitutions and one
26
27 new allelic variation leading to a splicing anomaly are discussed below.
28
29

30
31 The substitutions p.V903F (patient 6 and propositus 15) and p.S926L (patient 16) affect
32
33 the extra-cellular N-terminal part of the α IIb β -subunit that composes the calf-2 domain (see
34
35 Figure 4). In both cases they prevent α IIb β 3 complex expression at the Cos-7 cell surface.
36
37 They are predicted to impair the correct folding of the Calf-2 domain as the amino-acids V903
38
39 and S926 are located in its core (see Figure 4). To estimate their structural importance, the
40
41 human α IIb primary sequence has been aligned with several corresponding integrin sequences
42
43 - marsupial, fish, amphibian, prochordates and insect (Supp Figure S1A). These sequences
44
45 share at least 45 % of identity with the human α IIb (Psi-blast data from
46
47 <http://www.ncbi.nlm.nih.gov/BLAST>) and only 17 % of these residues are conserved through
48
49 all the seven organisms tested. It is noteworthy that V903 and S926 belong to these few
50
51 amino-acids, indicating that their conservation could be linked to a strong structural
52
53 evolutionary constraint. Furthermore, their poor conservation in other human integrin α -
54
55 subunits (data not shown) reinforces their α -IIb specificity.
56
57
58
59
60

1
2
3 The missense mutation c.1878G>C (patient 10), contiguous with the donor splice site of
4 intron 18, was predicted to be responsible for a p.Q595H substitution in α IIB. However this
5
6 p.Q595H mutation identified in a type I GT patient does not prevent the expression of the
7
8 mutant complex at the surface of transfected Cos-7 cells. Study of the patient's platelet
9
10 mRNA revealed full normal β 3 mRNA synthesis but α IIB mRNA remained undetectable. A
11
12 splicing anomaly can lead to mRNA decay and explain the lack of detection of the mRNA
13
14 coding for α IIB in patient 10. *In silico* splicing testing by using the Genscan and_NNSPLICE
15
16 0.9 web servers effectively predicted splicing anomalies. Similarly Jayo and coworkers [Jayo
17
18 et al., 2006] reported that the α IIB c.2829C>T mutation (exon 27) predicted to induce a
19
20 p.L912P substitution, partially leads to exon 28 skipping. The authors pointed out that rather
21
22 than the disruption of an ESE site, the aberrant mRNA processing would result from the
23
24 strengthening of the SR protein SC35 binding score by 0.65 units (ESEFinder analysis).
25
26 Interestingly the ESEFinder web server predicted the mutation c.1878G>C to increase the
27
28 score by 0.61 of an existing SF2/ASF binding site and to create two new binding sites for
29
30 SF2/ASF and SRp55 factors with significant high scores (respectively 2.5 and 4.33). It can
31
32 therefore be hypothesized that the 1878G>C mutation alters the normal splice factor pattern
33
34 necessary to accurately splice intron 18. This was confirmed by a minigene study that showed
35
36 the c.1878G>C alteration to provoke exon 18 skipping during mRNA splicing (data not
37
38 shown).
39
40
41
42
43
44
45
46
47
48

49 In the β 3 gene, the homozygous 4 bp deletion c.1125+3_1125+6del (patient 25) was the
50
51 single intronic mutation identified that does not directly affect a splice site but its contiguous
52
53 intronic sequence. Genscan analysis predict an altered splicing that correlates with the partial
54
55 loss of a sequence necessary to hybridize a small nuclear ribonucleoprotein involved in the
56
57 splicing process [Wang and Cooper, 2007].
58
59
60

1
2
3
4
5
6
7
8
9
10
11
12
13
14
15
16
17
18
19
20
21
22
23
24
25
26
27
28
29
30
31
32
33
34
35
36
37
38
39
40
41
42
43
44
45
46
47
48
49
50
51
52
53
54
55
56
57
58
59
60

The mutation c.191G>A (patient 21) responsible for a p.C38Y substitution, is located in the PSI domain of the $\beta 3$. Transient co-transfection of the $\beta 3$ mutated cDNA and with normal α IIb cDNA showed the mutation to reproduce a GT type I phenotype as no complex was detected on the cell surface by any of the anti- α IIb $\beta 3$ Moabs tested herein and including testing with a panel to $\beta 3$ (XIIF9, AP3 and Y2-51, data not shown). Computational analysis (Genscan, ESEFinder and NNSPLICE 0.9) indicated that the c.191G>A mutation would not induce splicing modification of the pre-mRNA. However, western-blot analyses showed that α IIb and the mutated $\beta 3$ were synthesized in transfected COS cells. The mutated Y38 $\beta 3$ presents a lower relative mobility than the wild type form which could result from structural modifications induced by the C16 – C38 disulfide bond disruption. A similar observation has been reported by Valentin and co-workers [Valentin et al., 1995], who have shown that alanine substitution of C38 lowered the relative mobility of $\beta 3$ expressed in COS cells. The structural modification of the PSI domain induced by the p.C38Y mutation was evidenced by the lack of reactivity of Moabs AP3 and SZ21 whose epitopes are respectively located in amino-acids stretches 49-98 [Peterson et al., 2003] and 28-35 [Honda et al., 1995]. So the structural anomaly of the PSI domain impairs the complex association and / or its post-translational maturation leading to its retention inside the cell.

The missense mutations p.M118R, p.G221D and p.K253M are located in the β -I domain of $\beta 3$ and are responsible for Type I GT. M118 and G221 are highly conserved in most integrin β subunits (data not Shown), and through invertebrates to vertebrates (see Supp Figure S1B). Most of the β -I domain mutations previously described, are responsible for Variant [Bajt et al., 1992; D'Andrea et al., 2002; D'Andrea et al., 2008; Lanza et al., 1992; Loftus et al., 1990] or type II GT [Jackson et al., 1998; Morel-Kopp et al., 2002; Nair et al., 2004; Tadokoro et al., 2002; Ward et al., 2000]. For comparison, Figure 5 locates the mutated amino-acids in the primary sequence (panel A) and in the β -I domain structure (panel B). Whatever the variant or

1
2
3 type II GT phenotype, the mutated complexes are located completely or partially at the
4 surface of the β -I domain, and some of them (see Figure 5B, underlined amino-acids) present
5
6 significant electrostatic or / and free energies interactions with the β -propeller (data not
7
8 shown). Variant mutations concern the MIDAS or SyMBS sites [Zhu et al., 2008] directly
9
10 affecting amino-acids involved in metal coordination (p.D119Y, p.S123P) or their very
11
12 nearby environment (p.R217W or Q) (see Figure 5A). Both variant and GT Type II mutations
13
14 support at least a 10 % expression of α IIB β 3 at the platelet surface. This contrasts with the
15
16 failure to detect the β 3-R118 and -D221 forms of the complex. Both the p.M118R and the
17
18 p.G221D mutations, located inside the β -I domain, should deeply alter its proper folding,
19
20 preventing complex expression at the platelet surface. Due to the nature of the amino-acid
21
22 involved, these mutations induce important changes of steric hindrance and charge. However,
23
24 in one case, the mutation p.L117W which affects a 100 % buried amino-acid still permits a 10
25
26 % expression of α IIB β 3 at the platelet surface [Basani et al., 1997]. We could hypothesize that
27
28 the characteristics of these residues, both hydrophobic and aliphatic, could explain this
29
30 behavior. In contrast the p.G221D mutation, changes of volume and charge of the side-chains
31
32 has a too deep impact. Similarly, such a steric hindrance change could explain the type I
33
34 phenotype induced by the p.M124V mutation [Gonzalez-Manchon et al., 2004], as the M124
35
36 side chain pointed inward the β -I domain.
37
38
39
40
41
42
43
44
45
46

47 Unlike for M118 and G221, K253 is not conserved in other integrin β subunits and if it is
48
49 present in β 3 from fish to primates it is replaced by an arginine in amphibian and absent from
50
51 invertebrate integrins (see Supp Figure S1B). However the p.K253M substitution results in
52
53 type I GT. The 3-dimensional structure of the α IIB β 3 headpiece revealed that the side chain of
54
55 K253 protrudes from the β -I domain toward a pocket at the surface of the α IIB β -propeller
56
57 (see Figure 3A and 5B). Analyses of electrostatic, desolvation and free energies revealed that
58
59 K253 is one of the most important amino-acids involved in the β -I domain interaction with
60

1
2
3
4
5
6
7
8
9
10
11
12
13
14
15
16
17
18
19
20
21
22
23
24
25
26
27
28
29
30
31
32
33
34
35
36
37
38
39
40
41
42
43
44
45
46
47
48
49
50
51
52
53
54
55
56
57
58
59
60

the β -propeller of α IIb. Furthermore, a mutant model of the α IIb β 3 headpiece shows that the p.K253M substitution does not introduce steric and energy clashes but only suppresses important contacts with the α IIb β -propeller. So, the p.K253M mutation revealed K253 as a key amino-acid for the association of the β -I domain with the β -propeller. The last mutation reported so far of the β -I domain that leads to type I GT was a substitution of the I325-P326-G327 sequence by a methionine. This mutation impairs the α IIb β 3 association [Morel-Kopp et al., 1997]. However, the mutated amino-acids locate between the α 6 helix and the β 7 sheet and do not directly face the α IIb β -propeller surface (see Figure 5B). Furthermore the I325-P326-G327 sequence does not seem to have significant if any, interaction with the β -propeller as indicated by free and electrostatic energy analysis (data not shown). It might indirectly impair the interaction(s) of the α 5 and / or α 6 helix with the β -propeller. The last four mutations described, p.K137Q [Peterson et al., 2008], p.T140I (HPA-16bw) [Jallu et al., 2002], p.R143Q (HPA-4) [Wang et al., 1992] and p.T195N (HPA-17bw) [Stafford et al., 2008] only lead to allelic variants involved in alloimmune thrombocytopenia that are not related to GT phenotype. All these amino-acids locate at the surface of the β -I domain with their side chains directed outside.

In conclusion, we have identified 29 mutations in a panel of 24 GT patients and 2 GT carriers. Seventeen of these mutations have not been described so far. We have confirmed by *in vitro* studies that the missense mutations identified impaired normal expression of the mutant complex (except for one case). Nonsense and splice site mutations were considered to be related to the GT phenotype. This study emphasizes the high allelic heterogeneity of both the α IIb and β 3 genes involved in GT phenotype. Finally the GT variant p.K253M of β 3 unveiled the key role of K253 in the β -propeller / β -I association of the α IIb β 3 headpiece.

Acknowledgments

We thank the following colleagues for sending us blood samples from the patients we have studied here-in : B. Bastenaire, C. Caron, M. Dreyfus, F. Dutrillaux, I Elalami, Nizard, G. Le Roux, J. Peynet, S. C. Saladin-Thiron / MT Bricquet, P. Tron, J.P. Vernant. We also thank the following technician students for their help in mutation identification : M. Brément, S. Cantais, E. Divron, F. Jacquinet, A. Ledigarcher, P. Moffat, V. Noel and F. Ping. The work done by M. Dusseaux was supported by a grant from Novo Nordisk. Conflict Interests: no conflict financial interests exist.

Reference List

- 1
2
3
4
5
6 Bajt ML, Ginsberg MH, Frelinger III AL, Berndt MC, Loftus JC. 1992. A spontaneous
7 mutation of integrin α IIB β 3 (platelet glycoprotein IIB-IIIa) helps define a ligand
8 binding site. *J Biol Chem* 267:3789-3794.
9
- 10 Basani RB, Brown DL, Vilaire G, Bennett JS, Poncz M. 1997. A Leu¹¹⁷--> Trp mutation
11 within the RGD-peptide cross-linking region of β 3 results in Glanzmann
12 thrombasthenia by preventing α IIB β 3 export to the platelet surface. *Blood* 90:3082-
13 3088.
14
- 15 Basani RB, French DL, Vilaire G, Brown DL, Chen F, Collier BS, Derrick JM, Gartner TK,
16 Bennett JS, Poncz M. 2000. A naturally occurring mutation near the amino terminus
17 of α IIB defines a new region involved in ligand binding to α IIB β 3. *Blood* 95:180-188.
18
- 19 Burge CB, Karlin S. 1998. Finding the genes in genomic DNA. *Curr Opin Struct Biol* 8:346-
20 354.
21
- 22 Caen JP. 1972. Glanzmann thrombasthenia. *J Clin Invest* 1:388-391.
23
- 24 Caen JP. 1989. Glanzmann's thrombasthenia. *Baillieres Clin Haematol* 2:609-625.
25
- 26 Camacho CJ, Zhang C. 2005. FastContact: rapid estimate of contact and binding free
27 energies. *Bioinformatics* 21:2534-2536.
28
- 29 Canutescu AA, Shelenkov AA, Dunbrack RL, Jr. 2003. A graph-theory algorithm for rapid
30 protein side-chain prediction. *Protein Sci* 12:2001-2014.
31
- 32 Cartegni L, Wang J, Zhu Z, Zhang MQ, Krainer AR. 2003. ESEfinder: A web resource to
33 identify exonic splicing enhancers. *Nucleic Acids Res* 31:3568-3571.
34
- 35 Champ PC, Camacho CJ. 2007. FastContact: a free energy scoring tool for protein-protein
36 complex structures. *Nucleic Acids Res* 35:W556-W560.
37
- 38 D'Andrea G, Bafunno V, Del VL, Amoriello A, Morabito P, Vecchione G, Grandone E,
39 Margaglione M. 2008. A beta3 Asp217-->Val substitution in a patient with variant
40 Glanzmann Thrombasthenia severely affects integrin α IIB β 3 functions. *Blood*
41 *Coagul Fibrinolysis* 19:657-662.
42
- 43 D'Andrea G, Colaizzo D, Vecchione G, Grandone E, Di Minno G, Margaglione M, on behalf
44 of the Glanzmann's thrombasthenia Italian team (GLATIT). 2002. Glanzmann's
45 thrombasthenia : Identification of 19 new mutations in 30 patients. *Thromb Haemost*
46 87:1034-1042.
47
- 48 French DL, Collier B. 1997. Hematologically important mutations: Glanzmann
49 thrombasthenia. *Blood Cells, Molecules, and Diseases* 23:39-51.
50
- 51 Gonzalez-Manchon C, Butta N, Larrucea S, Arias-Salgado EG, Alonso S, Lopez A, Parrilla
52 R. 2004. A variant thrombasthenic phenotype associated with compound
53 heterozygosity of integrin β 3-subunit : (Met124Val) β 3 alters the subunit dimerization
54 rendering a decreased number of constitutive active α IIB β 3 receptors. *Thromb*
55 *Haemost* 92:1377-1386.
56
57
58
59
60

- 1
2
3
4
5
6
7
8
9
10
11
12
13
14
15
16
17
18
19
20
21
22
23
24
25
26
27
28
29
30
31
32
33
34
35
36
37
38
39
40
41
42
43
44
45
46
47
48
49
50
51
52
53
54
55
56
57
58
59
60
- Gonzalez-Manchon C, Fernandez-Pinel M, Arias-Salgado EG, Ferrer M, Alvarez MV, Garcia-Munoz S, Ayuso MS, Parrilla R. 1999. Molecular genetic analysis of a compound heterozygote for the glycoprotein (GP) IIb gene associated with Glanzmann's thrombasthenia : disruption of the 674-687 disulfide bridge in GPIIb prevents surface exposure of GPIIb-IIIa complexes. *Blood* 93:866-875.
- Honda S, Honda Y, Bauer B, Ruan C, Kunicki TJ. 1995. The impact of three-dimensional structure on the expression of PI^A alloantigens on human integrin β_3 . *Blood* 86:234-242.
- Jackson DE, White MM, Jennings LK, Newman PJ. 1998. A ser₁₆₂--> Leu mutation within glycoprotein (GP) IIIa (integrin β_3) results in an unstable $\alpha_{IIb}\beta_3$ complex that retains partial function in a novel form of type II Glanzmann thrombasthenia. *Thromb Haemost* 80:42-48.
- Jallu V, Diaz-Ricart M, Ordinas A, Pico M, Vezon G, Nurden AT. 1994. Two human antibodies reacting with different epitopes on integrin β_3 of platelets and endothelial cells. *Eur J Biochem* 222:743-751.
- Jallu V, Meunier M, Brement M, Kaplan C. 2002. A new platelet polymorphism Duv(a+), localized within the RGD binding domain of glycoprotein IIIa, is associated with neonatal thrombocytopenia. *Blood* 99:4449-4456.
- Jayo A, Pabon D, Lastres P, Jimenez-Yuste V, Gonzalez-Manchon C. 2006. Type II Glanzmann thrombasthenia in a compound heterozygote for the alphaIIb gene. A novel missense mutation in exon 27. *Haematologica* 91:1352-1359.
- Jin Y, Dietz HC, Montgomery RR, Bell WR, McIntosh I, Collier B, Bray PF. 1996. Glanzmann thrombasthenia. Cooperative between sequence variants in *Cis* during splice site selection. *J Clin Invest* 98:1745-1754.
- Jin Y, Dietz HC, Nurden A, Bray PF. 1993. Single-strand conformation polymorphism analysis is a rapid and effective method for the identification of mutations and polymorphisms in the gene for GP IIIa. *Blood* 82:2281-2288.
- Kato A, Yamamoto K, Miyazaki S, Jung SM, Maoi M, Aoki N. 1992. Molecular basis for Glanzmann's thrombasthenia (GT) in a compound heterozygote with glycoprotein IIb gene : a proposal for the classification of GT based on the biosynthetic pathway of glycoprotein IIb-IIIa complex. *Blood* 79:3212-3218.
- Lanza F, Stierlé A, Fournier D, Morales M, André G, Nurden AT, Cazenave JP. 1992. A new variant of Glanzmann's thrombasthenia (Strasbourg I). Platelets with functionally defective glycoprotein IIb-IIIa. Complexes and a glycoprotein IIIa²¹⁴ Arg-->²¹⁴ Trp mutation. *J Clin Invest* 89:1995-2004.
- Laskowski RA, Mac Arthur MW, Moss DS, Thornton JM. 1993. PROCHECK : A program to check the stereochemical quality of protein structures. *J Appl Cryst* 26:286-291.
- Loftus JC, O'Toole TE, Plow EF, Glass A, Frelinger III AL, Ginsberg MH. 1990. A β_3 integrin mutation abolishes ligand binding and alters divalent cation-dependent conformation. *Science* 249:915-918.

- 1
2
3 Mitchell WB, Li JH, Singh F, Michelson AD, Bussel J, Collier BS, French DL. 2003. Two
4 novel mutations in the α IIB calcium-binding domains identify hydrophobic regions
5 essential for α IIB β 3 biogenesis. *Blood* 101:2268-2276.
6
7
8 Morel-Kopp MC, Kaplan C, Proulle V, Jallu V, Melchior C, Peyruchaud O, Aurousseau MH,
9 Kieffer N. 1997. A three amino acid deletion in glycoprotein IIIa is responsible for
10 type I Glanzmann's thrombasthenia : Importance of residues Ile325 Pro326 Gly327 for
11 B3 integrin subunit association. *Blood* 90:669-677.
12
13
14 Morel-Kopp MC, Melchior C, Chen P, Ammerlaan W, Lecompte T, Kaplan C, Kieffer N.
15 2002. A naturally occurring point mutation in the β 3 integrin MIDAS-like domain
16 affects differently α v β 3 and α IIB β 3 receptor function. *Thromb Haemost* 86:1425-
17 1434.
18
19
20 Nair S, Ghosh K, Shetty S, Mohanty D. 2004. A novel Ser123Pro substitution in the MIDAS
21 domain of integrin 3 associated with variant Glanzmann's thrombasthenia in an Indian
22 patient. *Haematologica* 89:1529-1530.
23
24
25
26
27
28
29
30
31
32
33
34
35
36
37
38
39
40
41
42
43
44
45
46
47
48
49
50
51
52
53
54
55
56
57
58
59
60
- Negrier C, Vinciguerra C, Attali O, Grenier C, Larcher ME, Dechavanne M. 1998.
Illegitimate transcription : its use for studying genetic abnormalities in lymphoblastoid
cells from patients with Glanzmann thrombasthenia. *Br J Haematol* 100:33-39.
- Newman PJ, Seligsohn U, Lyman S, Collier BS. 1991. The molecular genetic basis of
Glanzmann thrombasthenia in the iraqi-jewish and arab populations in Israel. *Proc
Natl Acad Sci USA* 88:3160-3164.
- Nurden AT. 2005. Qualitative disorders of platelets and megakaryocytes. *J Thromb Haemost*
3:1773-1782.
- Nurden AT, Breillat C, Jacquelin B, Combrie R, Freedman J, Blanchette VS, Schmutz M.
2004. Triple heterozygosity in the integrin α IIB subunit in a patient with Glanzmann's
thrombasthenia. *J Thromb Haemost* 2:813-819.
- Nurden AT, Ruan J, Pasquet JM, Gauthier B, Combrié R, Kunicki T, Nurden P. 2002. A
novel ¹⁹⁶Leu to Pro substitution in the β 3 subunit of the α IIB β 3 integrin in a patient
with a variant form of Glanzmann thrombasthenia. *Platelets* 13:101-111.
- Peretz H, Rosenberg N, Landau M, Usher S, Nelson EJ, Mor-Cohen R, French DL, Mitchell
BW, Nair SC, Chandy M, Collier BS, Srivastava A, Seligsohn U. 2006. Molecular
diversity of Glanzmann thrombasthenia in southern India: new insights into mRNA
splicing and structure-function correlations of alphaIIBbeta3 integrin (ITGA2B,
ITGB3). *Hum Mutat* 27:359-369.
- Peterson JA, Gitter M, Kanack A, Curtis B, McFarland J, Aster RH. 2009. New low-
frequency platelet glycoprotein polymorphisms associated with neonatal alloimmune
thrombocytopenia. *Transfusion* published online : 10 Oct 2009
- Peterson JA, Nyree CE, Newman PJ, Aster RH. 2003. A site involving the "hybrid" and PSI
homology domains of GPIIIa (beta 3-integrin subunit) is a common target for
antibodies associated with quinine-induced immune thrombocytopenia. *Blood*
101:937-942.

- 1
2
3 Quintanar A, Jallu V, Legros Y, Kaplan C. 1998. Human platelet antigen genotyping using a
4 fluorescent SSCP technique with an automatic sequencer. *Br J Haematol* 103:437-444.
5
6
7 Reese MG, Eeckman FH, Kulp D, Haussler D. 1997. Improved splice site detection in Genie.
8 *J Comput Biol* 4:311-323.
9
10 Ruan J, Peyruchaud O, Alberio L, Valles G, Clemetson K, Bourre F, Nurden AT. 1998.
11 Double heterozygosity of the GPIIb gene in a Swiss patient with Glanzmann's
12 thrombasthenia. *Br J Haematol* 102:918-925.
13
14
15 Schlegel N, Gayet O, Morel-Kopp MC, Wyler B, Hurtaud-Roux MF, Kaplan C, Mc Gregor J.
16 1995. The molecular genetic basis of Glanzmann thrombasthenia in a Gipsy
17 population in France. Identification of a new mutation on the α -IIb gene. *Blood*
18 86:977-982.
19
20
21 Scott JP, Chao YL, Newman PJ, Ward CM. 1998. A frameshift mutation at Gly975 in the
22 transmembrane domain of GPIIb prevents GPIIb-IIIa expression. Analysis of two
23 novel mutations in a kindred with type I Glanzmann thrombasthenia. *Thromb*
24 *Haemost* 80:546-550.
25
26
27 Smith PJ, Zhang C, Wang J, Chew SL, Zhang MQ, Krainer AR. 2006. An increased
28 specificity score matrix for the prediction of SF2/ASF-specific exonic splicing
29 enhancers. *Hum Mol Genet* 15:2490-2508.
30
31
32 Springer TA, Zhu J, Xiao T. 2008. Structural basis for distinctive recognition of fibrinogen
33 gammaC peptide by the platelet integrin alphaIIbbeta3. *J Cell Biol* 182:791-800.
34
35
36 Stafford P, Garner SF, Rankin A, Kekomaki R, Watkins NA, Ouwehand WH. 2008. A single-
37 nucleotide polymorphism in the human ITGB3 gene is associated with the platelet-
38 specific alloantigen Va (HPA-17bw) involved in fetal maternal alloimmune
39 thrombocytopenia. *Transfusion* 48:1432-1438.
40
41
42 Tadokoro S, Tomiyama Y, Honda S, Arai M, Yamamoto N, Shiraga M, Kosugi S, Kanakura
43 Y, Kurata Y, Matsuzawa Y. 1998. A Gln747 -->Pro substitution in the α IIb subunit is
44 responsible for a moderate α IIb β ₃ deficiency in Glanzmann thrombasthenia. *Blood*
45 92:2750-2758.
46
47
48 Tadokoro S, Tomiyama Y, Honda S, Kashiwagi H, Kosugi S, Shiraga M, Kiyoi T, Kurata Y,
49 Matsuzawa Y. 2002. Missense mutations in the β ₃ subunit have a different impact on
50 the expression and function between α IIb β ₃ and α _v β ₃. *Blood* 99:931-938.
51
52
53 Valentin N, Visentin GP, Newman PJ. 1995. Involvement of the cystein-rich domain of
54 glycoprotein IIIa in the expression of the human platelet alloantigen, PI^{A1} : Evidence
55 for heterogeneity in the humoral response. *Blood* 85:3028-3033.
56
57
58 Wang GS, Cooper TA. 2007. Splicing in disease: disruption of the splicing code and the
59 decoding machinery. *Nat Rev Genet* 8:749-761.
60
61
62
63
64
65
66
67
68
69
70
71
72
73
74
75
76
77
78
79
80
81
82
83
84
85
86
87
88
89
90
91
92
93
94
95
96
97
98
99
100

1
2
3 glycoprotein IIIa is responsible for the formation of the Pena/Penb alloantigen system.
4 J Clin Invest 90:2038-2043.
5

6
7 Ward CM, Kestin AS, Newman PJ. 2000. A Leu262Pro mutation in the integrin $\beta 3$ subunit in
8 an α IIB- $\beta 3$ complex that binds fibrin but not fibrinogen. Blood 96:161-169.
9

10 Xiao T, Takagi J, Coller BS, Wang JH, Springer TA. 2004. Structural basis for allostery in
11 integrins and binding to fibrinogen-mimetic therapeutics. Nature 432:59-67.
12

13
14 Zhu J, Luo BH, Xiao T, Zhang C, Nishida N, Springer TA. 2008. Structure of a complete
15 integrin ectodomain in a physiologic resting state and activation and deactivation by
16 applied forces. Mol Cell 32:849-861.
17
18
19
20
21
22
23
24
25
26
27
28
29
30
31
32
33
34
35
36
37
38
39
40
41
42
43
44
45
46
47
48
49
50
51
52
53
54
55
56
57
58
59
60

For Peer Review

Figure legends :**Figure 1 : Flow cytometry analysis of the expression of the mutant forms of α IIB β 3.**

Transiently transfected COS-7 cells expressing the wild type (WT) or the following mutants of α IIB β 3, p.S926L, p.Q595H, p.V903F (α IIB) and p.C38Y, p.M118R, p.G221D, p.K253 (β 3) were tested in flow cytometry by using Moabs SZ22 (hatched bar), SZ21 (dot bar) and P2 (white bar) or irrelevant mouse IgG (black bar). As control, untransfected cells (NT) were also tested. Following Moab incubation, bound IgGs were detected by using a polyclonal goat anti-mouse IgG conjugated to FITC. Washed cells were analyzed on an Epic-XML flow cytometer. Except for the α IIB p.Q595H one, all mutants showed a markedly decreased expression of the complex when compared to the wild type. Results represent the mean \pm SD of 3 experiments.

Figure 2 : Western blot analysis of the Y38 β 3 form of the α IIB β 3 complex expressed in COS-7 cells. SDS lysates from normal platelets (lane a), non-transfected COS-7 cells (lane b), or COS-7 cells expressing the wild-type (lane c) or the Y38 β 3 (lane d) forms of the complex were subjected to SDS-polyacrylamide gel electrophoresis under non reduced conditions and transferred to nitrocellulose membrane. Following incubation with Moabs SZ22, XIIF9, SZ21 and AP3, membrane bound IgGs were revealed by using anti-mouse IgG conjugated to peroxidase and the enhanced chemiluminescence detection technique.

Figure 3 : Location and modeling of the electrostatic effect of the p.K253M substitution. (A) and (B) Ribbon diagram of the α IIB β -propeller (in blue) and β 3 β I (in red) complexed domains. For the β I domain, The residue 253 is shown in green with (A) a lysine and (B) a methionine, while for β -propeller partners the P228 is shown in yellow, the F231 in

1
2
3
4
5
6
7
8
9
10
11
12
13
14
15
16
17
18
19
20
21
22
23
24
25
26
27
28
29
30
31
32
33
34
35
36
37
38
39
40
41
42
43
44
45
46
47
48
49
50
51
52
53
54
55
56
57
58
59
60

grey, the W262 in pink and the D232 in orange. (C) and (D) Electrostatics visualization of the single point mutation p.K253M with (C) wild K253 $\beta 3$ and (D) mutated M253 $\beta 3$ subunits. The visualizations are done using PyMol software (<http://www.pymol.org>).

Figure 4 : Localization of the V903 and S926 on a ribbon diagram of α IIIb**.** This α I**IIb** cartoon shows the internal location of the V903 and S926 side chains (orange, ball and stick representation) in the calf-2 composed by the N-terminal part of the β -subunit of α I**IIb** (blue) and the C-terminal part of the α -subunit (yellow and red respectively for β -sheet and α -helix). This ribbon diagram was obtained from the resolved structure of α I**IIb** (PDB code 3FCS) using the Swiss-PdbViewer software (version 4.0.1).

Figure 5 : Localization of the allelic variants of the β -I domain of $\beta 3$. Residues identified in this study and from the literature as involved in GT or neonatal alloimmune thrombocytopenia and that locate in the β -I domain of $\beta 3$ are reported on its primary sequence (panel A) and a ribbon diagrams showing the β -I domain and a partial view of the β -propeller of α I**IIb** (panel B). Allelic variants of residues highlighted in magenta, blue, green are respectively responsible for variant, II and I GT phenotypes or yellow for those concerning alloimmunisation. The GT type induced by the allelic variant of the H192 (grey) is not reported [Peretz et al., 2006]. Amino-acids identified by the energy analyze as in contact with the α I**IIb** β -propeller are underlined. The primary sequence (panel A) identifies the secondary structures [Xiao et al., 2004] and amino-acids responsible for MIDAS (\star), ADMIDAS (\bullet) and SyMBS (\blacklozenge) sites [Springer et al., 2008]. As known, variant GT mutations GT specifically affect metal ion binding sites but other mutants concern residues evenly distributed along the primary sequence. Boxed residues have been identified in this study. The ribbon diagram (panel B) was obtained from the resolved structure of α I**IIb** (PDB file 2VDL) using the Swiss-

1
2
3 PdbViewer software (version 4.0.1). α -helix (red) and β -sheet (yellow) are shown. All
4
5 residues whose mutants are involved in type II GT lay on the β -I surface except for the L117,
6
7 some of them (underlined amino-acids) directly contacting the β -propeller. The Met 118 and
8
9 Gly 221 locate inside the β -I domain and the Lys 253 points out to the β -propeller surface
10
11 however their identified allelic variants completely prevent the α IIb β 3 complex expression at
12
13 the platelet surface.
14
15
16
17
18
19
20
21
22
23
24
25
26
27
28
29
30
31
32
33
34
35
36
37
38
39
40
41
42
43
44
45
46
47
48
49
50
51
52
53
54
55
56
57
58
59
60

For Peer Review

Table 1 : α IIB mutations identified by genomic analysis^a

N ^o	Proband	GT ^b	Ex/IVS ^c	Sequence alteration ^d	Status ^e	Restriction enzyme	Predicted HGVS ^f	Predicted Classical ^g	Accession number ^h
1	CabGT-1	I	Ex 4	c.531T>A	Hom.	Hga I	p.C177*	p.C146*	ss131010618
2	CabGT-2	I	Ex 10	c.917dupA	Het.	Alw I	p.R307Efs*22	p.R276E + FS ^k	ss131007603
3			Ex 19	c.1912dup T	Het.	No enzyme	p.C639Mfs*22	p.C608M + FS ^{k,l}	ss 131007606
4	CabGT-3	I	Ex 13	c.1214T>C	Het.	Sfc I	p.I405T	p.I374T ^l	ss 131007609
5			IVS14	c.1442-13_1442-1del	Het.	Hph I	Splice ⁱ	Splice	ss 131007612
6	CabGT-4	I (car ^w)	IVS14	c.1442-13_1442-1del	Het.	Hph I	Splice	Splice	ss 131007612
7	CabGT-5	I	IVS14	c.1442-13_1442-1del NI ^j	Het.	Hph I	Splice	Splice	ss 131007612
8	CabGT-6	I	Ex 14	c.1413C>G	Het.	Dde I	p.Y471*	p.Y440* ^m	ss 131007615
9			Ex 27	c.2800G>T	Het.	Mae III / Tsp45 I	p.V934F	p.V903F	ss 131007617
10	CabGT-7	I	Ex 16	c.1563T>A NI	Het.	No enzyme	p.C521*	p.C490*	ss 131007620
11	CabGT-8	NC ^x	Ex 17	c.1672C>T	Hom.	Mse I	p.Q558*	p.Q527*	ss 131007623
12	CabGT-9	I	IVS18	c.1879-2A>G	Hom.	Xba I	Splice	Splice	ss 131007625
13	CabGT-10	I	Ex 18	c.1878G>C	Hom.	Hpy CH IV	p.Q626H / Splice	p.Q595H / Splice	ss 131007628
14	CabGT-11	I	Ex 18	c.1787T>C	Hom.	No enzyme	p.I596T	p.I565T ⁿ	ss 131008441
15	CabGT-12	II	Ex 21	c.2113T>C	Het.	Fok I	p.C705R	p.C674R ^{l,o,p}	ss 131007632
16			IVS29	c.3060+2T>C	Het.	Hph I	p.V982_K1020del	p.V951_K989del ^q	ss 131007634
17	CabGT-13	I	Ex 23	c.2333A>C	Hom.	Bsr I / TspR I	p.Q778P	p.Q747P ^r	ss 131007637
18	CabGT-14	I	Ex 27	c.2748_2757del	Hom.	Rsa I	p.T917Sfs181	p.T886S + FS	ss 131007639
19	CabGT-15	I (car)	Ex 27	c.2800G>T	Het.	Mae III / Tsp45 I	p.V934F	p.V903F	ss 131007617
20	CabGT-16	I	Ex 28	c.2870C>T	Hom.	Hpy CH4 V	p.S957L	p.S926L	ss 131007642
21	CabGT-17	II	Ex 28	c.2929C>T	Het.	Ava I	p.R977*	p.R946* ^{l,s,t}	ss 131007644
22			NI						
23	CabGT-18	NC	Ex 29	c.2944G>A	Het.	Bst NI	p.V982M	p.V951M ^{u,v}	ss 131007647
24			Ex 29	c.2965G>A	Het.	Nci I	p.A989T	p.A958T ^{u,v}	ss 131007649
25				NI					

^a cDNA ITGA2B GenBank accession number NM_000419.3^b GT : Glanzmann type

^c Ex : Exon ; IVS : intronic sequence

^d Nomenclature according to HGVS : +1 corresponds to the A of ATG translation initiation codon

^e Status : Het. : heterozygous; Hom. : Homozygous

^f Predicted effect, nomenclature according to HGVS : +1 corresponds to the initiating Met (signal peptide included)

^g Predicted effect, classical nomenclature : +1 corresponds to the first amino acid of the mature α IIb

^h NCBI dSNP accession number

ⁱ Splice defect based upon modeling with GENSCAN / NNSPLICE and ESEFinder

^j NI : 2nd mutation not Identified

^k FS : Frame shift

References : ¹ D'Andrea et al., 2002 [D'Andrea et al., 2002]; ^m Scott et al., 1998 [Scott et al., 1998]; ⁿ Ruan et al., 1998 [Ruan et al., 1998]; ^o Gonzalez-Manchon et al., 1999 [Gonzalez-Manchon et al., 1999]; ^p Mitchell et al., 2003 [Mitchell et al., 2003]; ^q French et al., 1997 [French and Coller, 1997]; ^r Tadokoro et al., 1998 [Tadokoro et al., 1998]; ^s Basani et al., 2000 [Basani et al., 2000]; ^t Peretz et al., 2006 [Peretz et al., 2006]; ^u Nurden et al., 2004 [Nurden et al., 2004]

^v V951M is responsible for GT but not A958T [Nurden et al., 2004]

^w Car : carrier

^x NC : Not communicated

Table 2 : $\beta 3$ mutations identified by genomic analysis^a

N°	Proband	GT ^b	Ex/IVS ^c	Sequence alteration ^d	Status ^e	Restriction enzyme	Predicted HGVS ^f	Predicted Classical ^g	Accession number ^h
19	CabGT-19	NC ^j	IVS2	c.166-2A>G	Hom.	Apa I	Splice ⁱ	Splice	ss 131007652
20	CabGT-20	I	Ex 2	c.100C>T	Hom.	Hpy CH4 IV	p.R34*	p.R8* ^l	ss 131007655
21	CabGT-21	I	Ex 3	c.191G>A NI ^k	Het.	Tsp 45I	p.C64Y	p.C38Y	ss 131007657
22	CabGT-22	I	Ex 4	c.431T>G	Hom.	Bsm F1	p.M144R	p.M118R	ss 131007660
23	CabGT-23	II	Ex 5	c.665 T>C	Hom.	Acy I	p.L222P	p.L196P ^m	ss 131007662
24	CabGT-24	I	Ex 5 Ex 6	c.740 G>A c.836 A>T	Het. Het.	Fok I Fat I	p.G247D p.K279M	p.G221D p.K253M	ss 131007665 ss 131007667
25	CabGT-25	NC	IVS8	c.1125+3_1125+6del NI	Het.	No enzyme	Splice	Splice	ss 131007670
26	CabGT-26	I	Ex 9 Ex 9	c.1143C>A c.1260G>A	Het. Het.	Taq I Hph I	p.K376_T420del	p.K350_T394del ⁿ	ss 131007672 ss 131007674

^acDNA ITGB3 GenBank accession number NM_000212.2

^bGT : Glanzmann type I or II (< 5% or 10 – 25 % of normal platelet α IIB β 3[Caen, 1989]). See Supp_mat tableS1 for detailed data.

^cEx : Exon ; IVS : intronic sequence

^dNomenclature according to HGVS : +1 corresponds to the A of ATG translation initiation codon

^eStatus : Het. : heterozygous; Hom. : Homozygous

^fNomenclature according to HGVS : +1 corresponds to the initiating Met (signal peptide included)

^gClassical nomenclature : +1 corresponds to the first amino acid of the mature $\beta 3$

^hNCBI dSNP accession number

ⁱSplice defect based upon modeling with GENSCAN, NNSPLICE and ESEFinder

^jNC : Not communicated

^kNI : 2nd mutation not Identified

^lReferences : ^lNegrier et al., 1998 [Negrier et al., 1998]; ^mNurden et al., 2002 [Nurden et al., 2002]; ⁿJin et al., 1996 [Jin et al., 1996]

Table 3 : Energetic effects of the Lys 253 Met substitution in the α I**II**b/ β 3 interaction

Energy of :		Lys 253			Met 253		
		Rank	Value (kcal/mol)	α I II b contact	rank	Value (kcal/mol)	α I II b contact
αIIIbβ3 Interface	Electrostatic (4r) Energy	/	-25.14	/	/	-16.08	/
	Desolvation Free Energy	/	-1.09	/	/	-6.81	/
	Van der Waals (CHARMm19)	/	-4894.63	/	/	-4903.95	/
Residue	Binding free energy	1 st negative	-5.541	/	12 th negative	-1.309	/
	Desolvation free energy	1 st positive	4.124	/	6 th negative	-1.347	/
	Electrostatic energy	1 st negative	-9.666	/	/	<i>low</i>	/
Contact	Electrostatic	1 st negative	-6.715	D232	/	/	/
		6 th negative	-2.076	F231	/	/	/
		13 th negative	-0.788	P228	/	/	/
		10 th positive	0.670	W262	6 th negative	-0.935	W262
	Free energy	1 st negative	-5.517	D232	/	/	/
		8 th negative	-1.166	F231	/	/	/
2 th positive		2.057	W262	/	/	/	




Figure 1 : Flow cytometry analysis of the expression of the mutant forms of α IIb β 3.
235x165mm (96 x 96 DPI)





Figure 2 : Western blot analysis of the Y38 β_3 form of the $\alpha_{IIb}\beta_3$ complex expressed in COS-7 cells.
155x177mm (96 x 96 DPI)



CMYK color scale

Figure 3 : Location and modeling of the electrostatic effect of the p.K253M substitution.
193x158mm (96 x 96 DPI)





Figure 3 : Location and modeling of the electrostatic effect of the p.K253M substitution.
193x158mm (96 x 96 DPI)




CYMK color scale

Figure 4 : Localization of the V903 and S926 on a ribbon diagram of aIIb.
135x115mm (96 x 96 DPI)

W



1
2
3
4
5
6
7
8
9
10
11
12
13
14
15
16
17
18
19
20
21
22
23
24
25
26
27
28
29
30
31
32
33
34
35
36
37
38
39
40
41
42
43
44
45
46
47
48
49
50
51
52
53
54
55
56
57
58
59
60



RGB color scale



Figure 4 : Localization of the V903 and S926 on a ribbon diagram of aIIb.
135x115mm (96 x 96 DPI)





CMYK color scale

Figure 5 : Localization of the allelic variants of the β -I domain of β 3.
212x200mm (96 x 96 DPI)



RGB color scale

Figure 5 : Localization of the allelic variants of the β -I domain of β 3.
212x200mm (96 x 96 DPI)