



Genetic interaction between Sox10 and Zfhx1b during enteric nervous system development.

Laure Stanchina, Tom van de Putte, Michel Goossens, Danny Huylebroeck,
Nadège Bondurand

► To cite this version:

Laure Stanchina, Tom van de Putte, Michel Goossens, Danny Huylebroeck, Nadège Bondurand. Genetic interaction between Sox10 and Zfhx1b during enteric nervous system development.. Developmental Biology, 2010, 341 (2), pp.416-28. 10.1016/j.ydbio.2010.02.036 . inserm-00481705

HAL Id: inserm-00481705

<https://inserm.hal.science/inserm-00481705>




Submitted on 7 May 2010

HAL is a multi-disciplinary open access archive for the deposit and dissemination of scientific research documents, whether they are published or not. The documents may come from teaching and research institutions in France or abroad, or from public or private research centers.

L'archive ouverte pluridisciplinaire **HAL**, est destinée au dépôt et à la diffusion de documents scientifiques de niveau recherche, publiés ou non, émanant des établissements d'enseignement et de recherche français ou étrangers, des laboratoires publics ou privés.







Zfhx1b/Sox10







E10.5/E11.5

Zfhx1b/Sox10





E13.5

Zfhx1b/Sox10 Version 0048170510








Zfhx1b/GFAP




Adult

Zfhx1b/GFAP



























Sox10^{LacZ/+}









Sox10^{LacZ/+}; Zfhx1b^{Δ/+}




A**Sox10^{LacZ/+}****Sox10^{LacZ/+}; Zfhx1b^{Δ/+}****B****Wild Type****Zfhx1b^{Δ/+}****Sox10^{LacZ/+}****Sox10^{LacZ/+}; Zfhx1b^{Δ/+}**


A***Sox10*^{LacZ/+}*****Sox10*^{LacZ/+}; *Zfhx1b*^{Δ/+}****A1****A2****A3****A4****C****Wild Type*****Zfhx1b*^{Δ/+}*****Sox10*^{LacZ/+}*****Sox10*^{LacZ/+}; *Zfhx1b*^{Δ/+}****C1****C2****C3****C4**

insertm-00481705, version 2 - 15 Jun 2010

B***Sox10*^{LacZ/+}*****Sox10*^{LacZ/+}; *Zfhx1b*^{Δ/+}****Head****Cranial****Tail**



Sox10^{LacZ/+}



Sox10^{LacZ/+}; Zfhx1b^{Δ/+}





Table 1: Genotypes distribution and viability analysis of animals resulting from $\text{Sox10}^{\text{LacZ}+/+}$ and $\text{Zfhx1b}^{\Delta/+}$ crosses.

Genotypes	Number of animals genotyped	Number of animals dead between P0 and 4 weeks	Number of animals dead between 4 and 6 weeks	% of mortality over total period
$\text{Sox10}^{+/+}; \text{Zfhx1b}^{+/+}$	14	0	0	0%
$\text{Sox10}^{\text{LacZ}+/+}; \text{Zfhx1b}^{+/+}$	13	1	0	8%
$\text{Sox10}^{+/+}; \text{Zfhx1b}^{\Delta/+}$	10	0	0	0%
$\text{Sox10}^{\text{LacZ}+/+}; \text{Zfhx1b}^{\Delta/+}$	10	7	3	100%

Pups mortality was quantified by identifying all animals post mortem or at weaning (number of animals genotyped) and counting the number of animals dead between P0 and 4 weeks or P0 and 6 weeks.