

## Quality of life assessment in HIV clinical research in resource-limited settings: better late than never.

Renaud Becquet, Juan Burgos-Soto, Maria Patrizia Carrieri, Bruno Spire

► **To cite this version:**

Renaud Becquet, Juan Burgos-Soto, Maria Patrizia Carrieri, Bruno Spire. Quality of life assessment in HIV clinical research in resource-limited settings: better late than never.. *Tropical Medicine and International Health*, Wiley-Blackwell, 2010, 15 (9), pp.1008-1010. <10.1111/j.1365-3156.2010.02583.x>. <inserm-00484884>

**HAL Id: inserm-00484884**

**<http://www.hal.inserm.fr/inserm-00484884>**

Submitted on 10 Jan 2012

**HAL** is a multi-disciplinary open access archive for the deposit and dissemination of scientific research documents, whether they are published or not. The documents may come from teaching and research institutions in France or abroad, or from public or private research centers.

L'archive ouverte pluridisciplinaire **HAL**, est destinée au dépôt et à la diffusion de documents scientifiques de niveau recherche, publiés ou non, émanant des établissements d'enseignement et de recherche français ou étrangers, des laboratoires publics ou privés.

**Quality of life assessment  
in HIV clinical research in resource-limited settings: better late than never**

Renaud BECQUET <sup>1,2</sup>, Juan BURGOS-SOTO <sup>1,3</sup>, Maria Patrizia CARRIERI <sup>4,5,6</sup>, Bruno SPIRE <sup>4,5,6</sup>

- <sup>1</sup> INSERM, Unité 897, Centre de Recherche "Epidémiologie et Biostatistique", Bordeaux, France
- <sup>2</sup> Institut de Santé Publique Epidémiologie Développement (ISPED), Université Victor Segalen Bordeaux 2, Bordeaux, France
- <sup>3</sup> ANRS site in Côte d'Ivoire (PAC-CI), Centre Hospitalier Universitaire de Treichville, Abidjan, Côte d'Ivoire
- <sup>4</sup> INSERM, Unité 912, "Economie et Sciences Sociales, Systèmes de Santé et Sociétés", Marseille, France
- <sup>5</sup> Université Aix Marseille, Marseille, France
- <sup>6</sup> Observatoire Régional de la Santé Provence Alpes Côte d'Azur (ORS PACA), Marseille, France

*Corresponding author*

Renaud Becquet

INSERM Unité 897, Institut de Santé Publique Epidémiologie Développement (ISPED),  
Université Victor Segalen Bordeaux 2, 146 rue Léo Saignat, 33076 Bordeaux, France

Tel: +33.(0)5.57.57.45.35, Fax: +33.(0)5.57.57.56.30

E-mail: Renaud.Becquet@isped.u-bordeaux2.fr

## **Summary**

Comprehensive and sustained optimal care for HIV-infected patients can now be achieved in resource-constrained settings thanks to the sustainability of programs providing antiretroviral therapy. However the primary goals of HIV virological suppression and improved survival need to be accompanied by a substantial improvement in patient's experience with HIV care and treatment. An assessment of both patient's quality of life and perceived toxicity and symptoms should now be systematically integrated into HIV clinical research in resource-constrained countries. This will allow treatment strategies aimed at optimizing the durability of response to antiretroviral therapy in these settings to be properly evaluated and compared.

The current scale-up of antiretroviral therapy (ART) in resource-constrained settings presents an unprecedented opportunity to radically reduce the burden of AIDS by improving survival rates and ensuring the virological control of HIV infection (1). ART access has transformed HIV into a chronic disease, but the burden of associated side-effects makes the sustainability of long-life treatment questionable. Despite these therapeutic progresses, stigma is frequently experienced by HIV-infected patients and has an adverse impact on health behaviours such as adherence and systematic condom use (2, 3).

The routine follow-up of HIV-infected patients in resource-constrained settings is solely based on a non-optimal set of clinical and biological indicators aimed at monitoring ART efficacy. This latter however is highly dependent on adherence, which in turn is mainly influenced by patients' perceptions of treatment experience (4). Patient experience with treatment can be measured either by generic (5) or HIV-specific (6) quality of life scales, or by other indicators of perceived health such as the HIV symptom index (7). Quality of life assessment is not only aimed at evaluating individual well-being, but also has public health purposes as high quality of life scores can positively influence health behaviours. Indeed, self-reported symptoms are associated with non-adherence to ART in both high- and low-income countries (8-10). Similarly, poor perceived health has been shown to be a major correlate of unprotected sex among HIV-positive patients in Europe (11, 12), and a similar relationship was recently shown in West-Africa (13).

In high-income countries, a wide variety of powerful ART regimens is available and the assessment of quality of life in clinical research is becoming an additional criterion to guide providers in selecting the best treatment option. For instance, in a recent trial comparing Lopinavir monotherapy with Lopinavir associated triple therapy (10), the number of self-reported symptoms was found to be an important discriminatory measure as it revealed major differences in patient well-being. Similarly, a significant improvement in quality of life scores has been reported in patients switching from enfuvirtide to raltegravir (14). While several trials comparing ART regimens are ongoing in resource-constrained settings, the lack of data collection of patient reported outcomes in such trials could have a serious influence on decisions regarding the best treatment options to take when different treatment strategies show similar efficacy.

To tackle this issue, there are at least three major reasons justifying the implementation of research on quality of life issues in resource-constrained settings. First, it is now conceivable that comprehensive and sustained optimal care can be achieved thanks to the sustainability of ART programs in such settings (15). It is therefore crucial to aim not only at achieving

virological suppression and improving survival, but also at optimizing patient perceived health. Second, the recently released WHO guidelines stress that, in resource-constrained settings, ART should be initiated among all HIV-infected patients with a CD4 count below 350 cells/ml (16). Similarly, all pregnant and breastfeeding women in these settings should now be offered ART to both improve maternal health and prevent mother-to-child transmission of HIV (17, 18). Hence, an increasing proportion of patients will soon be on ART without having had any experience of symptomatic HIV disease. In this context, research on therapeutic strategies which are able to guarantee both efficacy and improvement in quality of life is now urgently required. Third, mathematical models have recently shown that an intervention based on annual universal HIV screening followed by immediate ART initiation for all those infected would dramatically reduce HIV incidence in a generalised epidemic setting (19). One cannot seriously expect this intervention to be widely implemented without an adequate prior ascertainment of both the public and individual benefits. For this reason, quality of life assessment should be introduced in forthcoming population based trials aimed at evaluating this "treatment as prevention" hypothesis.

Tools aimed at measuring quality of life and other patient reported outcomes in the context of HIV infection were initially designed for high-income settings. There are no major methodological barriers to the expansion of these tools for the evaluation of patient well-being in lower income countries, as demonstrated recently by their successful implementation in Cameroon (20).

It would be unethical not to develop research on quality of life and patient perception of ART in developing settings, especially in sub-Saharan Africa. If "do no harm" remains the main principle of all medicine, the integration of quality of life assessment in comprehensive research programs on HIV care is a major ethical and public health priority.

## Acknowledgements

This group was supported by the French National Agency for Research on AIDS and viral hepatitis (ANRS), Paris, France and by the French Charity SIDACTION, Paris, France. We would like to thank Jude Sweeney for his helpful comments on this manuscript.

## References

1. Braitstein P, Brinkhof MW, Dabis F, Schechter M, Boulle A, Miotti P, et al. Mortality of HIV-1-infected patients in the first year of antiretroviral therapy: comparison between low-income and high-income countries. *Lancet*. 2006 Mar 11;367(9513):817-24.
2. Kippax SC, Aggleton P, Moatti JP, Delfraissy JF. Living with HIV: recent research from France and the French Caribbean (VESPA study), Australia, Canada and the United Kingdom. *Aids*. 2007 Jan;21 Suppl 1:S1-3.
3. Peretti-Watel P, Spire B, Obadia Y, Moatti JP. Discrimination against HIV-infected people and the spread of HIV: some evidence from France. *PLoS ONE*. 2007;2(5):e411.
4. Spire B, Duran S, Souville M, Lepout C, Raffi F, Moatti JP. Adherence to highly active antiretroviral therapies (HAART) in HIV-infected patients: from a predictive to a dynamic approach. *Soc Sci Med*. 2002 May;54(10):1481-96.
5. Ware J, Sherbourne C. The MOS 36-item short-form health survey (SF-36) : I. Conceptual framework and item selection. *Med Care*. 1992;30:473-83.
6. Wu AW, Revicki DA, Jacobson D, Malitz FE. Evidence for reliability, validity and usefulness of the Medical Outcomes Study HIV Health Survey (MOS-HIV). *Qual Life Res*. 1997 Aug;6(6):481-93.
7. Justice AC, Holmes W, Gifford AL, Rabeneck L, Zackin R, Sinclair G, et al. Development and validation of a self-completed HIV symptom index. *J Clin Epidemiol*. 2001 Dec;54 Suppl 1:S77-90.
8. Duran S, Spire B, Raffi F, Walter V, Bouhour D, Journot V, et al. Self-reported symptoms after initiation of a protease inhibitor in HIV-infected patients and their impact on adherence to HAART. *HIV Clin Trials*. 2001;2(1):38-45.
9. Marcellin F, Boyer S, Protopopescu C, Dia A, Ongolo-Zogo P, Koulla-Shiro S, et al. Determinants of unplanned antiretroviral treatment interruptions among people living with HIV in Yaounde, Cameroon (EVAL survey, ANRS 12-116). *Trop Med Int Health*. 2008 Dec;13(12):1470-8.
10. Spire B, Marcellin F, Cohen-Codar I, Flandre P, Boue F, Dellamonica P, et al. Effect of lopinavir/ritonavir monotherapy on quality of life and self-reported symptoms among antiretroviral-naïve patients: results of the MONARK trial. *Antivir Ther*. 2008;13(4):591-9.
11. Ammassari A, Murri R, Pezzotti P, Trotta MP, Ravasio L, De Longis P, et al. Self-reported symptoms and medication side effects influence adherence to highly active antiretroviral therapy in persons with HIV infection. *J Acquir Immune Defic Syndr*. 2001 Dec 15;28(5):445-9.
12. Vincent E, Bouhnik AD, Carrieri MP, Rey D, Dujardin P, Granier F, et al. Impact of HAART-related side effects on unsafe sexual behaviours in HIV-infected injecting drug users: 7-year follow up. *Aids*. 2004 Jun 18;18(9):1321-5.
13. Protopopescu C, Marcellin F, Preau M, Gabillard D, Moh R, Minga A, et al. Psychosocial correlates of inconsistent condom use among HIV-infected patients

- enrolled in a structured ART interruptions trial in Cote d'Ivoire: results from the TRIVACAN trial (ANRS 1269). *Trop Med Int Health*. 2010;15(6):706-12.
14. De Castro N, Braun J, Charreau I, Pialoux G, Cotte L, Katlama C, et al. Switch from Enfuvirtide to Raltegravir in Highly Treatment-experienced HIV-1-infected Patients: A Randomized Open-label Non-inferiority Trial, Easier ANRS 138. Abstract 572. The 16th conference on retroviruses and opportunistic infections; 2009; Montreal, Canada; 2009.
  15. UNAIDS. AIDS epidemic update. Geneva, Switzerland: United Nations program on HIV/AIDS; 2009.
  16. WHO. Rapid advice: Antiretroviral therapy for HIV infection in adults and adolescents. 2009 [cited May 5, 2010]; [http://www.who.int/hiv/pub/arv/rapid\\_advice\\_art.pdf](http://www.who.int/hiv/pub/arv/rapid_advice_art.pdf)
  17. Becquet R, Ekouevi DK, Arrive E, Stringer J, Meda N, Chaix ML, et al. Universal antiretroviral therapy for pregnant and breast-feeding HIV-infected women: towards the elimination of mother-to-child transmission of HIV-1 in resource-limited settings. *Clin Infect Dis*. 2009;49(12):1936-45.
  18. WHO. Rapid advice: Recommendations for Use of Antiretroviral Drugs for Treating Pregnant Women and Preventing HIV Infection in Infants. 2009 [cited May 5, 2010]; [http://www.who.int/hiv/pub/mtct/rapid\\_advice\\_mtct.pdf](http://www.who.int/hiv/pub/mtct/rapid_advice_mtct.pdf)
  19. Granich RM, Gilks CF, Dye C, De Cock KM, Williams BG. Universal voluntary HIV testing with immediate antiretroviral therapy as a strategy for elimination of HIV transmission: a mathematical model. *Lancet*. 2009 Jan 3;373(9657):48-57.
  20. Marcellin F, Bonono CR, Blanche J, Carrieri MP, Spire B, Koulla-Shiro S. Higher risk of unsafe sex and impaired quality of life among patients not receiving antiretroviral therapy in Cameroon: results from the EVAL survey (ANRS 12-116). *Aids*. 2010 Jan;24 Suppl 1:S17-25.