

**Combining naturally occurring polyphenols with
TNF-related apoptosis-inducing ligand: a promising
approach to kill resistant cancer cells?**

Guillaume Jacquemin, Sarah Shirley, Olivier Micheau

► **To cite this version:**

Guillaume Jacquemin, Sarah Shirley, Olivier Micheau. Combining naturally occurring polyphenols with TNF-related apoptosis-inducing ligand: a promising approach to kill resistant cancer cells?: TRAIL and polyphenols in anticancer therapy.. Cellular and Molecular Life Sciences, Springer Verlag, 2010, 67 (18), pp.3115-30. <10.1007/s00018-010-0407-6>. <inserm-00491262>

HAL Id: inserm-00491262

<https://www.hal.inserm.fr/inserm-00491262>

Submitted on 11 Jun 2010

HAL is a multi-disciplinary open access archive for the deposit and dissemination of scientific research documents, whether they are published or not. The documents may come from teaching and research institutions in France or abroad, or from public or private research centers.

L'archive ouverte pluridisciplinaire **HAL**, est destinée au dépôt et à la diffusion de documents scientifiques de niveau recherche, publiés ou non, émanant des établissements d'enseignement et de recherche français ou étrangers, des laboratoires publics ou privés.

Fig. 1 Schematic diagram of the TRAIL-induced apoptotic signaling. TRAIL binding on DR4 and/or DR5 triggers the recruitment of the adaptor protein FADD and the subsequent recruitment of the initiator Caspase-8 and/or -10, forming the DISC. The effector Caspase-3 is activated either directly by the initiator caspases (extrinsic pathway), or through the mitochondrial activation of Caspase-9 within the apoptosome (intrinsic pathway). The decoy receptors DcR1 and DcR2, as well as various anti-apoptotic proteins (in red), are involved in the regulation of the TRAIL signaling.

Fig. 2 Chemical structures of polyphenols that exhibit a synergistic effect with TRAIL.

Fig. 3 Chemical structure of the synthetic flavonoid flavopiridol.

Fig. 4 Schematic model of action of polyphenols on the TRAIL signaling pathway. Polyphenols modulate both the extrinsic and intrinsic pathways, and also tumor suppressors such as P53 and survival proteins such as Akt. The molecular targets vary depending on the structure of the polyphenolic compound and the cell type. This model reports only the main molecular targets of polyphenols.