

	aMCI patients		Converters		Non-converters	
	t0	t18	t0	t18	t0	t18
<b>Number</b>	17		7		10	
(women/men)	(11/6)		(5/2)		(6/4)	
<b>Follow-up duration</b> (months; mean ± SD)	17.9 ± 1.0		17.5 ± 0.91		18.2 ± 1.1	
(range)	(17.0 to 20.6)		(17.0 to 18.8)		(17.0 to 20.6)	
<b>Age at t0</b> (years; mean ± SD)	71.6 ± 8.9		73.3 ± 4.3		70.4 ± 11.2	
(range)	(55 to 87)		(73 to 80)		(55 to 87)	
<b>Education</b> (years; mean ± SD)	10.3 ± 3.9		11.0 ± 4.7		9.9 ± 3.6	
(range)	(7 to 17)		(7 to 17)		(7 to 17)	
<b>MMSE</b> (mean ± SD)	27.5 ± 1.2	25.6 ± 3.9 <sup>▣</sup>	26.7 ± 1.0	24.3 ± 5.1	28.1 ± 1.0	26.6 ± 2.6 <sup>▣</sup>
(range)	(25 to 29)	(17 to 30)	(25 to 28)	(17 to 29)	(26 to 29)	(23 to 30)
<b>MMSE-PAC</b> (mean ± SD)	-4.3 ± 9.1		-6.9 ± 12.3		-2.5 ± 6.1	
(range)	(-30.0 to 5.3)		(-30.0 to 3.0)		(-9.8 to 5.3)	
<b>Mattis</b> (mean ± SD)	137.1 ± 4.4	130.5 ± 10.1 <sup>▣▣</sup>	135.7 ± 4.9	121.3 ± 9.5 <sup>▣▣▣</sup>	138.1 ± 3.9	136.6 ± 4.3
(range)	(129 to 144)	(108 to 142)	(129 to 141)	(108 to 137)	(133 to 144)	(128 to 142)
<b>Mattis-PAC</b> (mean ± SD)	-3.3 ± 4.8		-7.1 ± 3.4		-0.7 ± 3.8	
(range)	(-11.8 to 2.9)		(-11.8 to -2.5)		(-7.4 to 2.9)	

**Table 1: Demographic and clinical data of aMCI patients at baseline (t0) and at follow-up (t18).** \*\* significant difference between converters and non-converters p<0.01, ▣ significant difference between t0 and t18 p<0.05, ▣▣ p<0.01, ▣▣▣ p<0.001.