



HAL
open science

Platelet serotonin level predicts survival in amyotrophic lateral sclerosis.

Luc Dupuis, Odile Spreux-Varoquaux, Gilbert Bensimon, Philippe Jullien, Lucette Lacomblez, François Salachas, Gaëlle Bruneteau, Pierre-François Pradat, Jean-Philippe Loeffler, Vincent Meininger

► **To cite this version:**

Luc Dupuis, Odile Spreux-Varoquaux, Gilbert Bensimon, Philippe Jullien, Lucette Lacomblez, et al.. Platelet serotonin level predicts survival in amyotrophic lateral sclerosis.. PLoS ONE, Public Library of Science, 2010, 5 (10), pp.e13346. 10.1371/journal.pone.0013346 . inserm-00531253

HAL Id: inserm-00531253

<https://www.hal.inserm.fr/inserm-00531253>

Submitted on 2 Nov 2010

HAL is a multi-disciplinary open access archive for the deposit and dissemination of scientific research documents, whether they are published or not. The documents may come from teaching and research institutions in France or abroad, or from public or private research centers.

L'archive ouverte pluridisciplinaire **HAL**, est destinée au dépôt et à la diffusion de documents scientifiques de niveau recherche, publiés ou non, émanant des établissements d'enseignement et de recherche français ou étrangers, des laboratoires publics ou privés.

Platelet Serotonin Level Predicts Survival in Amyotrophic Lateral Sclerosis

Luc Dupuis^{1,2*}, Odile Spreux-Varoquaux^{3,4,5}, Gilbert Bensimon^{6,7,8}, Philippe Jullien⁵, Lucette Lacomblez^{6,9,10}, François Salachas⁹, Gaëlle Bruneteau⁹, Pierre-François Pradat⁹, Jean-Philippe Loeffler^{1,2}, Vincent Meininger^{6,9}

1 INSERM U692, Strasbourg, France, **2** Université de Strasbourg, Strasbourg, France, **3** Faculté de Médecine Paris-Ile de France-Ouest, Paris, France, **4** Université de Versailles Saint-Quentin-en-Yvelines, Versailles, France, **5** Centre Hospitalier Versailles, Le Chesnay, France, **6** Université Pierre et Marie Curie, Paris, France, **7** UMR 7211, CNRS, Paris, France, **8** Service de Pharmacologie Clinique, Hôpital de la Pitié-Salpêtrière (AP-HP), Paris, France, **9** Département des Maladies du Système Nerveux, Centre Référent Maladie Rare SLA Hôpital de la Pitié-Salpêtrière (AP-HP), Paris, France, **10** INSERM U678, Paris, France

Abstract

Background: Amyotrophic lateral sclerosis (ALS) is a life-threatening neurodegenerative disease involving upper and lower motor neurons loss. Clinical features are highly variable among patients and there are currently few known disease-modifying factors underlying this heterogeneity. Serotonin is involved in a range of functions altered in ALS, including motor neuron excitability and energy metabolism. However, whether serotonergic activity represents a disease modifier of ALS natural history remains unknown.

Methodology: Platelet and plasma unconjugated concentrations of serotonin and plasma 5-HIAA, the major serotonin metabolite, levels were measured using HPLC with coulometric detection in a cohort of 85 patients with ALS all followed-up until death and compared to a control group of 29 subjects.

Principal Findings: Platelet serotonin levels were significantly decreased in ALS patients. Platelet serotonin levels did not correlate with disease duration but were positively correlated with survival of the patients. Univariate Cox model analysis showed a 57% decreased risk of death for patients with platelet serotonin levels in the normal range relative to patients with abnormally low platelet serotonin ($p = 0.0195$). This protective effect remained significant after adjustment with age, gender or site of onset in multivariate analysis. Plasma unconjugated serotonin and 5-HIAA levels were unchanged in ALS patients compared to controls and did not correlate with clinical parameters.

Conclusions/Significance: The positive correlation between platelet serotonin levels and survival strongly suggests that serotonin influences the course of ALS disease.

Citation: Dupuis L, Spreux-Varoquaux O, Bensimon G, Jullien P, Lacomblez L, et al. (2010) Platelet Serotonin Level Predicts Survival in Amyotrophic Lateral Sclerosis. PLoS ONE 5(10): e13346. doi:10.1371/journal.pone.0013346

Editor: Christophe Egles, Tufts University, United States of America

Received: July 14, 2010; **Accepted:** September 20, 2010; **Published:** October 13, 2010

Copyright: © 2010 Dupuis et al. This is an open-access article distributed under the terms of the Creative Commons Attribution License, which permits unrestricted use, distribution, and reproduction in any medium, provided the original author and source are credited.

Funding: Dr. Dupuis received research support from the Amyotrophic Lateral Sclerosis Association (Grant 1698, ALSA). Drs. Dupuis, Lacomblez, Pradat and Pr Meininger received support from the association pour la recherche sur la sclérose latérale amyotrophique (ARSLA). Drs. Loeffler, Lacomblez and Pradat received research support from the Association Française contre les Myopathies (AFM). Pr Meininger received research support from Teva Pharma, GSK and Trophos. Dr. Lacomblez received research support from Trophos, Eisai, GSK, Novartis and Janssen-Cilag. The funders had no role in study design, data collection and analysis, decision to publish, or preparation of the manuscript.

Competing Interests: Pr Meininger received research support from Teva Pharma, GSK and Trophos. Dr. Lacomblez received research support from Trophos, Eisai, GSK, Novartis and Janssen-Cilag. This does not alter the authors' adherence to all the PLoS ONE policies on sharing data and materials.

* E-mail: ldupuis@unistra.fr

Introduction

Amyotrophic lateral sclerosis (ALS) is a neurodegenerative disorder affecting both lower motor neurons (LMN) and upper motor neurons (UMN), and leads to death within 2 to 5 years of onset. Clinical features and progression are highly heterogeneous among patients (for review see [1]). There are currently few identified disease-modifying factors accounting for this heterogeneity. Among these, energy metabolism abnormalities have been shown in ALS patients [2,3,4,5] and their potential contribution to the course of the disease has been stressed [6], but little is known about the factors triggering these abnormalities.

The neurotransmitter serotonin is involved in a range of functions altered in ALS, including motor neuron excitability and

energy metabolism (reviewed in [7,8,9]). Early studies on very limited numbers of patients suggested that the levels of serotonin and its metabolite are decreased in brain tissues of ALS patient's post-mortem [10,11,12]. Moreover, recent imaging studies have shown decreased binding of serotonin 1A (5-HT_{1A}) ligands in ALS raphe and cortex [13]. However, definite evidence that serotonin itself is modified in ALS is lacking and whether serotonergic activity is a disease modifier of ALS natural history remains unknown. Since serotonergic neurons and platelets express similar serotonin related enzymes and receptor, alterations in central serotonin are likely to be reflected in platelet serotonin levels [14,15,16]. In this study we aimed to study the relationships between circulating serotonin and survival in a cohort of ALS patients.

Results

Decreased platelet serotonin levels in ALS patients

The demographic data of ALS patients (Table 1) are in close accordance with previous reports thus showing that the patients included in the present study are representative of the ALS population. Age and sex ratio of ALS and control groups did not differ (Table 1). Levels of platelet serotonin were significantly lower in ALS patients (median [range] ng/mL: 78.4 [5.2–232.1]) as compared to controls (median [range] ng/mL: 110.6 [41.4–239.6], $p = 0.0003$) with a global 30% decrease in ALS patients (Table 2). Normal values of the laboratory range from 60 to 200 ng/mL ($N = 69$ patients) [17,18]. Compared to normal value of the laboratory, 31% of ALS patients displayed values below the normal range, as compared to 6% of controls. Plasma unconjugated serotonin and 5-HIAA remained unchanged. The decreased platelet serotonin levels were found in patients with either bulbar or spinal onset (Table 3). Levels of plasma unconjugated serotonin were decreased in bulbar, but not in spinal onset patients compared to controls (Table 3). 5-HIAA plasma levels were similar in controls and in patients whatever was the site of onset, and the molar ratio between 5-HIAA and platelet serotonin (an index of MAO-A activity) was significantly increased in ALS patients compared to matched controls (Table 2), and this was mostly due to bulbar onset patients. This difference between bulbar and spinal onset patients was not due to a difference in nutritional status between these two groups since BMI did not differ significantly (bulbar onset patients: 24.6 ± 3.5 ; spinal onset patients: 24.7 ± 3.4 ; $p > 0.05$).

Decreased platelet serotonin is predictive of worsened survival of ALS patients

There was a significant decrease of platelet serotonin with increasing age at blood sampling in ALS patients (R squared = 0.053, $p = 0.0365$) but not in controls (R squared = 0.004825, $p = 0.72$). In the ALS patients group, there was no relationship of platelet serotonin with the length of disease prior to sampling (R squared = 0.016324 $p = 0.2527$). Platelet serotonin levels showed no correlation with either site of onset, sex, Norris score slopes, initial BMI or rate of BMI loss. Neither plasma serotonin nor 5-HIAA levels correlated with any of the clinical parameters.

Overall median survival, calculated from time of sampling to death, was 34 months [range: 1–149]. Univariate Cox model analysis of survival showed a significant increase of risk of death with increasing age (Cox model, RR [95%CI] (per year) = 1.028 [1.004–1.055], $p = 0.0210$) and significant decrease of the risk of death with increasing platelet serotonin levels (Cox model, RR [95%CI] (per

Table 1. Demographic parameters in ALS and control groups.

Item (+/–SD)	ALS patients	Controls
N	85	29
Mean age at blood sampling (years)	61.65 ± 10.5	61.0 ± 14.4
Sex ratio (M/F)	1.24	1.18
Bulbar site of onset (bulbar/spinal)	34% (29/56)	
Age at onset (years)	60.3 ± 9.9	
Age at death (years)	64.20 ± 9.8	
Survival (months)	45.9 ± 40.7	
BMI (kg/m ²)	24.7 ± 3.5	

doi:10.1371/journal.pone.0013346.t001

Table 2. Serotonin levels in ALS and control groups.

Variables (mean ± SD; n)	ALS patients	Controls	P value
Platelet serotonin (ng/mL)	83.4 ± 50.3; 82	124.3 ± 47.7; 29	0.0003
Plasma unconjugated serotonin (ng/mL)	1.35 ± 1.78; 82	1.41 ± 1.20; 29	NS
Plasma 5-HIAA (ng/mL)	7.35 ± 4.07; 72	7.71 ± 2.38; 29	NS
5-HIAA/platelet serotonin molar ratio	0.14 ± 0.20; 70	0.07 ± 0.04; 29	0.012

Groups were compared using Mann-Whitney test and the p-value is shown. doi:10.1371/journal.pone.0013346.t002

ng/mL) = 0.994 [0.989–0.999], $p = 0.0195$). We broke down patient population in 3 groups according to platelet serotonin levels. Values were considered as in a high normal range when superior to 100 ng/mL (i.e. approximately the mean of controls), in an abnormal low range when inferior to 50 ng/mL, and in a borderline range between 50 and 100 ng/mL. This analysis showed a good linearity of the platelet serotonin levels on survival (Figure 1). Patients in the high normal values range showed a 57% decreased risk of death relative to the abnormal low values range and 35% relative to the borderline one ($p = 0.023$ by the log-rank test, Figure). Platelet serotonin remained a significant independent predictor of survival following adjustment on known prognostic variables such as age, or site of onset or gender (RR [95%CI] (per ng/mL) = 0.995 [0.990–0.999], $p = 0.046$; RR [95%CI] (per ng/mL) = 0.984 [0.989–0.999], $p = 0.020$ and RR [95%CI] (per ng/mL) = 0.995 [0.989–0.999], $p = 0.026$ respectively).

Discussion

Our study provides two important results, (i) platelet serotonin levels are significantly decreased in ALS patients relative to matched controls, and (ii) platelet serotonin level is a significant predictor of survival independent of age.

We first show that platelet serotonin levels are globally decreased in ALS patients, with about one third of ALS patients displaying levels of platelet serotonin below the normal range.

Platelets are usually considered as an easily accessible model of central serotonergic neurons due to similarities in enzymatic and receptor equipments [14,16]. Thus, the decreased platelet serotonin levels could mirror what has been reported in ALS

Table 3. Serotonin levels in ALS patients as a function of site of onset.

Item (+/–SD; n)	ALS patients (Bulbar onset)	ALS patients (Spinal onset)	Controls
Platelet serotonin (ng/mL)	76.9 ± 47.5; 28**	86.8 ± 51.7; 54**	124.3 ± 47.7; 29
Plasma unconjugated serotonin (ng/mL)	0.88 ± 1.08; 26*	1.56 ± 2.00; 56	1.41 ± 1.20; 29
Plasma 5-HIAA (ng/mL)	8.00 ± 3.31; 23	7.04 ± 4.37; 49	7.71 ± 2.38; 29
5-HIAA/platelet serotonin molar ratio	0.21 ± 0.30; 22*	0.12 ± 0.13; 49	0.07 ± 0.04; 29

Groups were compared using Kruskal Wallis test. *, $p < 0.05$; **, $p < 0.01$. doi:10.1371/journal.pone.0013346.t003




Figure 1. Correlation between platelet serotonin levels and survival in patients with amyotrophic lateral sclerosis (ALS). Kaplan-Meier survival curves of patients with ALS stratified according to their platelet serotonin levels. Survival was calculated as the time period between blood sampling and death. The black line represents patients with ALS having abnormal serotonin values (serotonin <50 ng/mL, n = 22 patients); light grey line represents patients with high normal values (serotonin >100 ng/mL i.e. median of control group, n = 25 patients), and light black line patients with borderline values (50–100 ng/mL, n = 35 patients). Group comparison shows a statistically significant difference ($p = 0.024$ by the log-rank).

doi:10.1371/journal.pone.0013346.g001

patients, i.e., a spinal cord decrease in both serotonin and its metabolite as well as a decreased binding of 5-HT_{1A} ligand in the brain [9,10,11,12,13]. Patients with bulbar onset displayed a more profound impairment of the serotonergic system since plasma unconjugated serotonin levels were also decreased in this population, in contrast with patients with spinal onset. This is in line with a more pronounced loss of cortical 5-HT_{1A} binding observed in bulbar onset patients [13].

The causes of serotonin decrease remain elusive. Number of neurologic and psychiatric conditions, including Alzheimer's disease [19], are associated with decreases of the serotonergic system either centrally or in platelets or both. A similar decrease in serotonin has been described in frontotemporal dementia [20], which is now thought to form a continuum with ALS [21]. Platelet serotonin decrease was also found in depression, although to a lesser extent [18,22,23] and in violent suicide attempters [24]. Consistent with a pathogenic role of serotonin decrease in ALS, number of studies showed that depression in ALS worsens disease severity [25,26,27]. In all, our current study supports the existence of a marked impairment of the serotonergic system in a significant proportion of ALS patients. It should be pointed out that decreased levels of platelet serotonin have been reported in many different pathological or debilitating conditions, notably in psychiatric and neurologic diseases, and thus platelet serotonin levels in ALS, as observed decreased here, should not be considered as a specific biomarker of ALS. Another limitation of our study is that serotonin levels were assayed in blood, at one single time point. Future studies should be longitudinal and measure brain-derived serotonin in the CSF.

The second major result of this study is that platelet serotonin levels correlated positively with survival, as calculated from time of sampling to death. Numbers of studies have shown that serotonin levels drop with age [28]. Since age is a known modifier of ALS survival, [29] it was critical to determine whether the observed correlation was related to age and not to the disease process. In our

study, platelet serotonin did not correlate with age in controls. This is likely to be due to the relatively narrow age range of patients, since the most important variations of serotonin with age occur in the first two decades of life [30]. In the ALS population, after adjustment with age in a multivariate Cox model, platelet serotonin remained a significant predictor of survival, thus showing that the correlation between platelet serotonin and survival unlikely results from an effect of age on platelet serotonin levels. Another potential confounding factor in studies with serotonin is drug interactions. The observed correlations were not due to such drug interactions since patients taking serotonin reuptake inhibitors, antidepressants or any other serotonin related medications were excluded from the study. It should be noted that in this study, serotonin was measured at one single time point, when disease had already been diagnosed. Thus, serotonin decrease might be either preexisting disease onset and be associated with an intrinsic variation in serotonin metabolism, or be the consequence of the disease process itself. The observed positive correlation might, in this latter case be due to an upstream mechanism able to modify both ALS course and serotonin levels. In this case, serotonin decrease would be a surrogate marker of a crucial, still unknown, pathogenic event. In the current absence of longitudinal studies or of genetic studies on serotonin-related genes, it remains impossible to discriminate between these two hypothesis.

Previous research in animal models of ALS however supports that serotonin might directly modulate onset and survival in ALS. The administration of serotonin precursor 5-hydroxytryptophan delayed onset and mortality in a transgenic mice ALS model [31]. The mechanisms underlying the potential protection by serotonin in ALS clearly deserve further investigation. It seems unlikely that the potential positive effect of serotonin on survival is related to the excitatory action on motor neurons [7,8,32] since increased serotonin potentially exacerbates a glutamate-evoked excitotoxicity [9]. A more attractive hypothesis is a relationship between serotonin and energy metabolism. Serotonin modulates energy

homeostasis through complex and still incompletely characterized mechanisms [33]. A large number of ALS patients as well as transgenic ALS mice show increased energy expenditure [3,4,5,34], a phenotype which appears analogous to the effects of a chronic depletion in brain-derived serotonin [35]. Future studies should focus on the potential relationships between serotonin and energy metabolism in ALS.

Whether serotonin directly modulates disease course or constitute a surrogate marker of another crucial pathogenic event, our results show that platelet serotonin correlates with survival in ALS. To our knowledge, few biological factors have been demonstrated to be associated with survival in ALS. Among them are the increase of LDL/HDL ratio [6], of plasma ApoE [36], the polymorphisms in the *kifap3* gene [37] or more recently CSF glial markers [38]. In the current state of knowledge, our results do not indicate that serotonin replacement therapy could be useful for patients, but rather that investigations focused on the serotonergic system in ALS are warranted.

Materials and Methods

Ethics statement

Patients were informed of the aim of this non-interventional research and gave their written informed consent. The institutional review board of Salpêtrière hospital approved the research. Control patients gave their written informed consent for participating in this study.

Patients

Blood samples were collected from 90 patients followed at the Pitié Salpêtrière hospital (Paris, France) between 1994 and 1995. All patients met the El Escorial World Federation of Neurology criteria for the diagnosis of definite or probable ALS at the time of blood sampling. Patients underwent blood sampling after diagnosis and before riluzole treatment. Control samples were collected from 11 healthy volunteers and 18 patients hospitalized for orthopaedic surgery. There was no difference in any of the measured parameters between healthy volunteers and orthopaedic patients and they were thus considered as a single control group. Controls were age and sex matched and medical examinations excluded neurodegenerative, inflammatory, psychiatric (in particular depression and food intake disorders, such as anorexia) or neoplastic disorders of the nervous system. All blood samples were taken as a part of routine laboratory evaluations and were collected after a 12 h fasting period, with care taken to avoid possible medications and possible diets and dietary problems known to interfere with serotonin levels. For controls undergoing orthopaedic surgery, blood samples were taken before any surgery to avoid for the potential biases due to surgery or immobilization. All individuals included in the study, either patients or controls, did not use serotonin reuptake inhibitors or any medication with known interaction with serotonin (*e.g.* antidepressants). Death was documented by death certificate or written letter from relatives or physicians. Documented time of death was obtained for all patients but 5 who were lost to follow-up and had no information after sampling visit. Database was closed at 1st October 2009 and these 5 patients were excluded from the study. Survival was defined as time from blood sampling to death. Disease duration was calculated as the time interval between first symptoms and blood sampling. These values were calculated in months. Body mass index (BMI) was calculated at each visit and BMI loss was calculated using a linear regression model. Bulbar and Limbs Norris scale scores were assessed at sampling visit and at each subsequent visit.

Biochemical procedures

Circulating serotonin was measured in serum; while plasma unconjugated serotonin and 5-HIAA levels were measured in plasma. Platelet serotonin, which corresponds to about 98% of total circulating serotonin, was calculated as the difference between circulating and plasma unconjugated serotonin concentrations. The molar ratio between platelet serotonin and 5-HIAA levels was calculated and used as an indirect index of MAO-A activity. All the patients and the control subjects fasted overnight. Samples of blood were drawn from the antecubital vein between 7AM and 10AM. Five millilitres of blood were collected into Vacutainer tubes with no additive for circulating 5-HT, with citrate for plasma unconjugated serotonin and with dry heparin for plasma 5-HIAA levels. Serum and plasma were separated by centrifugation at 3000 g and stored at -80°C until further analysis without additional thawing. The results were unrelated to storing time. The three parameters were determined within 6 months after blood sampling by high-performance liquid chromatography with coulometric detection according to three different analytical procedures previously described [39,40]. The minimum quantifiable level of serotonin was 0.05 ng/mL in plasma and serum. The day-to-day variations were 2.0%, 3.0% and 2.7% for 25, 100 and 300 ng/mL respectively ($n=8$) for serum serotonin and 8.6%, 7.0% and 5.6% for 0.5, 5 and 10 ng/mL plasma serotonin levels ($n=8$). For plasma 5-HIAA assays, the minimum quantifiable level was 1 ng/mL with the day-to-day coefficient of variation of 14% and 11.2% for 6 and 21 ng/mL ($n=10$) respectively.

Statistical analysis

Data are expressed as mean \pm SD and/or median [range].

Based on normal values of the laboratory (mean [95% CI]: 92.7 [83.4–101.9] ng/ml) [17,18] we calculated that we could detect at least a 26% change in patients ($n=85$) as compared to controls ($n=29$) with 81% power and p -alpha (2-tailed) = 0.05, by the T test. For group comparisons (ALS *vs.* controls), we performed non-parametric Mann-Whitney tests. For comparisons of three groups (control *vs.* ALS patients with spinal onset or bulbar onset), we used Kruskal Wallis tests. Correlations were performed using the nonparametric Spearman's rank correlation test.

Univariate and multivariate Cox model analysis were performed to assess the effects of candidate variables on survival. To assess the linearity of platelet serotonin levels on survival, patient population was broken-down in 3 groups according to (i) high normal values (cut-off $>$ median of control group = 100 ng/mL *i.e.* circa 550 nM, 25 patients), (ii) abnormal low values (cut-off below the normal range <50 ng/mL *i.e.* circa 280 nM, 22 patients) and (iii) borderline values (50–100 ng/mL, 35 patients). Survival curves for the three groups were compared by the Mantel Cox (log rank) statistic. Data were analysed using JMP software (version 8.0; SAS Institute Inc., Cary, NC), and statistical threshold for statistical significance set at p (α 2-tailed) <0.05 .

Acknowledgments

The authors thank the patients and their families. Anne Grégoire is acknowledged for her technical HPLC assistance and Marie-José Ruivo and Annie Picchinenna for technical assistance.

Author Contributions

Conceived and designed the experiments: OSV GB VM. Performed the experiments: OSV. Analyzed the data: LD OSV GB VM. Contributed reagents/materials/analysis tools: OSV GB PJ LL FS GB PFP JPL VM. Wrote the paper: LD.

References

- Ravits JM, La Spada AR (2009) ALS motor phenotype heterogeneity, focality, and spread: deconstructing motor neuron degeneration. *Neurology* 73: 805–811.
- Pradat PF, Bruneteau G, Gordon PH, Dupuis L, Bonnefont-Rousselot D, et al. (2009) Impaired glucose tolerance in patients with amyotrophic lateral sclerosis. *Amyotroph Lateral Scler*. pp 1–6.
- Desport JC, Preux PM, Magy L, Boirie Y, Vallat JM, et al. (2001) Factors correlated with hypermetabolism in patients with amyotrophic lateral sclerosis. *Am J Clin Nutr* 74: 328–334.
- Desport JC, Preux PM, Truong TC, Vallat JM, Sautereau D, et al. (1999) Nutritional status is a prognostic factor for survival in ALS patients. *Neurology* 53: 1059–1063.
- Funalot B, Desport JC, Sturtz F, Camu W, Couratier P (2008) High metabolic level in patients with familial amyotrophic lateral sclerosis. *Amyotroph Lateral Scler*. pp 1–5.
- Dupuis L, Corcia P, Fergani A, Gonzalez De Aguilar JL, Bonnefont-Rousselot D, et al. (2008) Dyslipidemia is a protective factor in amyotrophic lateral sclerosis. *Neurology* 70: 1004–1009.
- Heckman CJ, Mottram C, Quinlan K, Theiss R, Schuster J (2009) Motoneuron excitability: The importance of neuromodulatory inputs. *Clin Neurophysiol*.
- Rekling JC, Funk GD, Bayliss DA, Dong XW, Feldman JL (2000) Synaptic control of motoneuronal excitability. *Physiol Rev* 80: 767–852.
- Sandyk R (2006) Serotonergic mechanisms in amyotrophic lateral sclerosis. *Int J Neurosci* 116: 775–826.
- Bertel O, Malessa S, Sluga E, Hornykiewicz O (1991) Amyotrophic lateral sclerosis: changes of noradrenergic and serotonergic transmitter systems in the spinal cord. *Brain Res* 566: 54–60.
- Forrest V, Ince P, Leitch M, Marshall EF, Shaw PJ (1996) Serotonergic neurotransmission in the spinal cord and motor cortex of patients with motor neuron disease and controls: quantitative autoradiography for 5-HT_{1A} and 5-HT₂ receptors. *J Neurol Sci* 139 Suppl: 83–90.
- Sofic E, Riederer P, Gsell W, Gavranovic M, Schmidtko A, et al. (1991) Biogenic amines and metabolites in spinal cord of patients with Parkinson's disease and amyotrophic lateral sclerosis. *J Neural Transm Park Dis Dement Sect* 3: 133–142.
- Turner MR, Rabiner EA, Hammers A, Al-Chalabi A, Grasby PM, et al. (2005) [11C]-WAY100635 PET demonstrates marked 5-HT_{1A} receptor changes in sporadic ALS. *Brain* 128: 896–905.
- Lesch KP, Wolozin BL, Murphy DL, Reiderer P (1993) Primary structure of the human platelet serotonin uptake site: identity with the brain serotonin transporter. *J Neurochem* 60: 2319–2322.
- Sneddon JM (1973) Blood platelets as a model for monoamine-containing neurones. *Prog Neurobiol* 1: 151–198.
- Ramamoorthy S, Bauman AL, Moore KR, Han H, Yang-Feng T, et al. (1993) Antidepressant- and cocaine-sensitive human serotonin transporter: molecular cloning, expression, and chromosomal localization. *Proc Natl Acad Sci U S A* 90: 2542–2546.
- Alvarez JC, Cremniter D, Lesieur P, Gregoire A, Gilton A, et al. (1999) Low blood cholesterol and low platelet serotonin levels in violent suicide attempters. *Biol Psychiatry* 45: 1066–1069.
- Alvarez JC, Gluck N, Arnulf I, Quintin P, Leboyer M, et al. (1999) Decreased platelet serotonin transporter sites and increased platelet inositol triphosphate levels in patients with unipolar depression: effects of clomipramine and fluoxetine. *Clin Pharmacol Ther* 66: 617–624.
- Muck-Seler D, Presecki P, Mimica N, Mustapic M, Pivac N, et al. (2009) Platelet serotonin concentration and monoamine oxidase type B activity in female patients in early, middle and late phase of Alzheimer's disease. *Prog Neuropsychopharmacol Biol Psychiatry* 33: 1226–1231.
- Huey ED, Putnam KT, Grafman J (2006) A systematic review of neurotransmitter deficits and treatments in frontotemporal dementia. *Neurology* 66: 17–22.
- Strong MJ (2008) The syndromes of frontotemporal dysfunction in amyotrophic lateral sclerosis. *Amyotroph Lateral Scler* 9: 323–338.
- Owens MJ, Nemeroff CB (1994) Role of serotonin in the pathophysiology of depression: focus on the serotonin transporter. *Clin Chem* 40: 288–295.
- Csernansky JG, Sheline YI (1993) Abnormalities of serotonin metabolism and nonpsychotic psychiatric disorders. *Ann Clin Psychiatry* 5: 275–281.
- Spreux-Varoquaux O, Alvarez JC, Berlin I, Batista G, Despiere PG, et al. (2001) Differential abnormalities in plasma 5-HIAA and platelet serotonin concentrations in violent suicide attempters: relationships with impulsivity and depression. *Life Sci* 69: 647–657.
- Hillemacher T, Grassel E, Tigges S, Bleich S, Neundorfer B, et al. (2004) Depression and bulbar involvement in amyotrophic lateral sclerosis. *Amyotroph Lateral Scler Other Motor Neuron Disord* 5: 245–249.
- McDonald ER, Wiedenfeld SA, Hillel A, Carpenter CL, Walter RA (1994) Survival in amyotrophic lateral sclerosis. The role of psychological factors. *Arch Neurol* 51: 17–23.
- Pailisse C, Lacomblez L, Dib M, Bensimon G, Garcia-Acosta S, et al. (2005) Prognostic factors for survival in amyotrophic lateral sclerosis patients treated with riluzole. *Amyotroph Lateral Scler Other Motor Neuron Disord* 6: 37–44.
- Guicheney P (1988) Human platelet serotonin content: methodological aspects and physiological variations. *Methods Find Exp Clin Pharmacol* 10: 253–258.
- Chio A, Logroscino G, Hardiman O, Swinger R, Mitchell D, et al. (2009) Prognostic factors in ALS: A critical review. *Amyotroph Lateral Scler* 10: 310–323.
- Flachaire E, Beney C, Berthier A, Salandre J, Quincy C, et al. (1990) Determination of reference values for serotonin concentration in platelets of healthy newborns, children, adults, and elderly subjects by HPLC with electrochemical detection. *Clin Chem* 36: 2117–2120.
- Turner BJ, Lopes EC, Cheema SS (2003) The serotonin precursor 5-hydroxytryptophan delays neuromuscular disease in murine familial amyotrophic lateral sclerosis. *Amyotroph Lateral Scler Other Motor Neuron Disord* 4: 171–176.
- Dunkley EJ, Isbister GK, Sibbritt D, Dawson AH, Whyte IM (2003) The Hunter Serotonin Toxicity Criteria: simple and accurate diagnostic decision rules for serotonin toxicity. *QJM* 96: 635–642.
- Tecott LH (2007) Serotonin and the orchestration of energy balance. *Cell Metab* 6: 352–361.
- Dupuis L, Oudart H, Rene F, Gonzalez de Aguilar JL, Loeffler JP (2004) Evidence for defective energy homeostasis in amyotrophic lateral sclerosis: benefit of a high-energy diet in a transgenic mouse model. *Proc Natl Acad Sci U S A* 101: 11159–11164.
- Yadav VK, Oury F, Suda N, Liu ZW, Gao XB, et al. (2009) A serotonin-independent mechanism explains the leptin regulation of bone mass, appetite, and energy expenditure. *Cell* 138: 976–989.
- Lacomblez L, Doppler V, Beucler I, Costes G, Salachas F, et al. (2002) APOE: a potential marker of disease progression in ALS. *Neurology* 58: 1112–1114.
- Landers JE, Melki J, Meininger V, Glass JD, van den Berg LH, et al. (2009) Reduced expression of the Kinesin-Associated Protein 3 (KIFAP3) gene increases survival in sporadic amyotrophic lateral sclerosis. *Proc Natl Acad Sci U S A* 106: 9004–9009.
- Sussmuth SD, Sperfeld AD, Hinz A, Bretschneider J, Endruhn S, et al. CSF glial markers correlate with survival in amyotrophic lateral sclerosis. *Neurology* 74: 982–987.
- Alvarez JC, Gluck N, Fallet A, Gregoire A, Chevalier JF, et al. (1999) Plasma serotonin level after 1 day of fluoxetine treatment: a biological predictor for antidepressant response? *Psychopharmacology (Berl)* 143: 97–101.
- Radat F, Berlin I, Spreux-Varoquaux O, Elatki S, Ferreri M, et al. (1996) Initial monoamine oxidase-A inhibition by moclobemide does not predict the therapeutic response in patients with major depression. A double blind, randomized study. *Psychopharmacology (Berl)* 127: 370–376.