



HAL
open science

Combined analysis of three genome-wide association studies on vWF and FVIII plasma levels.

Guillemette Antoni, Tiphaine Oudot-Mellakh, Apostolos Dimitromanolakis, Marine Germain, William Cohen, Philip Wells, Mark Lathrop, France Gagnon, Pierre-Emmanuel Morange, David-Alexandre Tregouet

► To cite this version:

Guillemette Antoni, Tiphaine Oudot-Mellakh, Apostolos Dimitromanolakis, Marine Germain, William Cohen, et al.. Combined analysis of three genome-wide association studies on vWF and FVIII plasma levels.. BMC Medical Genetics, 2011, 12 (1), pp.102. 10.1186/1471-2350-12-102 . inserm-00617599

HAL Id: inserm-00617599

<https://inserm.hal.science/inserm-00617599>

Submitted on 29 Aug 2011

HAL is a multi-disciplinary open access archive for the deposit and dissemination of scientific research documents, whether they are published or not. The documents may come from teaching and research institutions in France or abroad, or from public or private research centers.

L'archive ouverte pluridisciplinaire **HAL**, est destinée au dépôt et à la diffusion de documents scientifiques de niveau recherche, publiés ou non, émanant des établissements d'enseignement et de recherche français ou étrangers, des laboratoires publics ou privés.

RESEARCH ARTICLE

Open Access

Combined analysis of three genome-wide association studies on vWF and FVIII plasma levels

Guillemette Antoni^{1,2,3†}, Tiphaine Oudot-Mellakh^{2†}, Apostolos Dimitromanolakis³, Marine Germain^{1,2}, William Cohen^{4,5}, Philip Wells⁶, Mark Lathrop⁷, France Gagnon³, Pierre-Emmanuel Morange^{4,5} and David-Alexandre Tregouet^{1,2*}

Abstract

Background: Elevated levels of factor VIII (FVIII) and von Willebrand Factor (vWF) are well-established risk factors for cardiovascular diseases, in particular venous thrombosis. Although high, the heritability of these traits is poorly explained by the genetic factors known so far. The aim of this work was to identify novel single nucleotide polymorphisms (SNPs) that could influence the variability of these traits.

Methods: Three independent genome-wide association studies for vWF plasma levels and FVIII activity were conducted and their results were combined into a meta-analysis totalling 1,624 subjects.

Results: No single nucleotide polymorphism (SNP) reached the study-wide significance level of 1.12×10^{-7} that corresponds to the Bonferroni correction for the number of tested SNPs. Nevertheless, the recently discovered association of *STXBP5*, *STX2*, *TC2N* and *CLEC4M* genes with vWF levels and that of *SCARA5* and *STAB2* genes with FVIII levels were confirmed in this meta-analysis. Besides, among the fifteen novel SNPs showing promising association at $p < 10^{-5}$ with either vWF or FVIII levels in the meta-analysis, one located in *ACCN1* gene also showed weak association ($P = 0.0056$) with venous thrombosis in a sample of 1,946 cases and 1,228 controls.

Conclusions: This study has generated new knowledge on genomic regions deserving further investigations in the search for genetic factors influencing vWF and FVIII plasma levels, some potentially implicated in VT, as well as providing some supporting evidence of previously identified genes.

Background

Elevated plasma levels of factor VIII (FVIII) and von Willebrand factor (vWF), two key molecules of the coagulation cascade, are well-established risk factors for venous thrombosis (VT) [1-3]. More recent evidence shows that these plasma hemostatic proteins are also risk factors for other cardiovascular diseases (CVD) [4-8]. The broader role of FVIII and vWF is further supported by studies showing that genetic factors modulating the variability of these proteins are also associated with CVD. These include single nucleotide polymorphisms (SNPs) at the *BAI3* [9], *LDLR* [5,10], *VWF* [4] and

ABO [11] genes, the latter being associated with other quantitative risk factors for CVD [12,13].

The estimated heritability of FVIII and vWF levels range between 40% and 60% [14,15] among which about 20% is attributable to the *ABO* locus. A genome wide association study (GWAS) within the CHARGE consortium [16] has recently identified five new genes, apart from their structural genes and *ABO*, consistently influencing vWF and/or FVIII plasma levels. These include *CLEC4M*, *SCARA5*, *STX2*, *STXBP5* and *TC2N*, collectively explaining ~10% of the variability of each two traits. These observations suggest that there are additional genetic factors remaining to be identified and contributing to the hidden heritability of these quantitative traits.

* Correspondence: david.tregouet@upmc.fr

† Contributed equally

¹UMR_S 937, INSERM, Boulevard de l'Hopital, Paris, 75013, France

Full list of author information is available at the end of the article

The increased power of selected samples has long been recognized in family-based studies but more recently the putative advantages of carefully selected samples for quantitative trait analysis of unrelated subjects has also been highlighted [17]. Therefore, we undertook the combined analysis of individual data from three GWAS performed in samples of VT patients and in extended families ascertained on VT and Factor V Leiden (FVL) to identify novel genetic factors implicated in the variation of plasma levels of FVIII and vWF.

Methods

Overall strategy

To achieve our primary goal of identifying new genetic factors that could influence vWF and/or FVIII plasma levels, we used data from three carefully selected independent GWAS. Great attention was drawn to the homogeneity across samples in terms of - ethnic background (most individuals were of French origin), - exclusion criteria with respect to rare forms of inherited thrombophilia, - objectively diagnosed VT, - studied intermediate phenotypes (although some adjustments were done) and similar genotyping technologies (Illumina platform).

In the context of quantitative trait GWAS, individual genetic effect sizes are known to be small [18] and it is expected that a number of real associations do not reach genome-wide significance. Therefore, as part of our analytic strategy, we first tested for association in the individual studies, and results observed across samples were combined into a meta-analysis. We then focused on the consistency of associations across studies as our hypothesis was that real associations would more likely be consistently observed across studies given that each study samples were quite homogeneous with respect to the above-mentioned characteristics. Previously reported associations were also investigated using the above strategy.

As genetic variants associated to plasma levels of FVIII and vWF could be risk factors for VT, our secondary goal was to test the identified SNPs with VT using an *in silico* GWAS [19]. Analytic approaches and samples characteristics of the FVIII and vWF GWAS are described below.

FVL-families sample

Five extended French-Canadian families were ascertained through single probands with idiopathic VT diagnosed at the Thrombosis Clinic of the Ottawa Hospital, and carrying the FVL mutation. VT cases secondary to cancer as well as rare forms of inherited VT (protein S, protein C, AntiThrombin deficiencies) were excluded. A pedigree was drawn from interviews with each potential probands. The largest families were invited to participate

in the study - the family size and willingness to participate being the only criteria for the selection of the families (see Additional File 1, File S1 for the used questionnaire). The total number of family members was 255. Description of the extended families has been published elsewhere [9].

MARTHA samples

The MARseille THrombosis Association (MARTHA) project is composed of two independent samples of VT patients, named MARTHA08 (N = 1,006) and MARTHA10 (N = 586). MARTHA subjects are unrelated caucasians consecutively recruited at the Thrombophilia center of La Timone hospital (Marseille, France) between January 1994 and October 2005. All patients had a documented history of VT and free of well characterized genetic risk factors including AT, PC, or PS deficiency, homozygosity for FV Leiden or FII 20210A, and lupus anticoagulant. They were interviewed by a physician on their medical history, which emphasized manifestations of deep vein thrombosis and pulmonary embolism using a standardized questionnaire (see Additional file 2, File S2). The thrombotic events were confirmed by venography, Doppler ultrasound, spiral computed tomographic scanning angiography, and/or ventilation/perfusion lung scan. All the subjects were of European origin, with the majority being of French descent.

The main characteristics of the three samples are shown in Table 1.

In silico GWAS study on VT

In a previously published GWAS on VT [19], 419 early age of onset and the idiopathic character of VT (ie without environmental risk factors) (< 50 years) VT cases were compared to 1,228 healthy controls at 291,872

Table 1 Main Characteristics of the Studied Samples

	FVL Families N = 253	MARTHA08 N = 972	MARTHA10 N = 570
Age (SD)	40.4 (17.9)	45.7 (14.9)	49.2 (15.7)
Sex (% female)	50.6%	70.8%	58.2%
Smoking (%)	24.4%	24.9%	22.71%
History of VT (%)	5.95%	100%	100%
PT G20210A carriers	0.40%	15.9%	10.6%
FV Leiden carriers	24.9%	26.6%	14.1%
ABO blood group (%)			
O	40.6%	22.9%	22.4%
A	57.8%	61.8%	59.3%
B	1.6%	10.3%	14.4%
AB	-	5%	3.9%
FVIII (SD) IU/dL	118.6 (38.51)	138.70 (55.34)	130.2 (46.35)
vWF (SD) IU/dL	130.3 (53.24)	152.33 (68.23)	152.9 (63.93)

SNPs. Cases were patients from four different French medical centers (Grenoble, Marseille, Montpellier, Paris) selected according to the same criteria as the MARTHA samples, except with the restriction on age of onset. Controls were French subjects selected from the SUVI-MAX population [20].

Measurements

In the French-Canadian (FVL) sample, plasma levels of FVIII activity were measured by a clotting assay on the BCS instrument (Siemens Diagnostics, Marburg Germany) and vWF antigen was measured with a commercially available ELISA kit from Diagnostica Stago. The interassay coefficients of variation for FVIII were ~ 1% and 6.1% for vWF.

In MARTHA subjects, plasma coagulant activity and vWF antigen were assayed in an automated coagulometer (STA-R; Diagnostica Stago, Asnières, France). The interassay coefficients of variation for FVIII and vWF were 6.96% and 2.27% respectively.

Genotyping

The French-Canadian sample was genotyped with the Illumina 660W-Quad Beadchip. The raw datafile contained data for 547,886 autosomal SNPs genotyped on 255 individuals. From these SNPs, 490,083 passed the quality control (QC) criteria of genotyping rate > 90% and more than 20 observations of the minor allele among all individuals. After removing the 88,390 SNPs that failed QC, the overall genotyping rate was 99.88%. The maximum missing rate per sample for all the 255 samples was 3.9%, with an average missing rate of 0.13%. The family structures had previously been checked using 1079 microsatellite markers and RELPAIR [9]. To further verify the correctness of the family structure, we used PREST [21] and computed IBD estimates for all the sample pairs, within and across pedigrees. PREST reported 14,949 Mendelian errors, which is equivalent to a very low Mendelian error rate of 0.012% among all genotypes. Genotypes showing Mendelian inconsistencies were excluded from the analysis. Finally, phenotypic and genotypic data were available on a total of 253 individuals.

The MARTHA08 study sample was typed in 2008 with the Illumina Human610-Quad Beadchip containing 567,589 autosomal SNPs while the MARTHA10 sample was recently typed (beginning of 2010) with the same Illumina Human660W-Quad Beadchip as in the FVL study sample. SNPs showing significant ($P < 10^{-5}$) deviation from Hardy-Weinberg equilibrium, with minor allele frequency (MAF) less than 1% or genotyping call rate < 99%, in each study were filtered out. Individuals with genotyping success rates less than 95% were excluded from the analyses, as well as individuals

demonstrating close relatedness as detected by pairwise clustering of identity by state distances (IBS) and multi-dimensional scaling (MDS) implemented in PLINK software [22]. Non-European ancestry was also investigated using the Eigenstrat program [23] leading to the final selection of 972 and 570 patients left for analysis in MARTHA08 and MARTHA10, respectively. Plasma vWF levels were available in 834 and 537 MARTHA08 and MARTHA10 patients, respectively; corresponding numbers were 541 and 548 for plasma FVIII levels. A total of 442,728 SNPs were common to the three GWAS datasets (see Additional file 3, Figure S1).

Statistical analysis

In the FVL families, association of SNPs with vWF and FVIII levels was tested by means of measured genotype linear association analysis as implemented in the SOLAR (version 4.0, <http://solar.txbiomedgenetics.org/download.html>) program. In MARTHA subjects, association was tested using linear model as implemented in the PLINK program [22].

In order to handle differences in phenotype distributions across studies (Figure 1), and any possible deviation from normality, plasma levels of vWF and FVIII were first normalized before any statistical analysis using the normal quantile transformation [24], separately in the French-Canadian sample, MARTHA08 and MARTHA10. This transformation assigns to each observed measurement the quantile value of the standard normal distribution that corresponds to the rank of this measurement in the original untransformed distribution. Transformed variables are then normally distributed making linear models applicable, and linear regression coefficients comparable across studies. Association analyses were then carried out on the transformed variables assuming additive allele effects (0,1, 2 coding according to the number of minor alleles), and adjusting for age, sex and ABO blood group as tagged by the ABO rs8176746, rs8176704 and rs505922 [19].




Figure 1 Box Plot Distribution of FVIII (left) and vWF (right) Plasma Levels in the Three GWAS Datasets.

When appropriate, haplotype association analyses were carried out in MARTHA samples using THESIAS software [25] to handle the correlation between SNPs, that is linkage disequilibrium (LD). This widely used software implements a stochastic-EM algorithm that simultaneously estimates the frequencies and the effect on the studied phenotype of each inferred haplotype. Haplotype - phenotype associations are then assessed by means of likelihood ratio tests.

Results obtained in each GWAS datasets were combined in a meta-analysis using the GWAMA program [26] <http://www.sph.umich.edu/csg/abecasis/metal>. Both fixed-effect and random-effect models-based analyses were conducted. Regression coefficients characterizing the minor allele effect of each SNP were then combined (after having checked that the minor allele was the same in the different populations) using the inverse-variance method to provide an overall allelic estimate. All reported P values were 2-sided.

Results

A total of 442,728 QC-validated SNPs were common to the three GWAS and were tested through a meta-analysis for association with vWF and FVIII plasma levels. Quantile-quantile plots did not reveal any inflation from what was expected under the null hypothesis of no association (Figure 2), and no SNP reached the study-wide significance level of 1.12×10^{-7} that corresponds to the Bonferroni correction for the number of tested SNPs. Applying the less stringent Sidak correction corresponding to a significant threshold of $p = 1.16 \times 10^{-7}$ would not have modified this conclusion. We then further focused on genetic effects that were consistent across studies and with combined p-value of less than 10^{-5} . As fixed-effect and random-effect analyses provided similar results for most of the main associations (Tables 2 & 3),

the following discussion is based on results obtained from the fixed-effect model analysis.

Ten SNPs covering seven different genes (Figure 3 - Table 2) were associated with plasma vWF levels at $p < 10^{-5}$ with no strong evidence for heterogeneity across GWAS as the lowest Mantel-Haenszel observed p-value, $p = 0.036$, for the ANKDR6 rs645764 would not pass multiple testing correction for testing ten SNPs. The strongest association was observed for rs379440 ($P = 9.82 \times 10^{-6}$) mapping the *EPB41L4A* gene (Table 2). Another SNP at this locus was also associated with vWF, rs13361927 ($P = 4.51 \times 10^{-6}$), but its association was due to its complete LD with rs379440, with pairwise r^2 of 0.78, 0.69 and 0.62 in FVL, MARTHA08 and MARTHA10, respectively. Other vWF-associated SNPs included the *SAFB2* rs732505 ($P = 9.38 \times 10^{-6}$), *VPS8* rs4686760 ($P = 1.08 \times 10^{-6}$) and the *KRT18P24* rs1757948 ($P = 7.37 \times 10^{-6}$). The last three SNPs, rs1438993, rs10745527, rs2579103 (with $P \sim 6 \times 10^{-6}$), were located at the 12q21.33 locus with no known mapped gene and were in nearly complete association. Altogether, the independent signals derived from the rs4686760, rs379440, rs1757948, rs10745527 and rs732505 explained up to 5.7% and 3.8% of the variability of plasma vWF levels in MARTHA08 and MARTHA10, respectively, and 5.3% in the pooled MARTHA samples.

None of the ten vWF-associated SNPs were associated with plasma FVIII levels (all $p > 0.05$). However, six additional SNPs were specifically associated to FVIII levels with homogeneous effects (Mantel-Haenszel p-value > 0.05) across studies (Figure 4 - Table 3). The strongest effect ($P = 2.95 \times 10^{-6}$) was observed for rs7306642, a non synonymous Pro2039Thr variant within the *STAB2* gene, which was one of the recently identified genes by the CHARGE consortium. However, our hit rs7306642 was not in LD with any of the two *STAB2* SNPs recently identified, rs4981022 ($r^2 < 0.01$ in the three studies) and rs4981021 that served as a proxy for rs12229292 ($r^2 < 0.07$ in the three studies). Other FVIII-associated SNPs included the rs6708166 ($P = 1.30 \times 10^{-6}$) in the proximity of *LBH*, the rs1321761 ~ 300 kb apart from *FAM46A* ($P = 9.54 \times 10^{-6}$) and the intronic *VAV2* rs12344583 ($P = 7.92 \times 10^{-6}$) (Table 3). Lastly, two SNPs within the *ACCNI* gene, rs1354492 and rs12941510, were found modulating FVIII plasma levels, the A allele of the former being associated with increased FVIII levels ($\beta = +0.16$, $P = 2.42 \times 10^{-6}$) and the A allele of the latter being associated with decreased levels ($\beta = -0.17$, $P = 5.67 \times 10^{-6}$). These two SNPs were in complete negative LD generating three haplotypes, the sole carrying the rs1354492-A allele being associated with highest levels (see Additional file 4, Table S1). Altogether, these five SNPs (i.e. rs6708166, rs1321761, rs12344583, rs7306642, rs1354492) explained 8.2% and




Figure 2 Quantile-Quantile Plots of the Association Results from the Meta-Analysis of the Three GWAS Datasets.

Table 2 Ten SNPs Showing Association with vWF levels Across the Three GWAS Datasets With Combined Significance P-value < 10⁻⁵

Gene	SNP	Alleles*	MAF ⁺	β (SE)	p	I ²	P _{het}	Random Effect		Fixed Effect		
								β (SE)	p	β (SE)	p	
VPS8	rs4686760	A/G	FVL	0.47	-0.16 (0.08)	0.044	0	0.549	0.15 (0.03)	1.10 10 ⁻⁶	-0.15 (0.03)	1.08 10 ⁻⁶
			Martha08	0.46	-0.18 (0.04)	4.11 10 ⁻⁵						
			Martha10	0.45	-0.11 (0.05)	0.047						
EPB41L4A	rs13361927	G/A	FVL	0.15	0.44 (0.11)	3.08 10 ⁻⁴	0.53	0.119	-0.28 (0.09)	0.002	0.28 (0.06)	4.51 10 ⁻⁶
			Martha08	0.06	0.28 (0.09)	0.003						
			Martha10	0.05	0.11 (0.11)	0.316						
	rs379440	A/G	FVL	0.12	0.46 (0.12)	8.35 10 ⁻⁴	0	0.502	-0.34 (0.07)	9.99 10 ⁻⁷	0.34 (0.07)	9.82 10 ⁻⁷
			Martha08	0.04	0.31 (0.11)	0.004						
			Martha10	0.03	0.25 (0.14)	0.071						
ANKRD6	rs6454764	C/T	FVL	0.04	-0.01 (0.21)	0.977	0.70	0.036	-0.29 (0.14)	0.035	0.31 (0.07)	5.12 10 ⁻⁶
			Martha08	0.06	0.24 (0.09)	0.007						
			Martha10	0.05	0.54 (0.12)	8.97 10 ⁻⁶						
KRT18P24	rs1757948	T/G	FVL	0.27	0.34 (0.09)	2.82 10 ⁻⁴	0.62	0.071	-0.18 (0.06)	0.003	0.15 (0.03)	7.37 10 ⁻⁶
			Martha08	0.27	0.1 (0.05)	0.030						
			Martha10	0.30	0.15 (0.06)	0.009						
desert	rs1438993	G/A	FVL	0.19	0.15 (0.1)	0.127	0	0.666	-0.16 (0.03)	6.34 10 ⁻⁶	0.16 (0.03)	6.25 10 ⁻⁶
			Martha08	0.28	0.18 (0.05)	1.11 10 ⁻⁴						
			Martha10	0.27	0.12 (0.06)	0.052						
	rs10745527	T/G	FVL	0.20	0.19 (0.1)	0.062	0	0.663	-0.16 (0.03)	5.51 10 ⁻⁶	0.16 (0.03)	5.43 10 ⁻⁶
			Martha08	0.28	0.18 (0.05)	1.63 10 ⁻⁴						
			Martha10	0.27	0.11 (0.06)	0.056						
rs2579103	T/G	FVL	0.18	0.17 (0.11)	0.098	0	0.533	-0.16 (0.04)	7.72 10 ⁻⁶	0.16 (0.04)	7.61 10 ⁻⁶	
		Martha08	0.26	0.19 (0.05)	8.24 10 ⁻⁵							
		Martha10	0.25	0.1 (0.06)	0.090							
CDH2	rs2298574	A/G	FVL	0.04	-0.02 (0.19)	0.905	0.19	0.290	0.26 (0.07)	1.81 10 ⁻⁴	-0.27 (0.06)	5.67 10 ⁻⁶
			Martha08	0.08	-0.34 (0.08)	2.77 10 ⁻⁵						
			Martha10	0.07	-0.24 (0.1)	0.022						
SAFB2	rs732505	G/A	FVL	0.05	0.32 (0.18)	0.080	0	0.929	-0.25 (0.06)	9.50 10 ⁻⁶	0.25 (0.06)	9.38 10 ⁻⁶
			Martha08	0.09	0.24 (0.08)	0.001						
			Martha10	0.08	0.25 (0.1)	0.013						

*Common/rare alleles

⁺ Allele frequency of the minor allele

4.6% of the variability of FVIII levels in MARTHA08 and MARTHA10, respectively, and 6.3% in the combined MARTHA samples.

We then used our GWAS datasets to investigate SNPs that had previously been reported associated with vWF and/or FVIII [4,5,9,16]. As shown in Supplementary Table two, marginal associations (P < 0.05) with vWF levels at *STXBPS*, *VWF*, *STX2*, *TC2N* and *CLEC4M* were also observed in our study, the strongest (P = 1.3 10⁻⁴) being for SNP rs216335 at the structural *VWF* gene. All these associations were consistent (i.e the same allele was associated with a genetic effect in the same direction on the studied phenotype) with those previously reported. Together, these associations explained an additional 1.4% and 3.2% of the variance of

plasma levels of vWF in MARTHA08 and MARTHA10, respectively. We did not observe any evidence for an effect of *STAB2* rs4981022 or *BAI3* rs9363864, while the effect of *SCARA5* rs2726953 was heterogeneous across the studies. For FVIII levels, we observed marginal associations of *SCARA5* rs9644133 (P = 0.009) and *VWF* rs1063856 (P = 0.020) that were consistent with those previously reported (Table 4), these two SNPs explaining 0.7% and 0.2% of FVIII variability in MARTHA08 and MARTHA10, respectively. No trend for association was observed for the previously reported associations with *STXBPS*, *STAB2* nor *LDLR* SNPs (Table 5).

We have recently observed that, among the newly identified vWF and/or FVIII genes by the CHARGE consortium, *TC2N* could also be associated with VT risk [27]. Therefore

Table 3 Six SNPs Showing Association with FVIII Activity Across the Three GWAS Datasets With Combined Significance P-value < 10⁻⁵

Gene	SNP	Alleles*	MAF [†]	β (SE)	p	I ²	P _{het}	Random Effect		Fixed Effect		
								β (SE)	p	β (SE)	p	
LBH	rs6708166	G/A	FVL	0.41	-0.12 (0.09)	0.156	0	0.478	-0.17 (0.04)	1.32 10 ⁻⁶	-0.17 (0.04)	1.30 10 ⁻⁶
			Martha08	0.40	-0.23 (0.06)	8.98e-05						
			Martha10	0.42	-0.15 (0.05)	0.007						
FAM46A	rs1321761	T/C	FVL	0.42	-0.20 (0.08)	0.014	0	0.451	-0.15 (0.04)	9.67 10 ⁻⁶	-0.15 (0.04)	9.54 10 ⁻⁶
			Martha08	0.45	-0.10 (0.06)	0.074						
			Martha10	0.47	-0.19 (0.05)	5.93e-04						
VAV2	rs12344583	A/G	FVL	0.17	0.28 (0.11)	0.012	0	0.716	0.20 (0.04)	8.03 10 ⁻⁶	0.20 (0.04)	7.92 10 ⁻⁶
			Martha08	0.20	0.19 (0.07)	0.006						
			Martha10	0.18	0.17 (0.07)	0.012						
STAB2	rs7306642	C/A	FVL	0.16	0.52 (0.12)	1.36e-05	0.59	0.086	0.31 (0.10)	0.002	0.30 (0.06)	2.95 10 ⁻⁶
			Martha08	0.07	0.22 (0.11)	0.057						
			Martha10	0.07	0.20 (0.1)	0.052						
ACCN1	rs1354492	G/A	FVL	0.53	0.09 (0.08)	0.293	0.39	0.192	0.16 (0.04)	5.47 10 ⁻⁶	0.16 (0.03)	2.41 10 ⁻⁶
			Martha08	0.49	0.23 (0.05)	1.20e-05						
			Martha10	0.47	0.12 (0.05)	0.027						
	rs12941510	G/A	FVL	0.22	-0.29 (0.1)	0.004	0.12	0.321	-0.17 (0.04)	2.18 10 ⁻⁵	-0.17 (0.04)	5.67 10 ⁻⁶
			Martha08	0.31	-0.17 (0.06)	0.002						
			Martha10	0.33	-0.12 (0.06)	0.029						

*Common/rare alleles

† Allele frequency of the minor allele


we investigated the effect of the SNPs identified in our meta-analysis on the risk of VT. Our working hypothesis was that SNPs associated with increased (decreased, resp.) plasma levels of these two molecules could be associated with increased (decreased, resp.) risk of disease. For this, we used the results of our previously published GWAS based on 419 VT patients and 1228 healthy subjects (*in silico* association) [19]. As indicated in Table 6, only two SNPs, *VPS8* rs4686760 and *ACCN1* rs12941510, showed some trend of association consistent with our hypothesis. The rs4686760-G allele found associated with decreased vWF levels was slightly less frequent in VT patients than in controls (0.441 vs 0.475, $P = 0.101$) and the rs12941510-A allele, associated with decreased FVIII levels, was also less frequent in cases than in controls (0.310 vs 0.350, $P = 0.046$). These associations can only be considered as suggestive as they would not pass correction for multiple testing. Nevertheless, the observed homogeneity of the allele frequencies of these two SNPs across all genotyped patients is noteworthy. Combining all the VT patients ($n = 1946$), and comparing to the healthy controls of the *in silico* GWAS, the association of rs4686760 with VT remained (0.454 vs 0.475, $P = 0.108$), and that of rs12941510 was strengthened (0.314 vs 0.348, $P = 0.0056$) (Table 7).

Discussion

Theoretically, a sample size of 1,624 unrelated individuals should have a power of 95% to detect, at the

significant level of $1.12 \cdot 10^{-7}$, the additive allele effect of a SNP explaining at least 3% of the variability of a quantitative trait [28]. This power would decrease to 86% and 66% for a SNP explaining 2.5% and 2%, respectively. Our meta-analysis of 1,624 carefully selected samples did not reveal any genome-wide significant association suggesting that the additional common SNPs tagged by current GWAS array and influencing vWF and FVIII plasma levels left to be identified would, if any, individually explain less than 2% of the variability of these two traits.


By lowering the statistical stringency to $p < 10^{-5}$ but focusing on the homogeneity of the effects observed in three independent samples, we identified several novel candidate genes that could contribute to modulate the variability of vWF and FVIII, and that deserve to be further studied. The novel candidate genes for vWF are *VPS8*, *EBP41L4A*, *KRT18P24*, *SAFB2* and a region on 12q21.3 where no known gene maps. Unfortunately, little is known about the biology of the associated proteins and their role in cardiovascular diseases. Among these, *VPS8* stands out. The rs4686760-G allele of the *VPS8* gene, which was associated with decreased vWF levels, was also observed less frequently in VT cases than in healthy controls (0.45 vs 0.48) in the *in silico* GWAS, although this observation did not reach significance ($P = 0.10$). The vacuolar protein sorting 8 homolog gene (*VPS8*) is involved in protein traffic between the golgic



appartus and the vacuole [29] and could participate to the regulation of urokinase-type plasminogen activator [30], the latter known to be involved in thrombosis.

For FVIII levels, the candidate genes identified in our study were *LBH*, *FAM46A*, *VAV2*, *STAB2* and *ACCNI*. Both *LBH* and *VAV2* genes are thought to be involved in angiogenesis. The transcriptional cofactor limb-bud-and-heart (*Lbh*) was discovered as a small acidic nuclear protein highly conserved among species [31]. It has been demonstrated a dramatic suppression of VEGF mRNAs in cells that overexpress *Lbh* [32]. *Vav2* is a guanine nucleotide exchange factor for Rho family proteins. The expression of a dominant negative form of *Vav2* suppress the Vascular Endothelial-Protein Tyrosine Phosphatase (VE-PTP)-induced changes in endothelial cell morphology, such changes being implicated in regulation of angiogenesis [33].

Interestingly, we had previously shown that *STAB2* was located within a linkage peak for vWF levels in our FVL extended families [9] while almost



concomitantly *STAB2* SNPs were found associated with both FVIII and vWF in the CHARGE consortium GWAS [16]. However, the non-synonymous rs7306642 (Pro2039Thr) found associated here with FVIII levels did not show a homogeneous effect on vWF levels across the three GWAS datasets (data not shown), and was in very low LD with others *STAB2* SNPs found associated with these plasma levels. The substitution of a Proline by a Threonine at position 2039 is predicted to be damaging according to web resources <http://genetics.bwh.harvard.edu/pph/index.html>; <http://www.rostlab.org/services/SNAP>. Investigating the effect of this substitution on VT risk would have been relevant but the corresponding SNP did not pass quality control in our *in silico* GWAS. These observations nevertheless suggest that an in-depth haplotype analysis of the *STAB2* gene are required to gain better insight into which SNPs more likely influence plasma levels of FVIII and/or vWF.

ACCNI, encoding an amiloride-sensitive cation channel implicated in cell growth and migration [34], is another gene that deserves greater attention as its genetic variability was found here associated with both FVIII levels and VT risk. However, the SNP that seemed to modulate FVIII levels the most, rs1354492, was not the one that showed association with the disease. This could suggest that either different SNPs distinctly influence plasma levels and VT risk, or that the identified SNPs are in LD with unmeasured variant(s) that could simultaneously influence both phenotypes.

Table 4 Association of Previously Identified SNPs with vWF Levels in the three GWAS Datasets

Gene	SNP	Alleles*	MAF [†]	β (SE)	p	I ²	p _{het}	Random Effect		Fixed Effect		
								β (SE)	p	β (SE)	p	
BAI3	rs9363864	A/G	FVL	0.42	0.04 (0.08)	0.618	0	0.838	0.02 (0.03)	0.461	0.02 (0.03)	0.461
			Martha08	0.52	0.03 (0.04)	0.421						
			Martha10	0.49	-0.002 (0.05)	0.973						
STXBP5	rs9390459	G/A	FVL	0.43	-0.08 (0.08)	0.366	0	0.545	-0.09 (0.03)	0.005	-0.09 (0.03)	0.005
			Martha08	0.42	-0.06 (0.04)	0.197						
			Martha10	0.43	-0.13 (0.05)	0.011						
SCARA5	rs10866867 ⁽¹⁾	G/T	FVL	0.20	-0.08 (0.10)	0.446	0.71	0.03	0.05 (0.07)	0.466	0.09 (0.04)	0.015
			Martha08	0.25	0.17 (0.05)	4.88e-04						
			Martha10	0.25	0.01 (0.06)	0.830						
VWF	rs216335 ⁽²⁾	G/A	FVL	0.06	-0.28 (0.19)	0.141	0	0.945	-0.23 (0.06)	1.31 10 ⁻⁴	-0.23 (0.06)	1.30 10 ⁻⁴
			Martha08	0.08	-0.23 (0.08)	0.003						
			Martha10	0.06	-0.21 (0.11)	0.059						
VWF	rs1063856 ⁽³⁾	A/G	FVL	0.45	0.07 (0.08)	0.371	0	0.889	0.09 (0.03)	0.006	0.09 (0.03)	0.006
			Martha08	0.37	0.08 (0.05)	0.094						
			Martha10	0.38	0.11 (0.05)	0.041						
VWF	rs7306706	A/G	FVL	0.48	-0.04 (0.08)	0.612	0	0.754	0.01 (0.03)	0.664	0.01 (0.03)	0.664
			Martha08	0.45	0.02 (0.04)	0.634						
			Martha10	0.46	0.03 (0.05)	0.604						
STAB2	rs4981022	T/C	FVL	0.30	-0.05 (0.09)	0.601	0	0.541	-0.01 (0.03)	0.664	-0.01 (0.03)	0.664
			Martha08	0.30	0.02 (0.05)	0.652						
			Martha10	0.28	-0.06 (0.06)	0.333						
STX2	rs4334059 ⁽⁴⁾	C/T	FVL	0.33	0.01 (0.09)	0.863	0.01	0.363	0.1 (0.03)	0.004	0.1 (0.03)	0.003
			Martha08	0.37	0.08 (0.04)	0.067						
			Martha10	0.36	0.15 (0.06)	0.008						
TC2N	rs2402074 ⁽⁵⁾	G/A	FVL	0.52	0.05 (0.08)	0.548	0	0.509	0.07 (0.03)	0.033	0.07 (0.03)	0.033
			Martha08	0.48	0.04 (0.04)	0.382						
			Martha10	0.47	0.12 (0.05)	0.030						
CLEC4M	rs868875	A/G	FVL	0.22	-0.07 (0.1)	0.515	0	0.762	-0.08 (0.03)	0.026	-0.08 (0.03)	0.026
			Martha08	0.32	-0.10 (0.05)	0.036						
			Martha10	0.35	-0.05 (0.06)	0.424						

* Common/rare alleles

[†] Allele frequency of the minor allele

⁽¹⁾ rs10866867 serves as proxy for rs2726953 ($r^2 = 0.92$); ⁽²⁾ rs216335 serves as proxy for rs216318 ($r^2 = 1$)

⁽³⁾ rs1063856 serves as proxy for Rs1063857 ($r^2 = 1$); ⁽⁴⁾ rs4334059 serves as proxy for rs7978987 ($r^2 = 1.0$)

⁽⁵⁾ rs2402074 serves as proxy for rs10133762 ($r^2 = 0.96$); No good proxy with $r^2 > 0.5$ was available for the VWF rs4764478

Our meta-analysis was also able to replicate several of the previously reported associations between SNPs and vWF/FVIII levels. Replicated associations include vWF-associated SNPs at *STXBP5*, *VWF*, *STX2*, *TC2N* and *CLEC4M* genes, and FVIII-associated SNPs within *SCARA5* and *VWF* genes. Other previously reported associations were not replicated, such as those involving *LDLR*, *BAI3*, and *STAB2* SNPs [5,9,16]. In addition to a lack of power, as previously discussed, this could be due to differential effects of SNP in normal range of plasma levels compared to the higher levels observed in VT patients. This could apply to the association of *BAI3* with vWF levels observed in healthy nuclear families [9] where plasma levels were lower than those observed in

our VT samples. Conversely, this explanation does not completely hold for the *LDLR* SNPs that were found associated with FVIII activity in a population [5] where FVIII activity in healthy individuals were at higher levels than those observed in our VT patients. Besides, in these two studies, different methods from those we have used here were employed to measure vWF and FVIII activity, and this could also contribute to the discrepancies observed in our study.

Conclusions

In conclusion, a carefully planned meta-analysis of three independent samples gathering 1,624 individuals genotyped for more than 400,000 SNPs all over the genome

Table 5 Association of Previously Identified SNPs with FVIII Activity in the three GWAS Datasets

Gene	SNP	Alleles*	MAF [†]	β (SE)	p	I ²	P _{het}	Random Effect		Fixed Effect		
								β (SE)	p	β (SE)	p	
STXBPS	rs9390459	G/A	FVL	0.43	0.15 (0.08)	0.083	0.65	0.059	-0.02 (0.06)	0.795	-0.04 (0.03)	0.310
			Martha08	0.42	-0.08 (0.06)	0.158						
			Martha10	0.43	-0.07 (0.05)	0.199						
SCARA5	rs9644133	C/T	FVL	0.24	-0.08 (0.1)	0.433	0	0.753	-0.12 (0.05)	0.009	-0.12 (0.05)	0.009
			Martha08	0.17	-0.16 (0.07)	0.029						
			Martha10	0.18	-0.10 (0.07)	0.152						
VWF	rs1063856	A/G	FVL	0.45	0.11 (0.08)	0.170	0	0.843	0.08 (0.03)	0.020	0.08 (0.03)	0.020
			Martha08	0.37	0.09 (0.06)	0.114						
			Martha10	0.38	0.06 (0.05)	0.249						
STAB2	rs4981021 ⁽¹⁾	G/A	FVL	0.27	-0.13 (0.09)	0.146	0	0.389	-0.02 (0.04)	0.521	-0.02 (0.04)	0.521
			Martha08	0.32	-0.02 (0.06)	0.737						
			Martha10	0.29	0.02 (0.06)	0.782						
LDLR	rs2228671	C/T	FVL	0.14	-0.03 (0.11)	0.816	0.46	0.157	-0.01 (0.07)	0.890	-0.01 (0.05)	0.894
			Martha08	0.11	0.11 (0.09)	0.193						
			Martha10	0.10	-0.13 (0.09)	0.161						
	rs688	C/T	FVL	0.38	-0.25 (0.09)	0.005	0.79	0.010	-0.05 (0.08)	0.531	-0.02 (0.03)	0.652
			Martha08	0.45	0.06 (0.05)	0.235						
Martha10	0.45	-0.007 (0.05)	0.901									

* Common/rare alleles

[†] Allele frequency of the minor allele

⁽¹⁾ rs4981021 serves as proxy for rs12229292 ($r^2 = 0.88$)

Table 6 In Silico Association With Venous Thrombosis of the Identified vWF- and FVIII Associated SNPs

	Alleles*	Minor Allele Frequency		Cochran Armitage P-value	
		Cases	Controls		
vWF associated SNPs					
VPS8	rs4686760	A/G	0.441	0.475	P = 0.101
EPB41L4A	rs13361927	G/A	0.065	0.062	P = 0.797
KRT18P24	rs1634352†	G/A	0.284	0.318	P = 0.055
12q21.33	rs1438933	G/A	0.256	0.294	P = 0.051
CDH2	rs2298574	A/G	0.084	0.093	P = 0.444
SAFB2	rs732505	G/A	0.061	0.064	P = 0.713
FVIII associated SNPs					
VAV2	rs12344583	A/G	0.217	0.193	P = 0.133
ACCN1	rs1354492	G/A	0.476	0.469	P = 0.740
ACCN1	rs12941510	G/A	0.310	0.350	P = 0.046

*Common/minor alleles

† serves as proxy for rs1757948 ($r^2 = 1$).

No good proxy with $r^2 > 0.80$ was available for rs6708166 (LBH), rs1321761 (FAM46A) and rs7306642 (STAB2)

Table 7 Genotype Distributions of rs4686760 and rs12941510 Across VT Samples

	rs4686760			MAF ⁽²⁾
	AA	AG	GG	
MARTHA08	271	502	198	0.462
MARTHA10	173	281	115	0.449
GWAS patients	129	196	81	0.441
All VT patients	573	979	394	0.454
GWAS controls	354	581	292	0.475
Test of association $P = 0.108^{(1)}$				
	rs12941510			MAF
	AA	AG	GG	
MARTHA08	93	409	469	0.306
MARTHA10	67	243	259	0.331
GWAS patients	45	161	199	0.310
All VT patients	205	813	927	0.314
GWAS controls	139	576	512	0.348
Test of association $P = 0.0056$				

⁽¹⁾ Cochran Armitage trend test

⁽²⁾ Minor Allele Frequency

replicated very recent findings but did not reveal any new genetic factors that could individually explain at least 2% of the plasma variability of vWF and FVIII levels.

Additional material

Additional file 1: FVL Family Questionnaire.

Additional file 2: MARTHA questionnaire. Excel file illustrating the questionnaire used for selecting MARTHA VT patients.

Additional file 3: Figure S1. Genotype filtering strategy applied to the three GWAS datasets. ⁽¹⁾ A genotype calling rate of > 0.90 was used in the FVL families and a threshold of 0.99 was used for the MARTHA patients. ⁽²⁾ SNPs with minor allele frequency less than 0.04 and 0.01 in FVL families and MARTHA patients, respectively, were excluded from the analysis. ⁽³⁾ SNPs demonstrating deviation from Hardy-Weinberg equilibrium at $p < 10^{-5}$ were excluded. 217 SNPs failed the genotype calling criterion simultaneously in the three study samples and this number was 19,111 for the minor allele frequency criterion. 19 SNPs failed the Hardy-Weinberg criterion in MARTHA08 and MARTHA10.

Additional file 4: Table S1. Haplotype Association Analysis of ACCN1 rs1354492 and rs12941510 With Plasma FVIII levels in MARTHA08 and MARTHA10 Studies. ⁽¹⁾ Haplotypic effect associated with each haplotype by comparison to the most frequent AG haplotype under the assumption of haplotype additive effects. Analyses were adjusted for age, sex and ABO blood group.

Acknowledgements

The French-Canadian FVL family study was supported by grants from the Canadian Institutes of Health Research (MOP86466) and by the Heart and Stroke Foundation of Canada (T6484). The MARTHA studies were supported by a grant from the Program Hospitalier de la Recherche Clinique. G.A hold an "INSERM Poste d'accueil" position and T.O.M was supported by a grant from the Fondation pour la Recherche Médicale. F.G and P.W. hold Canada

Research Chairs. A France-Canada Research Fund 2008 provided opportunities for face-to-face meetings of lead collaborators.

Author details

¹UMR_S 937, INSERM, Boulevard de l'Hopital, Paris, 75013, France. ²UMR_S 937, ICAN Institute, Université Pierre et Marie Curie, Boulevard de l'Hopital, 75013, Paris, France. ³Dalla Lana School of Public Health, University of Toronto, College Street, Toronto, M5T 3M7, Ontario, Canada. ⁴UMR_S 626, INSERM, rue Saint-Pierre, Marseille, 13385, France. ⁵UMR_S 626, Université de la Méditerranée, rue Saint-Pierre, Marseille, 13385 France. ⁶Department of Medicine, Ottawa Hospital Research Institute, Carling Avenue, Ottawa, K1Y 4E9, Ontario, Canada. ⁷Institut de Génomique, Centre National de Génotypage, Commissariat à l'Energie Atomique, rue Gaston Crémieux, Evry, 91057, France.

Authors' contributions

PEM, ML, FG and DAT designed the study and directed its implementation. GA, TOM and AD carried out statistical analyses. MG and WC were responsible for data collection and database management. GA drafted the article that was further reviewed by PEM, FG and DAT. All authors read and approved the final manuscript.

Competing interests

The authors declare that they have no competing interests.

Received: 7 May 2011 Accepted: 2 August 2011

Published: 2 August 2011

References

- Koster T, Blann AD, Briet E, Vandenbroucke JP, Rosendaal FR: **Role of clotting factor VIII in effect of von Willebrand factor on occurrence of deep-vein thrombosis.** *Lancet* 1995, **345**:152-155.
- Kraaijenhagen RA, in't Anker PS, Koopman MM, Reitsma PH, Prins MH, van den Ende A, *et al*: **High plasma concentration of factor VIIIc is a major risk factor for venous thromboembolism.** *Thromb Haemost* 2000, **83**:5-9.
- Tsai AW, Cushman M, Rosamond WD, Heckbert SR, Tracy RP, Aleksic N, *et al*: **Coagulation factors, inflammation markers, and venous thromboembolism: the longitudinal investigation of thromboembolism etiology (LITE).** *Am J Med* 2002, **113**:636-642.
- van Schie MC, de Maat MP, Isaacs A, van Duin CM, Deckers JW, Dippel DW, *et al*: **Variation in the von Willebrand Factor gene is associated with VWF levels and with the risk of cardiovascular disease.** *Blood* 2011, **117**:1393-1399.
- Martinelli N, Girelli D, Lunghi B, Pinotti M, Marchetti G, Malerba G, *et al*: **Polymorphisms at LDLR locus may be associated with coronary artery disease through modulation of coagulation factor VIII activity and independently from lipid profile.** *Blood* 2010, **116**:5688-5697.
- Whincup PH, Danesh J, Walker M, Lennon L, Thomson A, Appleby P, *et al*: **von Willebrand factor and coronary heart disease: prospective study and meta-analysis.** *Eur Heart J* 2002, **23**:1764-1770.
- Folsom AR, Rosamond WD, Shahar E, Cooper LS, Aleksic N, Nieto FJ, *et al*: **Prospective study of markers of hemostatic function with risk of ischemic stroke. The Atherosclerosis Risk in Communities (ARIC) Study Investigators.** *Circulation* 1999, **100**:736-742.
- Cambronero F, Vilchez JA, Garcia-Honrubia A, Ruiz-Espejo F, Moreno V, Hernandez-Romero D, *et al*: **Plasma levels of von Willebrand factor are increased in patients with hypertrophic cardiomyopathy.** *Thromb Res* 2010, **126**:e46-50.
- Antoni G, Morange PE, Luo Y, Saut N, Burgos G, Heath S, *et al*: **A multi-stage multi-design strategy provides strong evidence that the BA13 locus is associated with early-onset venous thromboembolism.** *J Thromb Haemost* 2010, **8**:2671-2679.
- Vormittag R, Bencur P, Ay C, Tengler T, Vukovich T, Quehenberger P, *et al*: **Low-density lipoprotein receptor-related protein 1 polymorphism 663 C > T affects clotting factor VIII activity and increases the risk of venous thromboembolism.** *J Thromb Haemost* 2007, **5**:497-4502.
- Carpeggiani C, Coceani M, Landi P, Michelassi C, L'Abbate A: **ABO blood group alleles: A risk factor for coronary artery disease. An angiographic study.** *Atherosclerosis* 2010, **211**:461-466.
- Teupser D, Baber R, Ceglarek U, Scholz M, Illig T, Gieger C, *et al*: **Genetic regulation of serum phytosterol levels and risk of coronary artery disease.** *Circ Cardiovasc Genet* 2010, **3**:331-339.

13. Barbalic M, Dupuis J, Dehghan A, Bis JC, Hoogeveen RC, Schnabel RB, *et al.*: **Large-scale genomic studies reveal central role of ABO in sP-selectin and sICAM-1 levels.** *Hum Mol Genet* 2010, **19**:1863-1872.
14. Souto JC, Almasy L, Borrell M, Gari M, Martinez E, *et al.*: **Genetic determinants of hemostasis phenotypes in Spanish families.** *Circulation* 2000, **101**:1546-1551.
15. Morange PE, Tregouet DA, Frere C, Saut N, Pellegrina L, Alessi MC, *et al.*: **Biological and genetic factors influencing plasma factor VIII levels in a healthy family population: results from the Stanislas cohort.** *Br J Haematol* 2005, **128**:91-99.
16. Smith NL, Chen M-H, Dehghan A, Strachan DP, Basu S, Soranzo N, *et al.*: **Novel associations of multiple genetic loci with plasma levels of Factor VII, Factor VIII and von Willebrand Factor. The CHARGE (Cohorts for Heart and Aging Research in Genome Epidemiology) Consortium.** *Circulation* 2010, **121**:1392-1392.
17. Abecasis GR, Cookson WO, Cardon LR: **The power to detect linkage disequilibrium with quantitative traits in selected samples.** *Am J Hum Genet* 2001, **68**:1463-1474.
18. Teslovich TM, Musunuru K, Smith AV, Edmondson AC, Stylianou IM, Koseki M, *et al.*: **Biological, clinical and population relevance of 95 loci for blood lipids.** *Nature* 2010, **466**:707-713.
19. Tregouet DA, Heath S, Saut N, Biron-Andreani C, Scheved JF, Pernod G, *et al.*: **Common susceptibility alleles are unlikely to contribute as strongly as the FV and ABO loci to VTE risk: results from a GWAS approach.** *Blood* 2009, **113**:5298-5303.
20. Hercberg S, Galan P, Preziosi P, Bertrais S, Mennen L, *et al.*: **The SU.VI.MAX Study: a randomized, placebo-controlled trial of the health effects of antioxidant vitamins and minerals.** *Arch Intern Med* 2004, **164**:2335-2342.
21. Sun L, Wilder K, McPeck MS: **Enhanced pedigree error detection.** *Hum Hered* 2002, **54**:99-110.
22. Purcell S, Neale B, Todd-Brown K, Thomas L, Ferreira MA, Bender D, *et al.*: **PLINK: a tool set for whole-genome association and population-based linkage analyses.** *Am J Hum Genet* 2007, **81**:559-575.
23. Price AL, Patterson NJ, Plenge RM, Weinblatt ME, Shadick NA, Reich D: **Principal components analysis corrects for stratification in genome-wide association studies.** *Nat Genet* 2006, **38**:904-909.
24. Peng B, Yu RK, Dehoff KL, Amos CI: **Normalizing a large number of quantitative traits using empirical normal quantile transformation.** *BMC Proc* 2007, **1**(Suppl 1):S156.
25. Tregouet DA, Garelle V: **A new JAVA interface implementation of THESIAS: testing haplotype effects in association studies.** *Bioinformatics* 2007, **23**:1038-1039.
26. Magi R, Morris AP: **GWAMA: software for genome-wide association meta-analysis.** *BMC Bioinformatics* 2010, **11**:288.
27. Morange PE, Saut N, Antoni G, Emmerich J, Tregouet DA: **Impact on venous thrombosis risk of newly discovered gene variants associated with FVIII and VWF plasma levels.** *J Thromb Haemost* 2011, **9**:229-231.
28. Gauderman WJ, Morrison JM: **Quanto 1.1: a computer program for power and sample size calculations for genetic-epidemiology studies.** 2006 [<http://hydra.usc.edu/gxe>].
29. Chen YJ, Stevens TH: **The VPS8 gene is required for localization and trafficking of the CPY sorting receptor in *Saccharomyces cerevisiae*.** *Eur J Cell Biol* 1996, **70**:289-297.
30. Agaphonov M, Romanova N, Sokolov S, Iline A, Kalebina T, *et al.*: **Defect of vacuolar protein sorting stimulates proteolytic processing of human urokinase-type plasminogen activator in the yeast *Hansenula polymorpha*.** *FEMS Yeast Res* 2005, **5**:1029-1035.
31. Briegel KJ, Baldwin HS, Epstein JA, Joyner AL: **Congenital heart disease reminiscent of partial trisomy 2p syndrome in mice transgenic for the transcription factor *Lbh*.** *Development* 2005, **132**:3305-3316.
32. Conen KL, Nishimori S, Provot S, Kronenberg HM: **The transcriptional cofactor *Lbh* regulates angiogenesis and endochondral bone formation during fetal bone development.** *Dev Biol* 2009, **333**:348-358.
33. Mori M, Murata Y, Kotani T, Kusakari S, Ohnishi H, Saito Y: **Promotion of cell spreading and migration by vascular endothelial-protein tyrosine phosphatase (VE-PTP) in cooperation with integrins.** *J Cell Physiol* 2010, **224**:195-204.
34. Vila-Carriles WH, Kovacs GG, Jovov B, Zhou ZH, Pahwa AK, Colby G, *et al.*: **Surface expression of ASIC2 inhibits the amiloride-sensitive current and migration of glioma cells.** *J Biol Chem* 2006, **281**:19220-19232.

Pre-publication history

The pre-publication history for this paper can be accessed here:
<http://www.biomedcentral.com/1471-2350/12/102/prepub>

doi:10.1186/1471-2350-12-102

Cite this article as: Antoni *et al.*: Combined analysis of three genome-wide association studies on vWF and FVIII plasma levels. *BMC Medical Genetics* 2011 **12**:102.

**Submit your next manuscript to BioMed Central
and take full advantage of:**

- Convenient online submission
- Thorough peer review
- No space constraints or color figure charges
- Immediate publication on acceptance
- Inclusion in PubMed, CAS, Scopus and Google Scholar
- Research which is freely available for redistribution

Submit your manuscript at
www.biomedcentral.com/submit

