

**High glucose increases susceptibility to
oxidative-stress-induced apoptosis and DNA damage in
K-562 cells.**

Jan Hruđa, Vladimír Sramek, Xavier Leverve

► **To cite this version:**

Jan Hruđa, Vladimír Sramek, Xavier Leverve. High glucose increases susceptibility to oxidative-stress-induced apoptosis and DNA damage in K-562 cells.. Biomedical papers of the Medical Faculty of the University Palacký, Olomouc, Czechoslovakia, 2010, 154 (4), pp.315-20. <inserm-00628502>

HAL Id: inserm-00628502

<http://www.hal.inserm.fr/inserm-00628502>

Submitted on 4 Oct 2011

HAL is a multi-disciplinary open access archive for the deposit and dissemination of scientific research documents, whether they are published or not. The documents may come from teaching and research institutions in France or abroad, or from public or private research centers.

L'archive ouverte pluridisciplinaire **HAL**, est destinée au dépôt et à la diffusion de documents scientifiques de niveau recherche, publiés ou non, émanant des établissements d'enseignement et de recherche français ou étrangers, des laboratoires publics ou privés.

HIGH GLUCOSE INCREASES SUSCEPTIBILITY TO OXIDATIVE-STRESS-INDUCED APOPTOSIS AND DNA DAMAGE IN K-562 CELLS

Jan Hruđa^{a,b*}, Vladimír Sramek^b, Xavier Leverve^a

^a INSERM U884 Laboratoire de Bioénergétique Fondamentale et Appliquée, Université J. Fourier, Grenoble, France

^b Department of Anaesthesia and Intensive Care, St. Anne's University Hospital, Masaryk University, Brno, Czech Republic
E-mail: jan.hruda@fnusa.cz

Received: August 17, 2010; Accepted: November 11, 2010

Key words: K-562 cells/Oxidative stress/Apoptosis/Energy substrates/Hyperglycemia/DNA damage

Aim. The study was carried out to evaluate the effect of several substrates on oxidative stress induced apoptosis and in K-562 cells.

Methods. Glucose at 5, 11 and 30 mM concentrations was tested, as well as 5 mM glutamine and 5 mM fructose. The cells were exposed to tert-butylhydroperoxide (tBH) and apoptotic cells were evaluated by flow cytometry with FITC-Annexin V and propidium iodide. The effect of glucose concentration on DNA damage was evaluated using hydrogen peroxide and electrophoretic "DNA comets" assay at 5 mM and 30 mM glucose concentrations.

Results. The exposure of cells to tBH resulted in increased number of apoptotic cells, and this effect was prevented by administration of an antioxidant - N-Acetyl cysteine. Rising concentrations of glucose added to the toxic effect of tBH; we also observed some toxic effect of fructose and no effect of glutamine. We found higher susceptibility to hydrogen peroxide induced DNA damage with 30 mM glucose concentration.

Conclusion. Hyperglycemia increases the cell's susceptibility to oxidative stress and it also amplifies oxidative DNA damage. Glutamine - when used as a sole energetic substrate - showed no protective effect against oxidative stress.

INTRODUCTION

In the past decade considerable evidence has been shown that there is a tight link between energy metabolism of the cell, oxidative stress, and apoptosis^{1,2}.

Glycolysis is a universal metabolic pathway providing energy by converting glucose into pyruvate. Under aerobic conditions pyruvate is oxidized to CO₂ and H₂O in mitochondria providing ATP by oxidative phosphorylation. Using O₂ as final acceptor of electrons makes the breakdown of glucose a lot more efficient than in the case of lack of oxygen when pyruvate is converted to lactate. At the same time, however, it generates oxidative stress.

Oxidative stress is a result of overproduction of reactive oxygen species (ROS) which are produced in aerobic organisms by electron transport chain of mitochondria. The ROS are highly reactive agents readily attacking macromolecules including nucleic acids. Under physiological conditions, the complex I of the electron transport chain seems to be the major site of ROS production^{3,4}.

Apoptosis is energy-demanding cellular death which is regulated in several ways⁵. Mitochondria play a key role in the signalling pathway resulting in apoptosis⁶. Once the cascade is activated by the cell death stimuli (e.g. TNF- α), caspases-8 and -10 are cleaved to their active forms and mitochondria release cytochrome C and Apoptosis Inducing Factor into the cytoplasm⁷. Released cytochrome C then activates the initiator caspase-9 which consequently activates the effector caspases-3 and -7.

These effector caspases are also directly activated by the initiator caspases-8 and -10. Prolonged oxidative stress as well as DNA damage are some of the triggers of apoptosis that have been identified.

We hypothesized, that the susceptibility of the cell to oxidative stress and apoptosis is affected by the metabolic state of the cells. For this reason the ability of the cell to withstand the toxic effect of an oxidizing agent may vary significantly when different substrates are metabolised as a source of energy. In clinical practise a similar situation occurs in intensive care when a patient with varying blood glucose control and energy substrate availability is exposed to oxidative insults (e.g. ischaemia/reperfusion, hyperoxaemia, etc.).

MATERIAL AND METHODS

Cell culture and experimental condition

The K-562 commercial cell line was used in this experiment. The cells were cultured in RPMI-1640 medium supplied with 10% bovine fetal serum and 1% L-glutamine-streptomycin. They were maintained in the exponential growth phase by regular centrifugation and resuspension in fresh RPMI medium.

To avoid alteration of results by the constituents of RPMI medium the cells were centrifuged and resuspended in experimental medium for the experiment. This was prepared individually based on Krebs' medium and con-

Table 1. The percentage of surviving cells at different glucose and tBH concentrations.

Glucose	tBH	Median	95% Confidence interval	Min - Max
5 Mm	0	93.7	91.6 - 94.0	89.6 - 94.2
	250 μ M	89.1	88.0 - 90.7	86.9 - 92.5
	500 μ M	68.6	66.7 - 69.8	65.9 - 71.0
11 mM	0	92.9	91.1 - 93.8	89.7 - 94.7
	250 μ M	90.0	87.3 - 90.8	85.4 - 91.3
	500 μ M	64.5	59.5 - 68.1	53.4 - 72.1
30 mM	0	87.6	85.5 - 88.6	83.7 - 90.2
	250 μ M	77.0	71.3 - 78.7	67.2 - 81.5
	500 μ M	55.8	39.2 - 60.1	25.9 - 62.3
Glucose concentration	p<0.01			
tBH concentration	p<0.01			
Glucose-tBH interaction	p<0.01			

Table 2. The percentage of surviving cells at different tBH concentrations with different substrates.

Substrate	tBH	Median	95% Confidence interval	Min - Max
Glucose 5 mM	0	93.7	91.6 - 94.0	89.6 - 94.2
	250 μ M	89.1	88.0 - 90.7	86.9 - 92.5
	500 μ M	68.6	66.7 - 69.8	65.9 - 71.0
Glutamin 5 mM	0	93.8	91.2 - 94.0	90.2 - 95.0
	250 μ M	85.4	78.2 - 88.3	77.6 - 88.8
	500 μ M	66.9	38.2 - 69.7	35.4 - 71.5
Fructose 5 mM	0	92.4	90.2 - 93.3	89.9 - 93.0
	250 μ M	89.0	80.5 - 91.1	78.9 - 92.0
	500 μ M	50.9	34.7 - 63.3	33.1 - 63.8
Glucose vs.	Glutamine	Fructose		
no tBH	p=0.260	p=0.441		
tBH 250 μ M	p=0.040	p=0.44		
tBH 500 μ M	p=0.374	p<0.01		

Table 3. "Tail moments" in DNA comets assay at two different glucose concentrations and exposure to hydrogen peroxide. The differences between all groups are statistically significant (p<0.01).

Glucose	Ox.stress	Median	95% Confidence interval	Min - Max
5 mM	-	0.022	0.20 - 0.61	0.001 - 0.720
	H ₂ O ₂	7.481	1.18 - 3.63	5.28 - 9.85
30 mM	-	3.223	0.83 - 2.57	1.85 - 5.28
	H ₂ O ₂	47.672	4.54 - 13.99	32.81 - 56.56

taining 24 mM NaHCO₃, 120 mM NaCl, 4.8 mM KCl, 1.2 mM KH₂PO₄, 1.2 mM MgSO₄, and 1.2 mM CaCl₂.

All chemicals were purchased from Sigma-Aldrich, except for FITC-Annexin purchased from Molecular Probes via Interchim dealer.

Flow cytometry experimental protocol

The experimental medium was supplied with either 5 or 11 or 30 mM glucose, or 5 mM fructose, or 5 mM glutamine. An 8 hrs incubation in these media was followed by 20 min exposure to 250 or 500 μ M tert-butylhydroperoxide (tBH) with a control group not exposed to




Fig. 1. Cytometric detection of apoptosis with Annexin-FITC (FL1-H) and propidium iodide (FL3-H) – cells exposed to different concentrations of tBH; glucose 11 mM. The principle of the staining is based on high affinity of Annexin to phosphatidylserine (PhS). In normal cells, PhS is present solely on the cytoplasmic surface of the cell membrane where it cannot be accessed by Annexin. During the process of apoptosis, PhS is translocated to the outer layer of the cell membrane and the cell will become positive in Annexin staining. Propidium iodide (PI) does not pass through intact cell membrane and this is why it can be used to distinguish cells with compromised membrane. A cell positive in Annexin but negative in PI staining (lower right quadrant) can therefore be considered as apoptotic.

tBH. The cells were then centrifuged, washed with fresh RPMI, centrifuged again and resuspended in fresh RPMI. They were incubated overnight for 12 hrs then flow cytometry was performed to determine apoptosis.

Double staining with FITC-Annexin and propidium iodide was used to detect apoptotic cells. For flow cytometry the cells were harvested by centrifugation and washed with cold phosphate-saline buffer. The cells were then re-centrifuged and resuspended in Annexin-binding buffer containing 10 mM HEPES, 140 mM NaCl and 2.5 mM CaCl_2 at 1×10^6 cells/ml. FITC-Annexin was added to the cell suspension and incubated for 15 min in the dark at room temperature. Propidium iodide was added and cytometry was conducted using 488 nm laser. The FSC, SSC, FL1 and FL3 data was collected; compensation for FL1/FL3 spectral overlap was performed before cytometry reading. The Becton Dickinson “FACScan” analyzer with “CellQuest” BD acquisition/analysis software was used in this study.

DNA comets experimental protocol

The experimental medium was supplied with either 5 or 30 mM glucose; the cells were incubated in these media for 16 hrs. After the incubation the cell suspension was mixed with low-melt agarose at 37 °C, spread on prepared agarose-covered glass slides and cooled to 4 °C. The slides were then exposed to 1 mM Hydrogen Peroxide (H_2O_2) for 5 min with a control group not exposed to H_2O_2 . Hydrogen Peroxide was then washed off by three-




Fig. 2. The protective effect of N-Acetyl cysteine – percentage of viable cells (* $p < 0.05$ vs. tBH 1000 μM) at 1, 2 and 4 mM N-Acetyl cysteine concentrations.

fold rinsing in PBS and the cells were lysed on the slides with a solution containing 2.5 M NaCl, 100 mM EDTA, 10 mM Tris-Base, 10% DMSO and 1% Triton X-100 detergent. Subsequently electrophoresis was conducted for 30min at 25V/300mA. To visualize DNA fragments, the slides were stained with propidium iodide in a dark chamber and observed at 488 nm excitation wavelength in a fluorescence microscope (Fig. 3). Recorded images were analysed with the CometScore software available as freeware from TriTek corporation at http://autocomet.com/products_cometscore.php. “Tail moment” was used




Fig. 3. DNA Comet assay. DNA damage is clearly visible as “comet tail” and can be evaluated by special software.




Fig. 4. The metabolic connection between glucose and glutathione: pentose phosphate shuttle - an alternate pathway of glucose breakdown - is a key source of the reduced form of NADP necessary to restore the reduced form of glutathione. High glucose concentrations lead to inhibition of the pentose phosphate shuttle together with increased glucose flux through the glycolytic pathway. Thus high glucose concentrations lead to increased susceptibility to oxidative stress. (GS-SG oxidized glutathione, GSH reduced glutathione).

to imply the extent of DNA damage, as it includes both length of the tail and percentage of DNA contained in the tail⁸. Tail moment was calculated automatically by the evaluation software.

Statistical Analysis

A two-way ANOVA was applied to determine the statistical significance of the effect of glucose and tBH concentrations on the percentage of surviving cells. The Mann-Whitney U test was used to compare the effect of different substrates and the results of the DNA comets assay. All statistical tests were performed at the 5% significance level. All the results were obtained using Statistica software, StatSoft, Inc. (2007). STATISTICA (data analysis software system), version 8.0. www.statsoft.com.

Due to the non-normal character of the data (0–100%) variance homogeneity was violated. A Box-Cox transformation was used to normalise the variance and data analysis was repeated. The results of the analysis of the transformed data were identical.

RESULTS

The results of flow cytometry measurements are presented as the percentage of viable cells (detected by flow cytometry using Annexin-FITC and PI staining) after the final incubation of the experimental protocol. There were nine separate incubations and measurements for each particular condition. We observed a clear toxic effect of tBH

(Fig. 1). N-Acetylcysteine, a potent antioxidant clearly prevented the effect of tBH (Fig. 2). There was no effect on the viability of the cells if N-Acetylcysteine in 1, 2 and 4 mM concentrations was used without tBH (data not shown). In this way we confirmed that cell death was oxidative-stress related in our experiment layout.

The set of experiments with 5, 11 and 30 mM glucose as energetic substrate showed statistically significant ($p < 0.01$) decrease in percentage of surviving cells with rising tBH concentration. At 0, 250 and 500 μM tBH concentrations statistically significant ($p < 0.01$) decrease in percentage of surviving cells was found with rising glucose concentration. At the same time we proved ($p = 0.007$) that high concentrations of glucose add to the cytotoxicity of tBH (Table 1).

We found no significant change if 5 mM glutamine was used, compared to 5 mM glucose. With 5 mM of fructose there was no significant difference in cell mortality at 0 and 250 μM tBH concentrations, but the toxicity of high tBH concentrations was even more expressed (Table 2).

The results of DNA comets assay are presented as "Tail moment". Four separate incubations for each condition were performed with two slides prepared from each incubation and 50 cells evaluated on each slide. We found significantly greater DNA damage ($p < 0.01$) in cells incubated with 30 mM glucose compared to 5 mM glucose in both hydrogen peroxide and control group (Table 3).

DISCUSSION

In this study tBH had cytotoxic, apoptosis-inducing and dose-dependent effect on K-562 cells. We proved that high glucose concentrations add to the effect of tBH. As it was prevented by N-Acetyl cysteine – an antioxidant – it is clearly related to oxidative stress. The explanation for this may be the metabolic linkage between glucose breakdown and antioxidant protection of the cell (Fig. 4). Since tBH oxidizes the -SH groups it attenuates the cell's ability to protect against oxidative stress rather than directly inducing it. The -SH group is a key structure of action in glutathione – one of the most important mechanisms of cellular antioxidant defense. Reduced glutathione is essential for the breakdown of hydrogen peroxide via glutathione peroxidase. Once oxidized in this reaction, glutathione forms dimers and can only be restored with the reduced form of NADP. The source of reduced NADP within the cell is the pentose phosphate shuttle, an alternate metabolic pathway of glucose sharing its first step with glycolysis⁹. The next step of the pentose phosphate shuttle is catalysed by specific enzyme glucose-6-phospho-dehydrogenase which is notably inhibited by high glucose concentrations¹⁰. This is why high concentrations of glucose lead to increased glycolysis, mitochondrial metabolic activity and subsequently ROS production but at the same time they block production of NADP, thus attenuating the antioxidant protection of the cell. We suggest that in this way, high glucose concentration increases the toxic effect of an oxidizing agent. This finding is in agreement with other authors' observations that high rate of sub-

strate influx to mitochondria induces cell sensitivity to oxidant-induced apoptosis¹¹. The DNA damage observed in cells at high glucose concentrations can be explained in the same way as the additive effect of glucose to tBH – increased ROS production together with insufficient restoration of glutathione¹².

Fructose enters the glycolytic pathway down the flow at the level of triose-phosphates and therefore bypasses the first step of glycolysis – its most important regulatory mechanism¹³. As a result, the breakdown of fructose proceeds in a fast and rather uncontrolled manner and increases mitochondrial ROS production. This is probably not substantial under otherwise stable conditions but may become critical when the cell has to challenge further oxidative insults. This is a possible explanation why fructose proved to be toxic to cells only at high levels of tBH.

Glutamine has been reported to have certain anti-oxidant and anti-apoptotic capabilities^{14,15}. In our experiment, however, we found no protective effect of glutamine. The reason for this might be that glutamine was used here as the only available substrate for the cells and not just added as extra treatment.

The concentration of glucose of 30 mM is quite high, and may already have considerable osmotic effect. To ensure that this effect did not falsify our results, we conducted a limited series of experiments with media containing 5 mM D-glucose and 25 mM L-glucose which is not metabolized by the cells¹⁶. The results were similar to those obtained with 5 mM D-glucose only (data not shown).

CONCLUSION

We conclude, that higher glucose concentrations in culture media increase the demands of the antioxidative mechanisms and thereby make the cell more susceptible to oxidative-stress-induced apoptosis. The same effect also results in amplification of oxidative-stress-induced DNA damage in cells cultivated in high glucose media.

ACKNOWLEDGEMENT

The authors thank to Cecile Cottet and Christiane Chauvin from INSERM U844 for their willing help with flow cytometry and cell culture.

The authors claim no conflict of interests. The study was funded by INSERM; all the equipment used came from INSERM U884 laboratory.

REFERENCES

1. Ježek P, Hlavata L. Mitochondria in homeostasis of reactive oxygen species in cells, tissues and organism. *The International Journal of Biochemistry & Cell Biology* 2005;37:2478–2503.
2. Leverve XM. Mitochondrial function and substrate availability. *Crit Care Med* 2007;35(9 Suppl):S454–60
3. Leverve XM, Fontaine E. Role of substrates in the regulation of mitochondrial function in situ. *IUBMB Life* 2001;52(3–5):221–9

4. Magder S. Reactive oxygen species: toxic molecules or spark of life? *Critical Care* 2006;10:208
5. Steven HW. To die or not to die: an overview of apoptosis and its role in disease. *JAMA* 1998;279:300-307
6. Cavalli LR, Liang BC. Mutagenesis, tumorigenicity and apoptosis: are mitochondria involved? *Mutation Research* 1998;398:19-26
7. Reed JC, Green DR. Remodeling for Demolition: Changes in Mitochondrial Ultrastructure during Apoptosis. *Molecular Cell* 2002;9:1-9
8. Heaton PR, Ransley R, Charlton CJ, Mann SJ, Stevenson J, Smith BH, Rawlings JM, Harper EJ. Application of single-cell gel electrophoresis (comet) assay for assessing levels of DNA damage in canine and feline leukocytes. *J Nutr* 2002;132(6 Suppl 2):598S-603S.
9. Wamelink MM, Struys EA, Jakobs C. The biochemistry, metabolism and inherited defects of the pentose phosphate pathway: a review. *J Inherit Metab Dis* 2008;31(6):703-17.
10. Leverve X. Hyperglycemia and oxidative stress: complex relationships with attractive prospects. *Intensive Care Med* 2003;29(4):511-4.
11. Jeong D, Kim TS, Cho IT, Kim IY. Modification of glycolysis affects cell sensitivity to apoptosis induced by oxidative stress and mediated by mitochondria. *Biochemical and Biophysical Research Communications*. 2004;313:984-991
12. Hollins DL, Suliman HB, Piantadosi CA, Carraway MS. Glutathione regulates susceptibility to oxidant-induced mitochondrial DNA damage in human lymphocytes. *Free Radical Biology & Medicine* 2006;40:1220-1226
13. Mayes PA. Intermediary metabolism of fructose. *Am J Clin Nutr* 1993;58(5 Suppl):754S-765S.
14. Chang WK, Yang KD, Chuang H, Jan JT, Shaio MF. Glutamine protects activated human T cells from apoptosis by up-regulating glutathione and Bcl-2 levels. *Clin Immunol* 2002;104(2):151-60.
15. Petronini PG, Urbani S, Alfieri R, Borghetti AF, Guidotti GG. Cell susceptibility to apoptosis by glutamine deprivation and rescue: survival and apoptotic death in cultured lymphoma-leukemia cell lines. *J Cell Physiol* 1996;169(1):175-85.
16. Bühler I, Walter R, Reinhart WH. Influence of D- and L-glucose on erythrocytes and blood viscosity. *Eur J Clin Invest* 2001;31(1):79-85.