



A subset of dendritic cells as a major and constitutive source of IL-22BP

Jérôme Martin, Gaëlle Bériou, Michèle Heslan, Céline Bossard, Anne-Chantal Knol, Brigitte Dréno, Simon Milling, Miriam Merad, Régis Josien

► To cite this version:

Jérôme Martin, Gaëlle Bériou, Michèle Heslan, Céline Bossard, Anne-Chantal Knol, et al.. A subset of dendritic cells as a major and constitutive source of IL-22BP. 6th european workshop on immune-mediated inflammatory diseases, Nice, France. pp.P2. inserm-00643975

HAL Id: inserm-00643975

<https://www.hal.inserm.fr/inserm-00643975>

Submitted on 23 Nov 2011

HAL is a multi-disciplinary open access archive for the deposit and dissemination of scientific research documents, whether they are published or not. The documents may come from teaching and research institutions in France or abroad, or from public or private research centers.

L'archive ouverte pluridisciplinaire **HAL**, est destinée au dépôt et à la diffusion de documents scientifiques de niveau recherche, publiés ou non, émanant des établissements d'enseignement et de recherche français ou étrangers, des laboratoires publics ou privés.



POSTER PRESENTATION

Open Access

A subset of dendritic cells as a major and constitutive source of IL-22BP

Jérôme Martin^{1,2}, Gaëlle Bériou¹, Michèle Heslan¹, Céline Bossard³, Anne-Chantal Knol⁴, Brigitte Dréno⁴, Simon Milling⁵, Miriam Merad⁶, Régis Josien^{1,2*}

From 6th European Workshop on Immune-Mediated Inflammatory Diseases
Nice, France. 23-25 November 2011

Introduction

IL-22 is a cytokine produced by T cells and innate lymphocytes. Its receptor is exclusively expressed by non hematopoietic cells mostly epithelial cells and hepatocytes. IL-22 has pathogenic or protective effects depending on the model. A role for IL-22 is suggested in IMID such as psoriasis, rheumatoid arthritis and Crohn disease. IL-22BP is a soluble inhibitory receptor specific for IL-22 whose cellular source and physiological role are mainly unknown.

Aims

To identify the cellular source of IL-22BP in vivo and its regulation of expression.

Methods

Rat dendritic cells (DCs) were prepared from spleen. Human DCs were derived from monocytes (MDDC) in the presence of GM-CSF+IL-4. IL-22BP mRNA expression was assessed by q-PCR. Immunostaining experiments were performed with polyclonal and/or monoclonal Ab to IL-22BP. IL-22BP in serum was identified by WB.

Results

IL-22BP mRNA was expressed at high levels in rat secondary lymphoid organs and intestine. In spleen, IL-22BP mRNA was restricted to a CD4⁺SIRPA⁺ subset of DCs. This expression was constitutive. Similar results were obtained in lymph nodes. IL-22BP expression in DCs was confirmed at the protein level. An even stronger expression was found in a subset of lymph DCs migrating from gut. In mouse, IL-22BP expression appeared restricted to the CD103⁺ subsets of intestine DCs. In human, immunostaining experiments identified stellate IL-22BP⁺ cells in

colon lamina propria and dermis. In vitro, IL-22BP expression was induced during MDDC differentiation and retinoic acid receptor alpha agonist dramatically enhanced this expression. IL-22BP expression was rapidly down-regulated following maturation both in rat and human DC. Finally, WB analysis revealed high levels of serum IL-22BP in healthy volunteers.

Conclusion

These results indicate that tissue and lymphoid DCs are a main source of IL-22BP suggesting a role for these DCs in regulating IL-22 at epithelial barriers.

Author details

¹INSERM U643, ITUN, Nantes, France. ²Laboratoire d'Immunologie, Nantes, France. ³Service d'Anatomo-Pathologie, Nantes, France. ⁴INSERM U892, Laboratoire d'Immuno-Dermatologie, CHU Nantes, Nantes, France. ⁵University of Glasgow, Glasgow, UK. ⁶Mount Sinai School of Medicine, New York, USA.

Published: 23 November 2011

doi:10.1186/1479-5876-9-S2-P2

Cite this article as: Martin et al.: A subset of dendritic cells as a major and constitutive source of IL-22BP. *Journal of Translational Medicine* 2011 9(Suppl 2):P2.

¹INSERM U643, ITUN, Nantes, France

Full list of author information is available at the end of the article