

**Antibiotics induced commensal flora disruption favours
escherichia coli AIEC colonization and mesenteric
translocation in NOD2 knock-out mice**

Maryline Drouet, Cécile Vignal, Elisabeth Singer, Luc Dubreuil, Pierre
Desreumaux, Christel Neut

► **To cite this version:**

Maryline Drouet, Cécile Vignal, Elisabeth Singer, Luc Dubreuil, Pierre Desreumaux, et al.. Antibiotics induced commensal flora disruption favours escherichia coli AIEC colonization and mesenteric translocation in NOD2 knock-out mice. 6th european workshop on immune-mediated inflammatory diseases, Nice, France. pp.O8. inserm-00643978

HAL Id: inserm-00643978

<https://www.hal.inserm.fr/inserm-00643978>

Submitted on 23 Nov 2011

HAL is a multi-disciplinary open access archive for the deposit and dissemination of scientific research documents, whether they are published or not. The documents may come from teaching and research institutions in France or abroad, or from public or private research centers.

L'archive ouverte pluridisciplinaire **HAL**, est destinée au dépôt et à la diffusion de documents scientifiques de niveau recherche, publiés ou non, émanant des établissements d'enseignement et de recherche français ou étrangers, des laboratoires publics ou privés.



ORAL PRESENTATION

Open Access

Antibiotics induced commensal flora disruption favours *escherichia coli* AIEC colonization and mesenteric translocation in NOD2 knock-out mice

Maryline Drouet^{1,2*}, Cécile Vignal¹, Elisabeth Singer^{1,2}, Luc Dubreuil^{1,2}, Pierre Desreumaux^{1,3}, Christel Neut^{1,2}

From 6th European Workshop on Immune-Mediated Inflammatory Diseases
Nice, France. 23-25 November 2011

Background

Ileal lesions of Crohn's disease (CD) patients are colonized by pathogenic adherent-invasive *Escherichia coli* (AIEC). NOD2 gene mutations are risk factors for ileal CD and are associated with an abnormal AIEC-induced immune response.

Aim

Since innate immune system and commensal microbiota are critical to maintain intestinal barrier integrity, we evaluated the impact of commensal microbiota disruption induced by short term antibiotic treatment on AIEC colonization and translocation in wild type (WT) and NOD2 knock-out mice (NOD2KO).

Methods

Disruption of commensal microbiota was induced by a 3 days antibiotic treatment orally administered in WT and NOD2KO mice. At day 3, mice were infected with 10^9 CFU AIEC once a day for 2 days. Animals were sacrificed at day 1, 5, 30 and 60 after AIEC administration. Ileum, colon and mesenteric lymph nodes (MLN) were sampled for AIEC quantification in ileal and colonic tissues, bacterial translocation in MLN, and evaluation of histological abnormalities and intestinal inflammation.

Results

Without antibiotic treatment, AIEC was not able to colonize WT and NOD2KO mice. Compared to control animals, commensal microbiota disruption induced by antibiotics led to a significant increase of ileal and colonic adherent AIEC in both WT and NOD2KO mice at day 1.

Persistent AIEC colonization was still observed until day 30 in the ileum but not in the colon of NOD2KO mice, disappearing at day 60. Mesenteric translocation of AIEC was observed until day 30 only in NOD2KO mice. No histologic ileal and colonic abnormalities and/or inflammation were observed in WT and NOD2KO mice treated with antibiotics and infected with AIEC.

Conclusion

Sustained AIEC colonization in ileal tissues of mice was induced by commensal flora disruption promoted by short term antibiotic treatment, independently of NOD2 expression. In contrast, commensal flora disruption by antibiotics induced a long term AIEC bacterial translocation in a NOD2 dependent manner.

Author details

¹INSERM U995, Lille, France. ²Faculty of Pharmacy, Lille 2 University, Lille, France. ³Dept. of Gastroenterology, Claude Huriez Hospital, Lille, France.

Published: 23 November 2011

doi:10.1186/1479-5876-9-S2-O8

Cite this article as: Drouet et al.: Antibiotics induced commensal flora disruption favours *escherichia coli* AIEC colonization and mesenteric translocation in NOD2 knock-out mice. *Journal of Translational Medicine* 2011 **9**(Suppl 2):O8.

¹INSERM U995, Lille, France

Full list of author information is available at the end of the article