



Immunosuppressive role of fibrinogen-like protein 2 (FGL2) in CD8+regulatory T cells-mediated long-term graft survival

Séverine Bézie, Séverine Ménoret, Laurent Tesson, Xian-Liang Li, Claire Usal,
Ignacio Anegon, Lise Caron

► To cite this version:

Séverine Bézie, Séverine Ménoret, Laurent Tesson, Xian-Liang Li, Claire Usal, et al.. Immunosuppressive role of fibrinogen-like protein 2 (FGL2) in CD8+regulatory T cells-mediated long-term graft survival. 6th european workshop on immune-mediated inflammatory diseases, Nice, France. pp.O5. inserm-00643979

HAL Id: inserm-00643979

<https://inserm.hal.science/inserm-00643979>

Submitted on 23 Nov 2011

HAL is a multi-disciplinary open access archive for the deposit and dissemination of scientific research documents, whether they are published or not. The documents may come from teaching and research institutions in France or abroad, or from public or private research centers.

L'archive ouverte pluridisciplinaire **HAL**, est destinée au dépôt et à la diffusion de documents scientifiques de niveau recherche, publiés ou non, émanant des établissements d'enseignement et de recherche français ou étrangers, des laboratoires publics ou privés.



ORAL PRESENTATION

Open Access

Immunosuppressive role of fibrinogen-like protein 2 (FGL2) in CD8⁺regulatory T cells-mediated long-term graft survival

Séverine Bézie^{1*}, Séverine Ménoret², Laurent Tesson², Xian-Liang Li¹, Claire Usal¹, Ignacio Anegon¹, Lise Caron¹

From 6th European Workshop on Immune-Mediated Inflammatory Diseases
Nice, France. 23-25 November 2011

Background

We have previously reported that, in a model of cardiac allograft in rat, blockade of CD40-CD40L interaction induces long-term graft survival mediated by CD8⁺ CD45RC^{low} regulatory T cells (Tregs) [1]. Transcriptomic comparison of Tregs from AdCD40Ig-treated vs naïve rats highlighted the overexpression of FGL2 whose immunoregulatory properties are little known [2].

Material and methods

A lewis 1W rat heart is grafted in a MHC-mismatched Lewis 1A rat and infected with 2.10¹⁰ pi of adenovirus recombinant for CD40Ig molecule (AdCD40Ig) the day of the graft. Tregs, effector CD4⁺CD25⁻ T lymphocytes (TL), and plasmacytoid dendritic cells (pDC) from spleen are sorted by FACS Aria for *in vitro* tests. For *in vivo* studies, 4.5.10¹¹vg of FGL2-recombinant adenovirus associated virus (AAVFGl2) are intramuscularly or intravenously injected in recipients 30 days before the graft. Splenocytes are transferred to sublethally irradiated rats by *i.v* injection the day before the graft.

Results

We confirmed FGL2-overexpression in splenic Tregs and in the graft of AdCD40Ig-treated vs non-treated and naïve rats, at mRNA and protein level. FGL2 involvement in Tregs immunosuppressive function was proved by *in vitro* and *in vivo* experiments. Indeed, Tregs from AdCD40Ig-treated rats inhibit TL proliferation in response to allogeneic pDC. This inhibition is abrogated by FGL2-blocking antibodies [3] and can be mimicked by FGL2 protein alone. Moreover, AAV-mediated FGL2

overexpression in rat prolongs graft survival with a median of 18.5 days vs 11 days for controls by *i.m* injection and survival is improved when *i.v* injected. Furthermore, adoptive transfer of splenocytes from an AAVFGl2-treated tolerant rat, to irradiated rats, transmits long-term graft survival iteratively.

Conclusions

This is the first demonstration that the immunosuppressive molecule FGL2 is able to induce a long-term graft survival and that this tolerance is active and transferable by splenocytes. Work is under progress to identify the population responsible for this infectious tolerance.

Author details

¹INSERM UMR 643, Nantes, France. ²Plateform Transgenic Rats Nantes IBiSA-CNRS, Nantes, France.

Published: 23 November 2011

References

- Guillonneau C, Hill M, Hubert FX, Chiffolleau E, Hervé C, Li XL, et al: CD40Ig treatment results in allograft acceptance mediated by CD8CD45RC T cells, IFN-gamma, and indoleamine 2,3-dioxygenase. *J Clin Invest* 2007, **117**:1096-1106.
- Liu H, Shalev I: The FGL2-FcgammaRIIB pathway: a novel mechanism leading to immunosuppression. *Eur J Immunol* 2008, **38**(Suppl 11):3114-3126.
- Li XL, Menoret S: Mechanism and localization of CD8 regulatory T cells in a heart transplant model of tolerance. *J Immunol* 2010, **185**(Suppl 2):823-833.

doi:10.1186/1479-5876-9-S2-O5

Cite this article as: Bézie et al.: Immunosuppressive role of fibrinogen-like protein 2 (FGL2) in CD8⁺regulatory T cells-mediated long-term graft survival. *Journal of Translational Medicine* 2011 **9**(Suppl 2):O5.

¹INSERM UMR 643, Nantes, France

Full list of author information is available at the end of the article