

Purging the HIV-1 reservoir through the disruption of the PD-1 pathway

Sandrine Dafonseca, Nicolas Chomont, Mohamed El Far, Mohamed-Rachid Boulassel, Jean-Pierre Routy, Rafick Sékaly

► **To cite this version:**

Sandrine Dafonseca, Nicolas Chomont, Mohamed El Far, Mohamed-Rachid Boulassel, Jean-Pierre Routy, et al.. Purging the HIV-1 reservoir through the disruption of the PD-1 pathway. International AIDS Society's Workshop "Towards a Cure": HIV Reservoirs and Strategies to Control Them, Jul 2010, Vienne, Austria. pp.O15, 10.1186/1758-2652-13-S3-O15 . inserm-00663876

HAL Id: inserm-00663876

<https://www.hal.inserm.fr/inserm-00663876>

Submitted on 27 Jan 2012

HAL is a multi-disciplinary open access archive for the deposit and dissemination of scientific research documents, whether they are published or not. The documents may come from teaching and research institutions in France or abroad, or from public or private research centers.

L'archive ouverte pluridisciplinaire **HAL**, est destinée au dépôt et à la diffusion de documents scientifiques de niveau recherche, publiés ou non, émanant des établissements d'enseignement et de recherche français ou étrangers, des laboratoires publics ou privés.

ORAL PRESENTATION

Open Access

Purging the HIV-1 reservoir through the disruption of the PD-1 pathway

S DaFonseca^{1*}, N Chomont¹, M El Far^{2,3,4}, R Boulassel⁵, J Routy^{5,6}, R Sékaly^{1,2,3,4,6}

From International AIDS Society's Workshop "Towards a Cure": HIV Reservoirs and Strategies to Control Them Vienna, Austria. 16-17 July 2010

Background

The main obstacle to HIV-1 eradication is a small pool of T_{CM} (central memory) and T_{TM} (transitional memory) latently infected CD4⁺ T cells that persist in patients receiving HAART. The mechanisms implicated in the establishment and persistence of the HIV reservoir are still unknown. The PD-1 receptor is expressed by CD4⁺ T cells from HIV-infected patients, and efficiently inhibits T cell proliferation. Here, we investigate a possible role for this receptor in the establishment and maintenance of a cellular reservoir for HIV.

Methods

PBMCs were obtained by leukapheresis from HAART-naïve, chronically HIV-1-infected subjects and were subjected to total CD4⁺ T cells negative selection. Viral production was induced through TCR triggering (CD3/CD28) with or without co-triggering of the PD-1 pathway with an Ig-PD-L1 chimera. Cell culture supernatants were serially harvested; viral release was quantified by QRT-PCR or p24 ELISA. The frequency of CD4⁺ T cells harbouring HIV DNA was determined by Q-PCR.

Results

Cell sorting and Q-PCR experiments showed that PD-1^{high} cells from viremic donors preferentially harbour HIV-1 integrated DNA when compared with their PD-1^{low} counterparts, indicating that these cells constitute a preferential reservoir for the virus. Triggering of the PD-1 pathway inhibits 50% of HIV-1 production in primary CD4⁺ T cells at day 1 and up to 95% at day 3. Importantly, this inhibition was restricted to PD-1^{high} cells, demonstrating the specificity of this mechanism. Moreover, we observed

that the disruption of the PD-1/PD-L1 interaction enhances the spontaneous release of HIV-1 virions by CD4⁺ T cells.

Conclusions

Our results suggest that: (1) the PD-1 receptor can be used as a specific marker to target the HIV-1 reservoir; (2) viral production is inhibited after triggering of the PD-1 receptor; and (3) viral production was enhanced after blocking of this inhibitory pathway. Altogether, our results demonstrate a crucial role for PD-1 in the establishment.

Author details

¹Vaccine and Gene Therapy Institute, 11350 SW Village Parkway, Port St. Lucie, Florida 34987, USA. ²Laboratoire d'Immunologie, Centre de Recherche du Centre Hospitalier de l'Université de Montréal (CR-CHUM) Saint-Luc, 264 René Lévesque Est, Montréal, Québec H2X1P1, Canada. ³Laboratoire d'Immunologie, Département de Microbiologie et d'Immunologie, Université de Montréal, Québec, Canada. ⁴INSERM U743, CR-CHUM, Université de Montréal, 264 René Lévesque Est, Montréal, Québec H2X1P1, Canada. ⁵Immunodeficiency Service and Division of Hematology, Royal Victoria Hospital, McGill University Health Centre (MUHC), McGill University, 687 Pine Avenue West, Montréal, Québec H3A 1A1, Canada. ⁶Department of Microbiology and Immunology, 3775 University Street, McGill University, Montréal, Québec H3A2B4, Canada.

Published: 4 November 2010

doi:10.1186/1758-2652-13-S3-O15

Cite this article as: DaFonseca *et al.*: Purging the HIV-1 reservoir through the disruption of the PD-1 pathway. *Journal of the International AIDS Society* 2010 **13**(Suppl 3):O15.

* Correspondence: sdafonseca@vgti-fl.org

¹Vaccine and Gene Therapy Institute, 11350 SW Village Parkway, Port St. Lucie, Florida 34987, USA

Full list of author information is available at the end of the article