

Common breast cancer susceptibility alleles are associated with tumour subtypes in BRCA1 and BRCA2 mutation carriers: results from the Consortium of Investigators of Modifiers of BRCA1/2.

Anna Mulligan, Fergus Couch, Daniel Barrowdale, Susan Domchek, Diana Eccles, Heli Nevanlinna, Susan Ramus, Mark Robson, Mark Sherman, Amanda Spurdle, et al.

► **To cite this version:**

Anna Mulligan, Fergus Couch, Daniel Barrowdale, Susan Domchek, Diana Eccles, et al.. Common breast cancer susceptibility alleles are associated with tumour subtypes in BRCA1 and BRCA2 mutation carriers: results from the Consortium of Investigators of Modifiers of BRCA1/2.. Breast Cancer Research, BioMed Central, 2011, 13 (6), pp.R110. 10.1186/bcr3052 . inserm-00670601

HAL Id: inserm-00670601

<https://www.hal.inserm.fr/inserm-00670601>

Submitted on 15 Feb 2012

HAL is a multi-disciplinary open access archive for the deposit and dissemination of scientific research documents, whether they are published or not. The documents may come from teaching and research institutions in France or abroad, or from public or private research centers.

L'archive ouverte pluridisciplinaire **HAL**, est destinée au dépôt et à la diffusion de documents scientifiques de niveau recherche, publiés ou non, émanant des établissements d'enseignement et de recherche français ou étrangers, des laboratoires publics ou privés.

RESEARCH ARTICLE

Open Access

Common breast cancer susceptibility alleles are associated with tumour subtypes in *BRCA1* and *BRCA2* mutation carriers: results from the Consortium of Investigators of Modifiers of *BRCA1/2*

Anna Marie Mulligan¹, Fergus J Couch², Daniel Barrowdale³, Susan M Domchek⁴, Diana Eccles⁵, Heli Nevanlinna⁶, Susan J Ramus⁷, Mark Robson⁸, Mark Sherman⁹, Amanda B Spurdle¹⁰, Barbara Wappenschmidt¹¹, Andrew Lee³, Lesley McGuffog³, Sue Healey¹⁰, Olga M Sinilnikova^{12,13}, Ramunas Janavicius¹⁴, Thomas vO Hansen¹⁵, Finn C Nielsen¹⁵, Bent Ejlersen¹⁶, Ana Osorio¹⁷, Iván Muñoz-Repeto¹⁷, Mercedes Durán¹⁸, Javier Godino¹⁹, Maroulio Pertesi²⁰, Javier Benítez²¹, Paolo Peterlongo²², Siranoush Manoukian²³, Bernard Peissel²³, Daniela Zaffaroni²³, Elisa Cattaneo²³, Bernardo Bonanni²⁴, Alessandra Viel²⁵, Barbara Pasini²⁶, Laura Papi²⁷, Laura Ottini²⁸, Antonella Savarese²⁹, Loris Bernard³⁰, Paolo Radice²², Ute Hamann³¹, Martijn Verheus³², Hanne EJ Meijers-Heijboer³³, Juul Wijnen³⁴, Encarna B Gómez García³⁵, Marcel R Nelen³⁶, C Marleen Kets³⁷, Caroline Seynaeve³⁸, Madeleine MA Tilanus-Linthorst³⁹, Rob B van der Luijt⁴⁰, Theo van Os⁴¹, Matti Rookus³², Debra Frost³, J Louise Jones⁴², D Gareth Evans⁴³, Fiona Lalloo⁴³, Ros Eeles⁴⁴, Louise Izatt⁴⁵, Julian Adlard⁴⁶, Rosemarie Davidson⁴⁷, Jackie Cook⁴⁸, Alan Donaldson⁴⁹, Huw Dorkins⁵⁰, Helen Gregory⁵¹, Jacqueline Eason⁵², Catherine Houghton⁵³, Julian Barwell⁵⁴, Lucy E Side⁵⁵, Emma McCann⁵⁶, Alex Murray⁵⁷, Susan Peock³, Andrew K Godwin⁵⁸, Rita K Schmutzler¹¹, Kerstin Rhiem¹¹, Christoph Engel⁵⁹, Alfons Meindl⁶⁰, Ina Ruehl⁶¹, Norbert Arnold⁶², Dieter Niederacher⁶³, Christian Sutter⁶⁴, Helmut Deissler⁶⁵, Dorothea Gadzicki⁶⁶, Karin Kast⁶⁷, Sabine Preisler-Adams⁶⁸, Raymonda Varon-Mateeva⁶⁹, Ines Schoenbuchner⁷⁰, Britta Fiebig⁷¹, Wolfram Heinritz⁷², Dieter Schäfer⁷³, Heidrun Gevensleben⁷⁴, Virginie Caux-Moncoutier⁷⁵, Marion Fassy-Colcombet⁷⁵, François Cornelis⁷⁶, Sylvie Mazoyer¹³, Mélanie Léoné¹², Nadia Boutry-Kryza¹², Agnès Hardouin⁷⁷, Pascaline Berthet⁷⁷, Danièle Muller⁷⁸, Jean-Pierre Fricker⁷⁸, Isabelle Mortemousque⁷⁹, Pascal Pujol⁸⁰, Isabelle Coupier⁸⁰, Marine Lebrun⁸¹, Caroline Kientz⁸¹, Michel Longy⁸², Nicolas Sevenet⁸², Dominique Stoppa-Lyonnet^{75,83}, Claudine Isaacs⁸⁴, Trinidad Caldes⁸⁵, Miguel de la Hoya⁸⁵, Tuomas Heikkinen⁶, Kristiina Aittomäki⁸⁶, Ignacio Blanco⁸⁷, Conxi Lazaro⁸⁷, Rosa B Barkardottir⁸⁸, Penny Soucy⁸⁹, Martine Dumont⁸⁹, Jacques Simard⁹⁰, Marco Montagna⁹¹, Silvia Tognazzo⁹¹, Emma D'Andrea⁹², Stephen Fox⁹³, Max Yan⁹⁴, Tim Rebbeck⁹⁵, Olufunmilayo I Olopade⁹⁶, Jeffrey N Weitzel⁹⁷, Henry T Lynch⁹⁸, Patricia A Ganz⁹⁹, Gail E Tomlinson¹⁰⁰, Xianshu Wang², Zachary Fredericksen¹⁰¹, Vernon S Pankratz¹⁰¹, Noralane M Lindor¹⁰², Csilla Szabo¹⁰³, Kenneth Offit⁸, Rita Sakr⁸, Mia Gaudet¹⁰⁴, Jasmine Bhatia⁸, Noah Kauff⁸, Christian F Singer¹⁰⁵, Muy-Kheng Tea¹⁰⁵, Daphne Gschwantler-Kaulich¹⁰⁵, Anneliese Fink-Retter¹⁰⁵, Phuong L Mai¹⁰⁶, Mark H Greene¹⁰⁶, Evgeny Imyanitov¹⁰⁷, Frances P O'Malley¹⁰⁸, Hilmi Ozcelik¹⁰⁹, Gordon Glendon¹¹⁰, Amanda E Toland¹¹¹, Anne-Marie Gerdes¹¹², Mads Thomassen¹¹³, Torben A Kruse¹¹³, Uffe Birk Jensen¹¹⁴, Anne-Bine Skytte¹¹⁵, Maria A Caligo¹¹⁶, Maria Soller¹¹⁷, Karin Henriksson¹¹⁸, von Anna Wachenfeldt¹¹⁹, Brita Arver¹¹⁹, Marie Stenmark-Askmal¹²⁰, Per Karlsson¹²¹, Yuan Chun Ding¹²², Susan L Neuhausen¹²², Mary Beattie¹²³, Paul DP Pharoah¹²⁴, Kirsten B Moysich¹²⁵, Katherine L Nathanson⁴, Beth Y Karlan¹²⁶, Jenny Gross¹²⁶,

Esther M John¹²⁷, Mary B Daly¹²⁸, Sandra M Buys¹²⁹, Melissa C Southey¹³⁰, John L Hopper¹³¹, Mary Beth Terry¹³², Wendy Chung¹³², Alexander F Miron¹³³, David Goldgar¹³⁴, Georgia Chenevix-Trench¹⁰, Douglas F Easton³, Irene L Andrulis¹³⁵ and Antonis C Antoniou^{3*} and for Breast Cancer Family Registry¹³⁶ and for EMBRACE³ and for GEMO Study Collaborators¹³⁷ and for HEBON¹³⁸ and for kConFab Investigators⁹³ and for Ontario Cancer Genetics Network¹³⁹ and for SWE-BRCA¹⁴⁰, for CIMBA

Abstract

Introduction: Previous studies have demonstrated that common breast cancer susceptibility alleles are differentially associated with breast cancer risk for *BRCA1* and/or *BRCA2* mutation carriers. It is currently unknown how these alleles are associated with different breast cancer subtypes in *BRCA1* and *BRCA2* mutation carriers defined by estrogen (ER) or progesterone receptor (PR) status of the tumour.

Methods: We used genotype data on up to 11,421 *BRCA1* and 7,080 *BRCA2* carriers, of whom 4,310 had been affected with breast cancer and had information on either ER or PR status of the tumour, to assess the associations of 12 loci with breast cancer tumour characteristics. Associations were evaluated using a retrospective cohort approach.

Results: The results suggested stronger associations with ER-positive breast cancer than ER-negative for 11 loci in both *BRCA1* and *BRCA2* carriers. Among *BRCA1* carriers, single nucleotide polymorphism (SNP) rs2981582 (*FGFR2*) exhibited the biggest difference based on ER status (per-allele hazard ratio (HR) for ER-positive = 1.35, 95% CI: 1.17 to 1.56 vs HR = 0.91, 95% CI: 0.85 to 0.98 for ER-negative, *P*-heterogeneity = 6.5×10^{-6}). In contrast, SNP rs2046210 at 6q25.1 near *ESR1* was primarily associated with ER-negative breast cancer risk for both *BRCA1* and *BRCA2* carriers. In *BRCA2* carriers, SNPs in *FGFR2*, *TOX3*, *LSP1*, *SLC4A7/NEK10*, 5p12, 2q35, and 1p11.2 were significantly associated with ER-positive but not ER-negative disease. Similar results were observed when differentiating breast cancer cases by PR status.

Conclusions: The associations of the 12 SNPs with risk for *BRCA1* and *BRCA2* carriers differ by ER-positive or ER-negative breast cancer status. The apparent differences in SNP associations between *BRCA1* and *BRCA2* carriers, and non-carriers, may be explicable by differences in the prevalence of tumour subtypes. As more risk modifying variants are identified, incorporating these associations into breast cancer subtype-specific risk models may improve clinical management for mutation carriers.

Introduction

Germline mutations in *BRCA1* and *BRCA2* confer high risks of breast, ovarian and other cancers [1-3] and account for 15 to 20% of the excess familial risk of breast cancer among first degree relatives [4,5]. Breast cancer risks for *BRCA1* and *BRCA2* mutation carriers have been estimated to range between 40 and 87% by age 70 [6-12] with population-based estimates tending to be lower than estimates based on families with multiple affected individuals [6,8]. Moreover, breast cancer risks for mutation carriers were found to vary according to the age at diagnosis and the type of cancer of the index patient involved in the family ascertainment [6,7,11]. Such evidence suggests that genetic or other risk factors that cluster in families modify the cancer risks conferred by *BRCA1* and *BRCA2* mutations.

A substantial body of work indicates that tumours arising in patients with germline *BRCA1* mutations are morphologically and genetically distinct from those arising in carriers of *BRCA2* mutations and from tumours

in patients lacking mutations. In gene expression studies, *BRCA1*-associated tumours are often classified as basal subtype tumours [13,14]. This is reflected in their higher grade, and morphologic features including lymphocytic infiltrate, pushing margins and syncytial growth. Being basal-like they express several markers that are normally expressed in the basal/myoepithelial cells of the breast, including stratified epithelial cytokeratins 5/6, 14 and 17. *BRCA1*-associated tumours are more likely to be estrogen receptor (ER), progesterone receptor (PR) and HER2 negative and to harbor mutations in the *TP53* gene than age-matched sporadic breast cancers [15,16]. *BRCA2*-associated tumours are also predominantly high-grade invasive ductal carcinomas of no special type but they often demonstrate a luminal phenotype despite their high histologic grade [13,17]. Adjusting for grade, *BRCA2*-associated tumours are more often ER-positive and are less likely, compared with controls, to express the basal cytokeratin CK5 or to overexpress HER2/neu protein [17].

Establishing the estrogen receptor status of a breast cancer (positive or negative) reflects a major subdivision in breast cancer type (at least five major sub-types are recognized) and it is becoming clear that the risk factors associated with breast cancer, both genetic and epidemiological, differ according to sub-type. Genome-wide association studies (GWAS) in breast cancer have identified several common alleles (single nucleotide polymorphisms (SNPs)) associated with an increased risk of breast cancer in the general population [18-25]. Many of these SNPs are associated with risk for ER-positive breast cancer, fewer have so far been associated with ER-negative breast cancer risk [26,27].

Known risk breast cancer susceptibility alleles have been genotyped in a large series of female *BRCA1* and *BRCA2* mutation carriers assembled by the Consortium of Investigators of Modifiers of *BRCA1/2* (CIMBA) to evaluate their associations with risk of breast cancer for mutation carriers. Of the 12 SNPs (rs2981582 in *FGFR2*, rs3803662 in *TOX3/TNRC9*, rs889312 in *MAP3K1*, rs13281615 at 8q24, rs381798 in *LSP1*, rs13387042 at 2q35, rs4973768 in *SLC4A7/NEK10*, rs10941679 at 5p12, rs6504950 in *STXBP4/COX11*, rs999737/rs10483813 in *RAD51L1*, rs2046210 at 6q25.1 and rs11249433 at 1p11.2) investigated so far, eight were associated with breast cancer risk for *BRCA2* carriers (all but SNPs at 8q24, *RAD51L1*, 6q25.1 and *STXBP4/COX11*), whereas only three SNPs (6q25.1, *TOX3/TNRC9* and 2q35) were associated with risk for *BRCA1* mutation carriers [28-31]. Work from the Breast Cancer Association Consortium and subsequent studies have demonstrated differences in the associations between these susceptibility loci and tumour characteristics in the general population [27,32]. These results suggest that the observed differences in the associations between *BRCA1* and *BRCA2* mutation carriers may reflect differences in the distribution of tumour characteristics in mutation carriers. It is currently unclear whether these polymorphisms are associated with different tumour characteristics within *BRCA1* and *BRCA2* mutation carriers.

As an adjunct to predictive testing for a high risk *BRCA1* or *BRCA2* gene mutation, more individualized risk estimates that take into account additional genetic and environmental modifiers will require a more detailed understanding of how these various risk factors interact. Understanding whether common genetic variants modify the risks of developing ER-positive or ER-negative breast cancer in *BRCA1* and *BRCA2* mutation carriers could potentially influence the clinical management of these individuals. For example, knowing that a *BRCA1* mutation carrier is more likely to develop ER-positive breast cancer (than most *BRCA1* mutation carriers), may influence the choice of management strategies, such as chemoprevention. In this study, we used

data from the CIMBA consortium to evaluate the associations between the 12 common breast cancer susceptibility alleles and risk for breast cancer defined by ER and PR status.

Materials and methods

Subjects

Subjects were *BRCA1* and *BRCA2* mutation carriers recruited by 36 study centres in Europe, North America and Australia (Table 1). All carriers participated in clinical or research studies at the host institutions, which have been approved by local ethics committees (list provided in Additional file 1, Table S1). Each committee granted approval for access and use of the medical records for the present analyses.

The large majority of carriers were recruited through cancer genetics clinics offering genetic testing, and enrolled into national or regional studies. Eligibility to participate in CIMBA is restricted to female carriers of pathogenic *BRCA1* or *BRCA2* mutations who were 18 years old or older at recruitment. Information collected included the year of birth; mutation description, including nucleotide position and base change; age at last follow-up; ages at breast and ovarian cancer diagnoses; and age or date at bilateral prophylactic mastectomy. Information was also available on the country of residence. Related individuals were identified through a unique family identifier. Women were included in the analysis if they carried mutations that were pathogenic according to generally recognized criteria. Only studies that provided tumour pathology information and had genotype information were included in the analysis. However, to maximise the available information, genotyped mutation carriers within those studies missing information on tumour characteristics were included in the analysis and their disease subtype was assumed to be missing at random (see statistical methods for details). Further details about the CIMBA initiative can be found elsewhere [33].

Tumour pathology data collection

Tumour pathology data were amalgamated from a range of sources, specifically patient pathology reports, medical records, pathology review data, tumour registry records and results from tissue microarrays. Estrogen and progesterone receptor status was provided as negative or positive, with supplementary immunohistochemistry scoring data and methodology provided when available. Based on definitions supplied, most centres employed a cut off of $\geq 10\%$ of tumour cells stained positive to define receptor positivity. To ensure consistency across studies, when information on the proportion of cells stained was available, we used the same cut-off to define ER and PR positive tumours. For a small number of cases where composite scoring methods based on the

Table 1 Number of mutation carriers by country grouping affection status and tumour marker characteristics

Country Group	BRCA1	BRCA2	Unaffected	Breast Cancer	ER-	ER+	PR-	PR+
Austria ¹	465	179	318	326	76	51	76	44
Australia ²	660	552	541	671	235	200	297	121
Canada ³	443	358	386	415	107	70	89	68
Denmark ⁴	507	319	463	363	98	93	79	45
France-Belgium-Spain ⁵	1,673	1,256	1,217	1,712	140	165	1,661	127
Finland ⁶	103	105	91	117	59	54	74	39
Germany ⁷	1,231	589	648	1,172	443	336	457	311
Iceland ⁸	0	135	24	111	21	57	18	57
Italy ⁹	994	666	686	974	203	251	231	216
Latvia-Lithuania-Russia ¹⁰	190	0	79	111	21	6	18	7
Sweden ¹¹	537	177	396	318	86	54	89	50
Netherlands ¹²	804	319	611	512	72	41	69	29
UK-Eire ¹³	1,107	866	1,008	965	268	239	175	104
USA ¹⁴	2,707	1,559	2,118	2,048	482	366	512	297
Total	11,421	7,080	8,686	9,815	2,311	1,983	2,345	1,515

Studies in country groups:1:MUV; 2:BCFR/KCONFAB; 3:OCGN/BCFR/INHERIT; 4:CBCS/OUH; 5: GEMO/CNIO/HCSC/ICO/MOD-SQUAD; 6:HEBCS; 7:GC-HBOC/DKFZ; 8: ILUH; 9:CONSTIT-TEAM/IOVHBOCS/PBCS; 10: NNPIO/BFBOCC; 11: SWE-BRCA; 12: HEBON; 13:EMBRACE/UKGRFOCR; 14:FCCC/GEMO/GEORGETOWN/MAGIC/MAYO/MSKCC/NCI/OSU-CCG/UCI/UCSF/UKGRFOCR/UPENN/WCRI

proportion and intensity of staining were available (Allred score, Remmele score and H-score), widely-accepted cut-offs were used (Additional file 1, Table S2). Consistency checks were performed to validate receptor data against supplementary scoring information if provided.

Genotyping

This analysis included genotype data on 12 SNPs that had been previously assessed for their associations with the overall risk of breast cancer for *BRCA1* and *BRCA2* mutation carriers in CIMBA. Genotyping was performed using either the iPLEX or Taqman platforms and has been described in detail in the previous reports [28-31]. To ensure genotyping consistency, all genotyping centres were required to adhere to the CIMBA genotyping quality control criteria which are described in detail online [34]. The 12 SNPs genotyped were rs2981582 in *FGFR2*, rs3803662 in *TOX3/TNRC9*, rs889312 in *MAP3K1*, rs3817198 in *LSP1*, rs13387042 at 2q35, rs13281615 at 8q24, rs4973768 near *SLC4A7/NEK10*, rs6504950 in the *STXBP4/COX11* region, rs2046210 near *ESR1* at 6q25.1 and rs11249433 at 1p11.2. A Taqman assay could not be adequately designed for SNP rs999737 in the *RAD51L1* region and studies using this platform genotyped the surrogate SNP rs10483813 (pair-wise $r^2 = 1$ with rs999737 based on HapMap CEU data). Data for these two SNPs were combined and treated as a single locus in the analysis of associations.

Statistical analysis

The aim of this study was to evaluate the associations between each genotype and breast cancer subtypes defined by tumour characteristics in *BRCA1* and *BRCA2* mutation carriers separately. The phenotype of each individual was defined by the age at diagnosis of breast cancer and its subtype or by age at last follow-up. Individuals were censored at the age of the first breast cancer diagnosis, ovarian cancer diagnosis, or bilateral prophylactic mastectomy or the age at last observation. Mutation carriers censored at ovarian cancer diagnosis were considered unaffected.

The analysis of risk modifiers in *BRCA1* and *BRCA2* mutation carriers is complicated by the fact that mutation carriers are not randomly sampled with respect to their disease status. Many carriers are sampled through families seen in genetic clinics. The first tested individual in a family is usually someone diagnosed with cancer at a relatively young age. Such study designs, therefore, tend to lead to an over-sampling of affected individuals, and standard analytical methods like Cox regression or case-control analysis may lead to biased estimates of the risk ratios [35]. This can be illustrated by considering an individual affected at age t . In a standard analysis of a cohort study or a case-control analysis, the SNP genotype for the individual will be compared with those of all individuals at risk at age t or in a case-control analysis, with controls randomly sampled from all possible at risk individuals. This analysis leads to consistent estimates of the hazard ratio or

odds ratio estimates. However, in the present design, mutation carriers are already selected on the basis of disease status (where affected individuals are over-sampled). If standard cohort analysis were applied to these data, it would lead to affected individuals at age t being compared to unaffected carriers selected on the basis of their future disease status. If the genotype is associated with the disease, the risk estimate will be biased to zero because too many affected individuals (in whom the at-risk genotype is overrepresented) are included in the comparison group. Simulation studies have shown that this effect can be quite marked [35]. To address this, a retrospective likelihood approach was previously proposed, which models the observed genotypes conditional on the disease phenotypes [36]. For the current analyses we have extended this method to model the simultaneous effect of each SNP on more than one tumour subtype. We briefly describe this method for the analysis of associations with ER-positive and ER-negative breast cancer, but the same principles apply for the analysis of associations with other tumour characteristics.

We modelled the likelihood of the observed genotypes and tumour subtype conditional on the disease status, that is:

$$\prod_{i=1}^n P(g_i, d_i | y(t_i)) = \prod_{i=1}^n \frac{P((y(t_i), d_i | g_i)) P(g_i)}{P(y(t_i))} \quad (1)$$

Where $y(t_i)$ is the disease phenotype for individual i at censoring age t_i (breast cancer at age t_i or unaffected at age t_i), d_i is the ER status (0 = negative, 1 = positive) and g_i the observed genotype of individual i ($g_i = 0, 1$ or 2 minor alleles) and n the number of subjects in the analysis. To allow for tumour characteristics we assumed that breast cancer consists of different disease subtypes, such that the total breast cancer incidence at age t_i , $\lambda(t_i)$, is the sum of the disease incidence for the subtypes, that is $\lambda(t_i) = \nu(t_i) + \mu(t_i)$, where $\nu(t_i)$ is the incidence for ER-negative disease and $\mu(t_i)$ is the incidence of ER-positive disease. We assumed that the subtype-specific incidences depend on the underlying genotype through a Cox-proportional hazards model: $\nu(t_i) = \nu_0(t_i) \exp(\beta^t \mathbf{z}_g)$ and $\mu(t_i) = \mu_0(t_i) \exp(\gamma^t \mathbf{z}_g)$ where $\nu_0(t_i)$ and $\mu_0(t_i)$ and are the baseline incidences for disease subtypes (ER-negative and ER-positive respectively), \mathbf{z}_g is the genotype vector for individual i and β and γ are the subtype specific genotype log-risk ratios (for ER-negative and ER-positive breast cancer respectively). The probabilities of developing ER-positive and ER-negative breast cancer conditional on the underlying genotype were assumed to be independent. We further assumed that, if tumour

subtype is unknown, the information is missing at random with respect to genotype. Then for each individual:

$$P((y(t_i), d_i | g_i)) = \begin{aligned} &= (\nu_0(t_i) \exp(\beta^t \mathbf{z}_g))^{O_i(1-d_i)} \times (\mu_0(t_i) \exp(\gamma^t \mathbf{z}_g))^{O_i d_i} \times \\ &\exp\left(-\sum_{u=0}^{t_i-1} (\mu_0(u) \exp(\gamma^t \mathbf{z}_g) + \nu_0(u) \exp(\beta^t \mathbf{z}_g))\right) \text{ if } d_i = 0, 1 \\ &= (\mu_0(t_i) \exp(\gamma^t \mathbf{z}_g) + \nu_0(t_i) \exp(\beta^t \mathbf{z}_g))^{O_i} \times \\ &\exp\left(-\sum_{u=0}^{t_i-1} (\mu_0(u) \exp(\gamma^t \mathbf{z}_g) + \nu_0(u) \exp(\beta^t \mathbf{z}_g))\right) \text{ if } d_i = \text{unknown} \end{aligned}$$

were $O_i = 0$ if unaffected and $O_i = 1$ if affected. Thus, the above formulation allows use of all mutation carriers irrespective of whether the tumour subtype is observed or not. The baseline incidences for each disease subtype ($\nu_0(t_i)$ and $\mu_0(t_i)$) are unknown. However, it is possible to solve for those recursively by constraining the overall breast cancer incidence for mutation carriers $\lambda(t)$, to agree with external estimates as previously demonstrated [37,38] and by imposing a further constraint on the ratio of the observed ER-positive to ER-negative breast cancers in each age group:

$$\begin{aligned} \frac{P(\text{ERpositive at age } t)}{P(\text{ERnegative at age } t)} &= \frac{\pi_+(t)}{\pi_-(t)} = \\ &= \frac{\sum_g P(g) \mu_0(t) \exp(\gamma^t \mathbf{z}_g) \exp\left(-\sum_{u=0}^{t-1} (\mu_0(u) \exp(\gamma^t \mathbf{z}_g) + \nu_0(u) \exp(\beta^t \mathbf{z}_g))\right)}{\sum_g P(g) \nu_0(t) \exp(\beta^t \mathbf{z}_g) \exp\left(-\sum_{u=0}^{t-1} (\mu_0(u) \exp(\gamma^t \mathbf{z}_g) + \nu_0(u) \exp(\beta^t \mathbf{z}_g))\right)} \end{aligned}$$

The likelihood in equation 1 can then be maximised jointly over the log-risk ratios β and γ , genotype frequencies $P(g)$ and the age and subtype-specific frequencies $\pi_+(t)$ and $\pi_-(t)$. This likelihood is based on the assumption that the ascertainment of mutation carriers is dependent on the overall disease phenotype (breast cancer) but not on tumour subtypes. This allows the subtype frequencies $\pi_+(t)$ and $\pi_-(t)$ to be estimated within the dataset. Relaxing this assumption and conditioning also on tumour subtype requires external estimates for the age and subtype-specific frequencies $\pi_+(t)$ and $\pi_-(t)$.

The effect of each SNP was modelled either as a per-allele HR (multiplicative model) or as separate HRs for heterozygotes and homozygotes, and these were estimated on the logarithmic scale. Heterogeneity in the hazard ratios between tumour subtypes was examined by fitting models where $\nu(t_i) = \nu_0(t_i) \exp(\beta_{1g})$ and $\mu(t_i) = \mu_0(t_i) \exp(\beta_1 + \beta_2)g$ with $g = 0, 1$ and 2 (for 0, 1, 2 copies of the minor allele respectively) and testing for $\beta_2 = 0$. Analyses were carried out with the pedigree-analysis software MENDEL [39]. All analyses were stratified by country of residence and used calendar-year- and cohort-specific cancer incidences for *BRCA1* and *BRCA2* [40]. For this purpose, a stratified version of the retrospective likelihood (equation 1) was derived as described previously [36]. Countries with small numbers

of mutation carriers were grouped together. We used a robust variance-estimation approach to allow for the non-independence among related mutation carriers [41].

Predicted breast cancer risks by ER status

Based on our results we computed the predicted absolute risk of developing ER-negative and ER-positive breast cancer for *BRCA1* and *BRCA2* mutation carriers by the combined 12 SNP profile. For each individual we derived an empirical score, based on the per-allele log-relative hazard estimates for each genotype, which was of the form $\sum_{j=1}^{12} \beta_j g_j$ where β_j is the per-allele log-hazard estimate for locus j and g_j is the genotype at the same locus (taking values 0, 1 and 2). This assumes a multiplicative model for the combined SNP associations. This is a reasonable assumption given that previous analyses found no evidence of departure from the multiplicative model [35]. Scores were calculated for ER-positive and ER-negative disease, separately for *BRCA1* and *BRCA2* mutation carriers. The empirical distribution of the derived score was then used to compute the subtype specific incidence associated with each multilocus genotype as described previously [31]. We reported the absolute risks of developing ER-specific breast cancer at the 5th, 50th and 95th percentiles of the empirical distribution of the SNP profile.

Results

A total of 11,421 *BRCA1* and 7,080 *BRCA2* mutation carriers from 36 studies had been successfully genotyped for at least one of the 12 SNPs and were eligible for inclusion in these analyses. 9,815 *BRCA1* and *BRCA2* mutation carriers were censored at a first invasive breast cancer diagnosis, of whom 4,310 had information on either ER or PR (Table 1).

Associations with ER status - *BRCA1* mutation carriers

There were significant differences in the HR for ER-positive and ER-negative disease for *BRCA1* mutation carriers for two SNPs (Table 2). The *FGFR2* SNP rs2981582 exhibited the clearest difference with a strong association for ER-positive disease but not ER-negative disease (per allele HR = 1.35, 95% CI: 1.17 to 1.56, for ER-positive compared with HR = 0.91, 95% CI: 0.85 to 0.98 for ER-negative, P -heterogeneity = 6.5×10^{-6}). The *SLC4A7/NEK10* SNP rs4973768 also exhibited a similar pattern (ER-positive: per-allele HR = 1.17, 95% CI: 1.03 to 1.133, compared with ER-negative: per-allele HR = 0.99, 95% CI: 0.93 to 1.06, P -heterogeneity = 0.027). Although there was no significant evidence of differences between the HRs for ER-positive and ER-negative breast cancer, the *TOX3/TNRC9* SNP rs3803662 was

significantly associated with the risk of ER-positive disease (per-allele HR = 1.25, 95% CI: 1.06 to 1.46, P -trend = 0.0062) but not with risk for ER-negative breast cancer (per-allele HR = 1.05, 95% CI: 0.97 to 1.13, P -trend = 0.21). *LSP1* SNP rs3817198 was associated with the risk of ER-negative breast cancer (per-allele HR = 1.07, 95% CI: 1.00 to 1.07, P -trend = 0.047) but not with risk of ER-positive breast cancer (per-allele HR = 1.07, 95% CI: 0.93 to 1.22, P -trend = 0.33, P -het = 0.98). The 6q25.1 SNP rs2046210 near *ESR1* was associated with the risk for both ER-negative (per-allele HR = 1.19, 95% CI: 1.11 to 1.27, P -trend = 2.4×10^{-7}) and ER-positive (per-allele HR = 1.14, 95% CI: 1.01 to 1.30, P -trend = 0.043) breast cancer. There was no significant evidence of association with either ER-negative or ER-positive breast cancer for any of the other SNPs, although the HR estimates tended to be higher for ER-positive breast cancer (for example, SNPs rs13387042 at 2q35 and rs13281615 at 8q24).

Associations with ER status - *BRCA2* mutation carriers

Only SNP rs2046210 at 6q25.1 exhibited differential associations between ER-positive and ER-negative breast cancer for *BRCA2* mutation carriers (P -heterogeneity = 0.045, Table 3). The per-allele HR for ER-negative disease was estimated to be 1.17 (95% CI: 0.99 to 1.38) whereas the per-allele HR for ER-positive breast cancer was 0.97 (95% CI: 0.89 to 1.05). Although there were no significant differences in the associations between the two types of disease for *BRCA2* mutation carriers, the HR estimates for ER-positive disease tended to be larger compared to ER-negative breast cancer. SNPs at/near *FGFR2*, *TOX3/TNRC9*, *MAP3K1*, *LSP1*, 2q35, *SLC4A7/NEK10*, 5p12 and 1p11.2 were associated with ER-positive breast cancer for *BRCA2* mutation carriers (using either a per-allele or 2 df genotype test). The strongest associations were for the *FGFR2* rs2981582 SNP (HR for ER-positive breast cancer = 1.35, 95% CI: 1.23 to 1.48, P -trend = 1.4×10^{-10}) and *TOX3/TNRC9* SNP rs3803662 (HR for ER-positive breast cancer = 1.28, 95% CI: 1.16 to 1.41, P -trend = 1.5×10^{-6}). Only SNPs at or near *MAP3K1*, *STXBP4/COX11* and 6q25.1 were associated with the risk of ER-negative breast cancer for *BRCA2* mutation carriers.

Associations with PR status - *BRCA1* mutation carriers

The general pattern of associations with PR-positive and PR-negative breast cancer for *BRCA1* mutation carriers (Additional file 1, Table S3) was similar to that seen for ER status. Significant differences in the associations between PR-positive and PR-negative breast cancer were observed for three SNPs. The minor allele of *FGFR2* SNP rs2981582 was associated with a significantly higher risk for PR-positive breast cancer for *BRCA1*

Table 2 Genotype and per-allele hazard ratio estimates by estrogen receptor status for *BRCA1* mutation carriers

Genotype	Unaffected, N (%)	Affected by subtype, N (%)			ER-			ER+			P-het
		ER-	ER+	Unknown	HR	95% CI	P-trend	HR	95% CI	P-trend	
<i>FGFR2</i> rs2981582											
GG	1301 (36.2)	447 (40.0)	104 (29.6)	869 (35.1)	1.00			1.00			
GA	1,721 (47.9)	516 (46.2)	166 (47.3)	1190 (48.1)	0.93	0.83 to 1.03		1.24	0.98 to 1.57		
AA	573 (15.9)	154 (13.8)	81 (23.1)	416 (16.8)	0.82	0.70 to 0.96		1.85	1.40 to 2.44		
Per-allele					0.91	0.85 to 0.98	0.01	1.35	1.17 to 1.56	4 × 10⁻⁵	6.5 × 10⁻⁶
<i>TOX3/TNRC9</i> rs3803662											
CC	1,811 (52.0)	545 (49.5)	154 (45.7)	1195 (50.2)	1.00			1.00			
CT	1,405 (40.3)	461 (41.8)	143 (42.4)	987 (41.5)	1.06	0.96 to 1.18		1.20	0.97 to 1.50		
TT	269 (7.7)	96 (8.7)	40 (11.9)	199 (8.4)	1.09	0.90 to 1.31		1.61	1.14 to 2.27		
Per-allele					1.05	0.97 to 1.13	0.21	1.25	1.06 to 1.46	0.0062	0.07
<i>MAP3K1</i> rs889312											
AA	1,858 (49.6)	569 (49.7)	186 (52.1)	1319 (51.6)	1.00			1.00			
AC	1,552 (41.4)	480 (41.9)	136 (38.1)	987 (38.6)	0.97	0.87 to 1.07		0.86	0.69 to 1.06		
CC	336 (9.0)	97 (8.5)	35 (9.8)	250 (9.8)	0.97	0.81 to 1.16		0.97	0.69 to 1.35		
Per-allele					0.98	0.91 to 1.06	0.56	0.97	0.83 to 1.13	0.69	0.92
<i>LSP1</i> rs3817198											
TT	1,894 (47.4)	652 (45.4)	195 (44.5)	1205 (43.7)	1.00			1.00			
TC	1,680 (42.0)	629 (43.8)	197 (45.0)	1239 (44.9)	1.09	0.99 to 1.20		1.11	0.92 to 1.35		
CC	422 (10.6)	154 (10.7)	46 (10.5)	315 (11.4)	1.13	0.97 to 1.33		1.10	0.80 to 1.50		
Per-allele					1.07	1.00 to 1.15	0.047	1.07	0.93 to 1.22	0.33	0.98
2q35 rs13387042											
GG	924 (24.0)	301 (22.1)	93 (22.2)	576 (21.4)	1.00			1.00			
GA	1,855 (48.3)	723 (53.1)	194 (46.3)	1370 (50.9)	1.17	1.04 to 1.32		1.02	0.81 to 1.30		
AA	1,064 (27.7)	338 (24.8)	132 (31.5)	745 (27.7)	0.96	0.84 to 1.11		1.25	0.97 to 1.62		
Per-allele					0.98	0.91 to 1.04	0.48	1.13	0.99 to 1.28	0.075	0.065
8q24 rs13281615											
AA	1,319 (32.8)	502 (35.9)	143 (33.7)	897 (32.5)	1.00			1.0			
AG	2,008 (50.0)	657 (47.0)	198 (46.7)	1,364 (49.5)	0.98	0.88 to 1.08		0.97	0.79 to 1.19		
GG	691 (17.9)	238 (17.0)	83 (19.6)	496 (18.0)	1.00	0.87 to 1.16		1.17	0.89 to 1.53		
Per-allele					1.00	0.93 to 1.07	0.93	1.06	0.93 to 1.22	0.38	0.43
SLC4A7/NEK10 rs4973768											
CC	1,148 (26.2)	406 (27.2)	103 (22.2)	691 (24.7)	1.00			1.00			
CT	2,205 (50.4)	736 (49.3)	235 (50.5)	1,399 (50.1)	0.98	0.88 to 1.08		1.20	0.96 to 1.51		
TT	1,024 (23.4)	350 (23.5)	127 (27.3)	703 (25.2)	0.99	0.87 to 1.12		1.38	1.07 to 1.77		
Per-allele					0.99	0.93 to 1.06	0.83	1.17	1.03 to 1.33	0.013	0.027
STXBP4/COX11 rs6504950											
GG	2,346 (53.1)	814 (53.2)	252 (52.9)	1,502 (53.1)	1.00			1.00			
GA	1,737 (39.3)	593 (37.8)	191 (40.1)	1,127 (39.8)	1.00	0.91 to 1.10		1.03	0.86 to 1.24		
AA	333 (7.5)	122 (8.0)	33 (6.9)	200 (7.1)	1.04	0.88 to 1.23		0.94	0.65 to 1.34		
Per-allele					1.01	0.94 to 1.09	0.77	1.00	0.87 to 1.15	0.97	0.87
5p12 rs10941679											
AA	2,211 (55.8)	815 (57.3)	271 (61.0)	1,529 (56.5)	1.00			1.00			
AG	1,472 (37.1)	517 (36.4)	145 (32.7)	1,001 (37.0)	0.99	0.90 to 1.09		0.84	0.69 to 1.02		
GG	280 (7.1)	90 (6.3)	28 (6.3)	177 (6.5)	0.89	0.73 to 1.08		0.84	0.58 to 1.20		
Per-allele					0.97	0.90 to 1.04	0.39	0.88	0.75 to 1.02	0.08	0.26
6q25.1 - rs2046210											
CC	1,886 (43.3)	567 (38.2)	158 (36.0)	1,007 (37.0)	1.00			1.00			
TC	1,919 (44.1)	718 (48.3)	232 (52.9)	1,305 (48.0)	1.21	1.10 to 1.33		1.37	1.13 to 1.67		
TT	547 (12.6)	201 (13.5)	49 (11.2)	409 (15.0)	1.39	1.21 to 1.59		1.11	0.81 to 1.53		
Per-allele					1.19	1.11 to 1.27	2.4 × 10⁻⁷	1.14	1.01 to 1.30	0.043	0.60

Table 2 Genotype and per-allele hazard ratio estimates by estrogen receptor status for *BRCA1* mutation carriers (Continued)

1p11.2 - rs11249433										
TT	1,491 (34.1)	506 (33.7)	144 (32.4)	973 (35.1)	1.00			1.00		
CT	2,133 (48.7)	745 (49.7)	245 (55.2)	1,342 (48.5)	0.97	0.88 to 1.07		1.10	0.90 to 1.34	
CC	752 (17.2)	248 (16.5)	55 (12.4)	455 (16.4)	0.99	0.87 to 1.13		0.76	0.56 to 1.03	
Per-allele					0.99	0.93 to 1.06	0.79	0.92	0.81 to 1.05	0.22 0.35
<i>RAD51L1</i> - rs999737/rs10483813										
CC/TT	2,335 (61.5)	760 (64.1)	212 (62.5)	1,551 (62.5)	1.00			1.00		
TC/AT	1,294 (34.1)	370 (31.2)	113 (33.3)	819 (33.0)	0.92	0.83 to 1.03		1.00	0.80 to 1.24	
TT/AA	170 (4.5)	56 (4.7)	14 (4.1)	110 (4.4)	1.02	0.81 to 1.28		0.93	0.56 to 1.56	
per allele					0.96	0.88 to 1.04	0.34	0.98	0.82 to 1.18	0.87 0.81

P-het, Heterogeneity *P*-value; ER-, estrogen receptor negative; ER+, estrogen receptor positive

mutation carriers (per-allele HR for PR-positive = 1.29, 95% CI: 1.10 to 1.51, HR for PR-negative = 0.93, 95% CI: 0.87 to 1.00, *P*-heterogeneity = 7×10^{-4}). Allele "A" of SNP rs13387042 at 2q35 was associated with a significantly higher risk of PR-positive breast cancer for *BRCA1* mutation carriers (HR for PR-positive breast cancer = 1.16, 95% CI: 1.01 to 1.33; HR for PR-negative = 0.97, 95% CI: 0.91 to 1.04, *P*-heterogeneity = 0.034). Although the *RAD51L1* SNP showed no differential associations with ER-status, there was evidence that the minor allele of this SNP was associated with a lower risk of PR-positive breast cancer (HR for PR-positive = 0.79, 95% CI: 0.61 to 0.95; HR for PR-negative = 1.02, 95% CI: 0.94 to 1.11; *P*-heterogeneity = 0.027). The only SNPs associated with risk for PR-negative breast cancer were SNPs at 6q25.1 (per-allele HR = 1.19, 95% CI: 1.11 to 1.27, *P*-trend = 3.7×10^{-7}), and in *LSP1* (per-allele HR = 1.09, 95% CI: 1.01 to 1.16, *P*-trend = 0.017), but these were not significantly different from the associations with PR-positive breast cancer for *BRCA1* mutation carriers.

Associations with PR status - *BRCA2* mutation carriers

Only two SNPs demonstrated significant differences in the associations with PR-positive and PR-negative breast cancer for *BRCA2* mutation carriers (Additional file 1, Table S4). SNP rs13387042 at 2q35 was associated with a higher risk of PR-positive breast cancer (per-allele HR for PR-positive = 1.14, 95% CI: 1.03 to 1.25; per-allele HR for PR-negative = 0.92, 95% CI: 0.81 to 1.04, *P*-heterogeneity = 0.009). SNP rs10941679 at 5p12 was also associated with a higher risk of PR-positive breast cancer for *BRCA2* mutation carriers (per-allele HR for PR-positive = 1.15, 95% CI: 1.03 to 1.27, HR for PR-negative = 0.94, 95% CI: 0.81 to 1.09, *P*-heterogeneity = 0.028). SNPs near or at *FGFR2*, *TOX3/TNRC9*, *LSP1* were associated with both PR-negative and PR-positive breast cancer, whereas the 1p11.2 SNP was associated with risk for PR-negative breast cancer. *MAP3K1* and *SLC4A7*/

NEK10 were associated only with risk of PR-positive breast cancer among *BRCA2* mutation carriers.

Absolute risks of developing ER-positive and ER-negative breast cancer by SNP profile

Using the estimated HRs for ER-positive and ER-negative breast cancer for *BRCA1* and *BRCA2* mutation carriers, we computed the predicted absolute risk of developing ER-negative and ER-positive breast cancer at various percentiles of the combined SNP distribution. The SNP profile distribution is different for each disease subtype and mutation. We note that SNPs for which the per-allele HR estimates are close to 1.0 contribute little to the predicted ER-specific risks. Figure 1 shows the predicted risks of developing ER-negative and ER-positive breast cancer for *BRCA1* and *BRCA2* mutation carriers at the 5th, 50th and 95th percentiles of the empirical risk distribution of the combined SNP profile. A *BRCA1* mutation carrier at the 5th percentile of the SNP profile distribution would be at 43% risk of developing ER-negative breast cancer by age 80 compared with 60% for *BRCA1* mutation carriers at the 95th percentile of the risk distribution. The risks of developing ER-positive breast cancer would be 18% and 46% by age 80 at the 5th and 95th percentiles of the ER-positive breast cancer risk distribution. *BRCA2* mutation carriers at the 5th percentile of the ER-negative breast cancer risk distribution are predicted to have a 22% risk of developing ER-negative breast cancer by age 80 compared with 39% for the 95th percentile of the risk distribution. The risks of developing ER-positive breast cancer by age 80 for *BRCA2* carriers varied from 33% to 70% at the 5th and 95th percentiles of the ER-positive risk distribution respectively.

Discussion

This is the first report to investigate the associations between 12 common breast cancer susceptibility alleles and ER and PR status of breast tumours in *BRCA1* and

Table 3 Genotype and per-allele hazard ratio estimates by estrogen receptor status for *BRCA2* mutation carriers

Genotype	Unaffected, N (%)	Affected by subtype, N (%)			HR	ER- 95%CI	P-trend	ER+		P-trend	P-het
		ER-	ER+	Unknown				HR	95% CI		
<i>FGFR2 rs2981582</i>											
GG	794 (37.8)	86 (32.7)	248 (29.5)	457 (29.8)	1.00			1.00			
GA	987 (47.0)	137 (52.1)	419 (49.8)	755 (49.3)	1.28	0.99 to 1.67		1.35	1.17 to 1.55		
AA	321 (15.3)	40 (15.2)	174 (20.7)	320 (20.9)	1.23	0.85 to 1.78		1.81	1.51 to 2.18		
Per-allele					1.14	0.97 to 1.35	0.12	1.35	1.23 to 1.48	1.4 × 10⁻¹⁰	0.097
<i>TOX3/TNRC9 rs3803662</i>											
CC	1,088 (53.4)	136 (53.3)	377 (46.3)	702 (48.2)	1.00			1.00			
CT	792 (38.9)	96 (37.7)	361 (44.3)	604 (41.5)	0.98	0.75 to 1.27		1.33	1.17 to 1.53		
TT	157 (7.7)	23 (9.0)	77 (9.5)	150 (10.3)	1.27	0.83 to 1.93		1.54	1.22 to 1.95		
Per-allele					1.06	0.88 to 1.29	0.53	1.28	1.16 to 1.41	1.5 × 10⁻⁶	0.11
<i>MAP3K1 rs889312</i>											
AA	1,107 (51.1)	121 (45.7)	430 (50.3)	746 (47.7)	1.00			1.00			
AC	888 (41.0)	120 (45.3)	349 (40.8)	646 (41.3)	1.23	0.96 to 1.59		1.03	0.90 to 1.17		
CC	170 (7.9)	24 (9.1)	76 (8.9)	172 (11.0)	1.42	0.93 to 2.16		1.29	1.03 to 1.62		
Per-allele					1.21	1.01 to 1.45	0.039	1.09	0.99 to 1.21	0.08	0.35
<i>LSP1 rs3817198</i>											
TT	1,075 (46.1)	142 (44.4)	429 (42.0)	718 (42.7)	1.00			1.00			
TC	1,005 (43.1)	146 (45.6)	466 (45.6)	759 (45.2)	1.08	0.86 to 1.36		1.14	1.01 to 1.29		
CC	252 (10.8)	32 (10.0)	127 (12.4)	203 (12.1)	1.02	0.68 to 1.51		1.39	1.14 to 1.70		
Per-allele					1.03	0.87 to 1.22	0.70	1.17	1.07 to 1.28	5.5 × 10⁻⁴	0.20
<i>2q35 rs13387042</i>											
GG	571 (25.3)	71 (23.0)	216 (22.0)	382 (23.1)	1.00			1.00			
GA	1,080 (47.8)	156 (50.5)	500 (50.8)	809 (48.8)	1.12	0.85 to 1.47		1.18	1.01 to 1.36		
AA	608 (26.9)	82 (26.5)	268 (27.2)	466 (28.1)	1.06	0.78 to 1.45		1.13	0.95 to 1.34		
Per-allele					1.03	0.87 to 1.19	0.71	1.06	0.97 to 1.15	0.20	0.75
<i>8q24 rs13281615</i>											
AA	794 (34.1)	99 (31.6)	317 (31.7)	524 (31.3)	1.00			1.00			
AG	1,156 (49.6)	165 (52.7)	511 (51.1)	837 (49.9)	1.08	0.85 to 1.38		1.05	0.92 to 1.21		
GG	382 (16.4)	49 (15.7)	172 (17.2)	315 (18.8)	1.05	0.75 to 1.46		1.13	0.94 to 1.35		
Per-allele					1.04	0.89 to 1.21	0.66	1.06	0.97 to 1.16	0.19	0.80
<i>SLC4A7/NEK10 rs4973768</i>											
CC	669 (26.5)	82 (24.9)	251 (22.6)	401 (23.5)	1.00			1.00			
CT	1,241 (49.1)	164 (49.9)	546 (49.4)	829 (48.7)	1.05	0.81 to 1.36		1.14	0.99 to 1.31		
TT	618 (24.5)	83 (25.2)	311 (28.0)	474 (27.8)	1.04	0.77 to 1.41		1.27	1.08 to 1.50		
Per-allele					1.02	0.88 to 1.19	0.78	1.13	1.04 to 1.22	0.0043	0.25

Table 3 Genotype and per-allele hazard ratio estimates by estrogen receptor status for *BRCA2* mutation carriers (Continued)


STXBP4/COX11 rs6504950										
GG	1,420 (55.6)	171 (51.0)	601 (53.1)	896 (52.5)	1.00					1.00
GA	951 (37.2)	145 (43.3)	444 (39.3)	684 (40.1)	1.27	1.03 to 1.58			1.09	0.97 to 1.23
AA	184 (7.2)	19 (5.7)	86 (7.6)	127 (7.4)	0.84	0.54 to 1.30			1.08	0.87 to 1.34
Per-allele					1.07	0.92 to 1.25	0.36	1.06	0.97 to 1.16	0.19 0.91
5p12 rs10941679										
AA	1,372 (58.6)	176 (54.8)	584 (54.8)	924 (56.3)	1.00					1.00
AG	824 (35.2)	122 (38.0)	425 (39.9)	622 (37.9)	1.08	0.86 to 1.37			1.15	1.01 to 1.30
GG	146 (6.2)	23 (7.2)	57 (5.4)	94 (5.7)	1.25	0.82 to 1.91			0.94	0.73 to 1.22
Per-allele					1.10	0.92 to 1.31	0.28	1.06	0.96 to 1.17	0.23 0.70
6q25.1 - rs2046210										
CC	985 (39.8)	121 (39.2)	466 (42.1)	634 (37.7)	1.00					1.00
TC	1,165 (47.1)	132 (42.7)	499 (45.1)	802 (47.7)	0.99	0.78 to 1.12			0.96	0.85 to 1.08
TT	324 (13.1)	56 (18.1)	141 (12.8)	247 (14.7)	1.47	1.08 to 2.01			0.94	0.78 to 1.12
Per-allele					1.17	0.99 to 1.38	0.059	0.97	0.89 to 1.05	0.41 0.045
1p11.2 - rs11249433										
TT	895 (35.9)	107 (33.6)	345 (31.4)	599 (34.7)	1.00					1.00
CT	1,226 (49.2)	160 (50.3)	553 (50.3)	843 (48.9)	1.00	0.79 to 1.28			1.08	0.96 to 1.23
CC	371 (14.9)	51 (16.0)	202 (18.4)	282 (16.4)	1.00	0.73 to 1.40			1.27	1.08 to 1.50
Per-allele					1.00	0.86 to 1.17	0.98	1.12	1.03 to 1.22	0.0065 0.23
RAD51L1 - rs999737/ rs10483813										
<i>BRCA1</i>	CC/TT	1,368 (59.5)	167 (59.0)	589 (61.4)	1,000 (62.0)	1.00				1.00
	TC/AT	789 (34.3)	104 (36.8)	323 (33.6)	534 (33.1)	1.06	0.84 to 1.35		0.94	0.83 to 1.07
	TT/AA	141 (6.1)	12 (4.2)	48 (5.0)	80 (4.9)	0.74	0.42 to 1.32		0.87	0.67 to 1.14
Per-allele					0.97	0.80 to 1.17	0.73	0.94	0.85 to 1.03	0.20 0.77

P-het, Heterogeneity P-value; ER-, estrogen receptor negative; ER+, estrogen receptor positive

BRCA2 mutation carriers. The analysis was made possible by the availability of a large, combined dataset with genotype and tumour pathology information in mutation carriers collated through the CIMBA consortium.

The majority of the SNPs examined demonstrated stronger associations with ER-positive breast cancer for both *BRCA1* and *BRCA2* mutation carriers (Figure 2). Only rs2046210 on 6q25.1 exhibited stronger evidence for ER-negative disease. Among *BRCA1* mutation carriers, the most marked difference was for SNP rs2981582 in *FGFR2*, which was strongly associated with ER-positive breast cancer and exhibited no evidence of an association with ER-negative breast cancer ($P = 6.5 \times 10^{-6}$). Previous analyses of this polymorphism in


mutation carriers failed to find an association with the overall risk of breast cancer for *BRCA1* mutation carriers, but found an association with risk for *BRCA2* mutation carriers [29,31]. Our results suggest that rs2981582 in *FGFR2* also modifies ER-positive breast cancer risk for *BRCA1* mutation carriers to a similar relative extent as in *BRCA2* mutation carriers and ER-positive disease in the general population [27,32]. Similar patterns were observed for SNPs rs3803662 in *TOX3/TNRC9* and rs4973768 in *SLC4A7/NEK10* in which the associations were predominantly with ER-positive breast cancer for both *BRCA1* and *BRCA2* mutation carriers, in line with results from studies of breast cancer in the general population [18,21,27]. The



HR estimates for ER-positive breast cancer in *BRCA1* and *BRCA2* mutation carriers for these SNPs were very similar.

Among the 12 SNPs investigated in this report, SNP rs2046210 at 6q25.1 exhibited the strongest association with the risk of breast cancer for *BRCA1* mutation carriers in previous analyses, and was not associated with risk for *BRCA2* mutation carriers [28]. The current results suggest that this was mainly driven by an

association with ER-negative breast cancer risk. This observation is again consistent with the effects seen in population-based studies, in which the relative risk is higher for ER-negative than ER-positive disease [42,43] (Alison Dunning, personal communication). There was some evidence that the 6q25.1 SNP is also associated with ER-negative disease cancer subtype in *BRCA2* mutation carriers, although the estimates for ER-negative breast cancer in *BRCA2* mutation carriers are



imprecise due to the relatively small sample size. In addition to the 12 loci investigated in this report, a recently identified locus at 19p13 also appears to be predominantly associated with ER-negative breast cancer [44].

The patterns of association between the SNPs and PR tumour status were similar to those observed for ER, which is not surprising given that ER and PR expression are highly correlated. There were, however, two notable exceptions. The 2q35 SNP rs13387042 demonstrated significantly stronger associations with PR-positive than PR-negative breast cancer for both *BRCA1* and *BRCA2* mutation carriers ($P = 0.034$ and $P = 0.0086$, for PR-positive for *BRCA1* and *BRCA2* respectively), suggesting

this SNP may be more relevant for *BRCA1* and *BRCA2* tumours expressing PR. However, a population-based study has found this SNP is also associated with PR-negative breast cancer [45]. Furthermore, the *RAD51L1* locus was associated with PR-positive breast cancer for *BRCA1* mutation carriers and the magnitude of the association was similar to that observed in the general population [23] (A.B. Spurdle, personal communication).

Previous studies demonstrated that SNPs, which are associated with ER-positive breast cancer in the general population, tend to be associated with the breast cancer risk for *BRCA2* mutation carriers and SNPs, which are associated with ER-negative breast cancer in the general population, tend to be associated with the breast cancer

risk for *BRCA1* mutation carriers [27,31,44-46]. The current results demonstrate that despite lack of an association between a SNP and the overall breast cancer risk for *BRCA1* or *BRCA2* mutation carriers, residual associations exist with specific disease subtypes. Figure 2 summarises the association patterns in *BRCA1* and *BRCA2* mutation carriers. The HR estimates for ER-positive and ER-negative breast cancer among *BRCA1* mutation carriers appear to be different (intraclass correlation coefficient (ICC) approximately 0), as are the HR estimates for ER-positive and ER-negative breast cancer among *BRCA2* mutation carriers (ICC = 0.13). On the other hand the HR estimates for ER-positive breast cancer among *BRCA1* and *BRCA2* mutation carriers appear to be more similar (ICC = 0.65). There is, however, little correlation in the HR estimates for ER-negative breast cancer among *BRCA1* and *BRCA2* mutation carriers (ICC = 0.05). However, SNP 6q25.1, which is mainly associated with ER-negative disease in *BRCA1* mutation carriers, is estimated to confer similar HRs for ER-negative breast cancer for both *BRCA1* and *BRCA2* mutation carriers. These associations are mainly in the same direction and of similar magnitude to those observed with breast cancer in the general population stratified by ER expression status. Taken together, these findings are consistent with a model in which these SNPs and *BRCA1* or *BRCA2* mutations combine multiplicatively on the risk for ER-positive or ER-negative breast cancer [47]. Hence, the apparent differences in the strength of the SNP associations by *BRCA1* and *BRCA2* mutation status can be explained once tumour subtype is taken into account.

The major strength of the current study is the large sample of *BRCA1* and *BRCA2* mutation carriers with SNP and tumour marker information. Despite the large sample size, ER and PR marker information was only available for approximately 30% of the mutation carriers that had been diagnosed with breast cancer. The sample sizes for tumour subtypes, while still large, were, therefore, much smaller than were available for analyses of breast cancer risk overall, particularly for ER-positive breast cancer in *BRCA1* carriers and ER-negative breast cancer in *BRCA2* carriers. However, by analysing the data using a retrospective cohort approach and analysing the associations with ER-positive and ER-negative disease simultaneously we were able to include all mutation carriers in the analysis, including affected individuals with missing ER status, thus maximizing the available information. Ongoing efforts by CIMBA aim to increase the proportion of mutation carriers diagnosed with breast cancer who also have available tumour pathology information. This will enable us to assess the associations with breast cancer subtypes with greater precision.

The majority of the mutation carriers in CIMBA are identified through clinical genetics centers and, therefore, the source of information or definition of tumour marker status could vary across studies. This heterogeneity in classification may attenuate some of the differences by tumour type. For example, most commonly, a cut-off of 10% of cells staining was taken to denote positivity for ER and PR by the centers without further information on intensity or proportion of positive tumour nuclei and this was used for all our analyses; however, in centers that use the Allred score, a value of > 2 denoted positivity, which may reflect as few as 1% of cells staining. In fact, recent recommendations suggest that ER and PgR assays be considered positive, for therapeutic purposes, if there are at least 1% positive tumour nuclei [48], but these data were not available for the majority of carriers in our samples to enable reclassification. It has been shown, however, that ER is almost always diffusely positive or completely negative (that is, it shows a bimodal staining pattern) with few cases falling between these extremes [49]. Given the small number of tumours likely to fall into the 1 to 9% of cells staining category, the impact of changing the cutoff to 1% on our results would be limited. Furthermore, there was no evidence of variation in the distributions of ER or PR status across the studies separately for *BRCA1* and *BRCA2* tumours (Mavaddat N, Antoniou AC, personal communication, manuscript in preparation) and all analyses were stratified by country. Finally, the clear differences observed for some SNPs (most notably for *FGFR2* rs2981582, where the association was limited to ER-positive disease) suggest that the effect of misclassification in tumour subtype on the SNP associations is likely to have been small.

BRCA1 and *BRCA2* tumours have also been found to differ in terms of other tumour characteristics compared to breast cancers in the general population. For example, tumours in mutation carriers are more likely to be of higher grade in comparison to breast cancers in the general population. The distribution of grade has been found to vary between ER-positive and ER-negative tumours in both *BRCA1* and *BRCA2* mutation carriers (Mavaddat N, Antoniou AC, personal communication, manuscript in preparation). Although the number of carriers with information on grade, ER status and SNPs was too small to permit combined analysis, our results are unlikely to have been influenced after adjusting for tumour grade. Case-only analysis to test for differences in associations between the SNPs and tumour grade (using ordinal logistic regression) revealed no significant associations between any of the SNPs and grade for both *BRCA1* and *BRCA2* mutation carriers ($P > 0.05$ for all tests, results not shown).

The analysis was performed within a retrospective cohort approach, by extending the retrospective likelihood approach described previously [36] to model the simultaneous effects on different breast cancer subtypes defined by ER/PR. Under this approach the associations were estimated simultaneously for the tumour subtypes under investigation. This method depends on the assumption that ascertainment of mutation carriers does not depend on tumour subtypes. This is a reasonable assumption since more than 90% of mutation carriers in our sample were recruited prior to 2007, when it was uncommon to use tumour pathology in selecting individuals for *BRCA1* and *BRCA2* mutation screening. Furthermore, the results were virtually identical in a case only, logistic regression analysis for testing for differences in the associations with tumour subtypes which included only individuals with known tumour characteristics (results not shown).

The average risks of developing ER-positive and ER-negative breast cancer in both *BRCA1* and *BRCA2* mutation carriers are substantially higher compared to the general population [38]. Therefore, in combination, these SNPs lead to much bigger differences in the absolute risk of developing the disease subtypes between the extremes of the combined SNP genotype distributions [50]. Based on the SNP profiles investigated in this report, the absolute risk difference between mutation carriers at the top 5% of the risk distribution compared to the bottom 5% is much greater for ER-positive breast cancer than for ER-negative breast cancer for both *BRCA1* and *BRCA2* (Figure 1). Recent GWAS have identified several other common breast cancer susceptibility variants which have not been investigated in *BRCA1* and *BRCA2* mutation carriers yet [24,51]. Moreover, ongoing GWAS in *BRCA1* and *BRCA2* mutation carriers [44,52] may also identify further modifiers of breast cancer risk for mutation carriers. It will be important to investigate the associations of these variants with different disease subtypes in *BRCA1* and *BRCA2* mutation carriers. Currently, it is unusual for the risks of different disease subtypes to be taken into account in the genetic counseling process. However, as more risk modifying variants are identified in the future, provided these have different associations with different disease subtypes in mutation carriers and confer relative risks which are greater (or smaller) than 1, having precise breast cancer subtype risks may be useful for the planning of the clinical management of both *BRCA1* and *BRCA2* mutation carriers. For example, knowing that a female *BRCA1* mutation carrier was primarily at risk of ER-positive breast cancer based on her associated SNP profile (rather than ER-negative breast cancer, as is the case for the majority of cases) might potentially

influence the choice of clinical management by screening, chemoprevention or prophylactic surgery.

Conclusions

In summary, in this report we investigated the associations of common breast cancer polymorphisms with ER and PR status. Our results indicate there are differential associations between these SNPs and the risk of developing ER-positive or ER-negative breast cancer in *BRCA1* and *BRCA2* mutation carriers that mirror similar differences seen in the general population. The findings add to our understanding of the biology of tumour development in mutation carriers and as more risk variants are identified in the future they may improve clinical management of these individuals.

Additional material

Additional file 1: Supplementary Tables. Supplementary Table 1: List of local ethics committees that approved studies. Supplementary Table 2: Methods and thresholds used to define the final marker variables. Supplementary Table 3: Per-allele hazard ratio estimates by progesterone receptor status for *BRCA1* mutation carriers. Supplementary Table 4: Per-allele hazard ratio estimates by progesterone receptor status for *BRCA2* mutation carriers.

Abbreviations

CMBA: Consortium of Investigators of Modifiers of *BRCA1/2*; ER: estrogen receptor; GWAS: genome-wide association studies; HR: hazard ratio; PR: progesterone receptor; SNPs: single nucleotide polymorphisms.

Acknowledgements

This work was supported by Cancer Research UK grants C12292/A11174 and C1287/A10118. The research leading to these results has received funding from the European Community's Seventh Framework Programme under grant agreement n° 223175 (HEALTH-F2-2009-223175). ACA is a CR-UK Senior Cancer Research Fellow, DFE is CR-UK Principal Research Fellow. Study specific acknowledgments

Breast Cancer Family Registry (BCFR)

This work was supported by the National Cancer Institute, National Institutes of Health under RFA-CA-06-503 and through cooperative agreements with members of the Breast Cancer Family Registry (BCFR) and Principal Investigators, including Cancer Care Ontario (U01 CA69467), Columbia University (U01 CA69398), Fox Chase Cancer Center (U01 CA69631), Huntsman Cancer Institute (U01 CA69446), Cancer Prevention Institute of California (formerly the Northern California Cancer Center) (U01 CA69417), University of Melbourne (U01 CA69638), and Research Triangle Institute Informatics Support Center (RFP No. N02PC45022-46). Samples from the FCCC, HCI, and CPIC were processed and distributed by the Coriell institute for Medical Research. The content of this manuscript does not necessarily reflect the views or policies of the National Cancer Institute or any of the collaborating centers in the BCFR, nor does mention of trade names, commercial products or organizations imply endorsement by the US Government or the BCFR.

The Baltic Familial Breast and Ovarian Cancer Consortium (BFOCC Latvia and Lithuania)

Lithuania: This work is supported by the Research Council of Lithuania grant LIG-19/2010 to Ramunas Janavicius. Latvia: We acknowledge Genome Database of Latvian Population, Latvian Biomedical Research and Study Center for providing data and DNA samples. This work is supported by Liepaja's municipal council to Laima Tihomirova (Latvian Biomedical Research and Study Centre).

Copenhagen Breast Cancer Study (CBCS)

We thank the NEYE Foundation for financial support
The Deutsches Krebsforschungszentrum (DKFZ) study
The study was supported by the DKFZ.
The CONSIT TEAM is supported by grants from Associazione Italiana per la Ricerca sul Cancro (4017) and by funds from Italian citizens who allocated the 5 × 1,000 share of their tax payment in support of the Fondazione IRCCS Istituto Nazionale Tumouri, according to Italian laws (INT-Institutional strategic projects "5 × 1000") to PP and grants from Fondazione Italiana per la Ricerca sul Cancro (Special Project "Hereditary tumours"), Ministero della Salute (Extraordinary National Cancer Program 2006 "Alleanza contro il Cancro", and "Progetto Tumouri Femminili") and Ministero dell'Università e Ricerca (RBLAO3-BETH) to PR. CONSIT TEAM acknowledges the contribution of Gaia Roversi, Carla B. Ripamonti, Marilena Morganti and Marco Pierotti of the of the Fondazione IRCCS Istituto Nazionale dei Tumouri, Milan, Italy; Monica Barile of the Istituto Europeo di Oncologia, Milan, Italy and Liliana Varesco of the Istituto Nazionale per la Ricerca sul Cancro, Genoa, Italy.
Epidemiological study of BRCA1 and BRCA2 mutation carriers (EMBRACE)
EMBRACE Collaborating Centers are: Coordinating Centre, Cambridge: Steve Ellis, Elena Fineberg, Radka Platte. North of Scotland Regional Genetics Service, Aberdeen: Zosia Miedzybrodzka. Northern Ireland Regional Genetics Service, Belfast: Patrick Morrison, Lisa Jeffers. West Midlands Regional Clinical Genetics Service, Birmingham: Trevor Cole, Kai-ren Ong, Jonathan Hoffman. South West Regional Genetics Service, Bristol: Margaret James. East Anglian Regional Genetics Service, Cambridge: Joan Paterson, Sarah Downing, Amy Taylor. Medical Genetics Services for Wales, Cardiff: T. Rogers., St James's Hospital, Dublin & National Centre for Medical Genetics, Dublin: M. John Kennedy, David Barton. South East of Scotland Regional Genetics Service, Edinburgh: Mary Porteous, Sarah Drummond. Peninsula Clinical Genetics Service, Exeter: Carole Brewer, Emma Kivuva, Anne Searle, Selina Goodman, Kathryn Hill. West of Scotland Regional Genetics Service, Glasgow: Victoria Murday, Nicola Bradshaw, Lesley Snadden, Mark Longmuir, Catherine Watt, Sarah Gibson, Eshika Haque, Ed Tobias, Alexis Duncan. South East Thames Regional Genetics Service, Guy's Hospital London: Chris Jacobs, Caroline Langman, Anna Whaithe. North West Thames Regional Genetics Service, Harrow. Leicestershire Clinical Genetics Service, Leicester. Yorkshire Regional Genetics Service, Leeds: Carol Chu, Julie Miller. Cheshire and Merseyside Clinical Genetics Service, Liverpool: Ian Ellis. Manchester Regional Genetics Service, Manchester: Jane Taylor. North East Thames Regional Genetics Service, NE Thames, London: Alison Male, Cheryl Berlin. Nottingham Centre for Medical Genetics, Nottingham: Rebecca Collier. Northern Clinical Genetics Service, Newcastle: Fiona Douglas, Oonagh Claber, Irene Jobson. Oxford Regional Genetics Service, Oxford: Lisa Walker, Diane McLeod, Dorothy Halliday, Sarah Durell, Barbara Stayner. The Institute of Cancer Research and Royal Marsden NHS Foundation Trust: Susan Shanley, Nazneen Rahman, Richard Houlston, Elizabeth Bancroft, Lucia D'Mello, Elizabeth Page, Audrey Ardern-Jones, Kelly Kohut, Jennifer Wiggins, Elena Castro, Anita Mitra, Lisa Robertson. North Trent Clinical Genetics Service, Sheffield: Oliver Quarrell, Cathryn Bardsley. South West Thames Regional Genetics Service, London: Shirley Hodgson, Sheila Goff, Glen Brice, Lizzie Winchester, Charlotte Eddy, Vishakha Tripathi, Virginia Attard. Wessex Clinical Genetics Service, Princess Anne Hospital, Southampton: Anneke Lucassen, Gillian Crawford, Donna McBride, Sarah Smalley. EMBRACE is supported by Cancer Research UK Grants C1287/A10118 and C1287/A11990. D. Gareth Evans and Fiona Laloo are supported by an NIHR grant to the Biomedical Research Centre, Manchester. The Investigators at The Institute of Cancer Research and The Royal Marsden NHS Foundation Trust are supported by an NIHR grant to the Biomedical Research Centre at The Institute of Cancer Research and The Royal Marsden NHS Foundation Trust. Ros Eeles, Elizabeth Bancroft and Lucia D'Mello are also supported by Cancer Research UK Grant C5047/A8385. Fox Chase Cancer Center Study (FCCC)
Fox Chase Cancer Center Biosample Repository: We acknowledge M. Pat Gilroy, Lesley Cruz, Diane Faison, Barbara Dettore, Mary Donovan, and Meghan Butler for their help collecting patient data and samples. University of Kansas Medical Center (KUMC):
A.K.G. was funded by U01CA69631, 5U01CA113916, and the Eileen Stein Jacoby Fund. The author acknowledges support from The University of Kansas Cancer Center and the Kansas Bioscience Authority Eminent Scholar Program. A.K.G. is the Chancellors Distinguished Chair in Biomedical Sciences endowed Professor.
Georgetown University

Work is supported by the Familial Cancer Registry (CI) and the Tissue Culture Shared Registry at Georgetown University (NIH/NCI grant P30-CA051008), the Cancer Genetics Network (HHSN261200744000C) (CI), and Swing Fore the Cure (CI).
The German Consortium of Hereditary Breast and Ovarian Cancer (GC-HBOC) GC-HBOC is supported by a grant of the German Cancer Aid (grant 109076), the Centre of Molecular Medicine Cologne (CMMC).
Genetic Modifiers of cancer risk in BRCA1/2 mutation carriers (GEMO) study: Cancer Genetics Network "Groupe Génétique et Cancer", Fédération Nationale des Centres de Lutte Contre le Cancer, France
The study was supported by the Ligue National Contre le Cancer; Association for International Cancer Research Grant (AICR-07-0454); and the Association "Le cancer du sein, parlons-en!" Award. We wish to thank all the GEMO collaborating groups for their contribution to this study. GEMO Collaborating Centers are: Coordinating Centres, Unité Mixte de Génétique Constitutionnelle des Cancers Fréquents, Centre Hospitalier Universitaire de Lyon/Centre Léon Bérard, et Equipe «Génétique du cancer du sein», Centre de Recherche en Cancérologie de Lyon: Laure Barjhoux, Carole Verny-Pierre, Sophie Giraud, and Service de Génétique Oncologique, Institut Curie, Paris: Marion Gauthier-Villars, Bruno Buecher, Claude Houdayer, Muriel Belotti, Carole Tirapo, Antoine de Pauw. Institut Gustave Roussy, Villejuif: Brigitte Bressac-de-Paillerets, Audrey Remenieras, Véronique Byrde, Olivier Caron, Gilbert Lenoir. Centre Jean Perrin, Clermont-Ferrand: Yves-Jean Bignon, Nancy Uhrhammer. Centre Léon Bérard, Lyon: Christine Lasset, Valérie Bonadona. Centre François Baclesse, Caen. Institut Paoli Calmettes, Marseille: Hagay Sobol, Violaine Bourdon, Tetsuro Noguchi, François Eisinger. Groupe Hospitalier Pitié-Salpêtrière, Paris: Florence Coulet, Chrystelle Colas, Florent Soubrier. CHU de Arnaud-de-Villeneuve, Montpellier. Centre Oscar Lambret, Lille: Jean-Philippe Peyrat, Joëlle Fournier, Françoise Révillon, Philippe Vennin, Claude Adenis. Hôpital René Huguenin/Institut Curie, St Cloud: Etienne Rouleau, Rosette Lidereau, Liliane Demange, Catherine Nogues. Centre Paul Strauss, Strasbourg. Institut Bergonié, Bordeaux: Emmanuelle Barouk-Simonet, Françoise Bonnet, Virginie Buben. Institut Claudius Regaud, Toulouse: Christine Toulas, Rosine Guimbaud, Laurence Gladieff, Viviane Feillel. CHU de Grenoble: Dominique Leroux, Hélène Dreyfus, Christine Rebischung, Magalie Peysselon. CHU de Dijon: Fanny Coron, Laurence Faivre. CHU de St-Etienne: Fabienne Prieur. Hôtel Dieu Centre Hospitalier, Chambéry: Sandra Fert Ferrer. Centre Antoine Lacassagne, Nice: Marc Fréney. CHU de Limoges: Laurence Vénat-Bouvet. CHU de Nantes: Capucine Delnatte. CHU Bretonneau, Tours. Creighton University, Omaha, USA: Henry T.Lynch, Carrie L.Snyder.
The Hereditary Breast and Ovarian Cancer Research Group Netherlands (HEBON)
HEBON Collaborating Centers: Coordinating center: Netherlands Cancer Institute, Amsterdam, NL: FBL Hogervorst, S Verhoef, M Verheus, LJ van 't Veer, FE van Leeuwen; Erasmus Medical Center, Rotterdam, NL: M Collée, AMW van den Ouweland, A Jager, MJ Hoening; Leiden University Medical Center, NL, Leiden: CJ van Asperen, JT Wijnen, MP Vreeswijk, RA Tollenaar, P Devilee; Radboud University Nijmegen Medical Center, Nijmegen, NL: MJ Ligtenberg, N Hoogerbrugge; University Medical Center Utrecht, Utrecht, NL: MG Ausems; Amsterdam Medical Center, NL: CM Aalfs; VU University Medical Center, Amsterdam, NL: JJP Gille, Q Waisfisz, HEJ Meijers-Heijboer; University Hospital Maastricht, Maastricht, NL: EB Gomez-Garcia, CE van Roozendaal, Marinus J Blok, B Caanen; University Medical Center Groningen University, NL: JC Oosterwijk, AH van der Hout, MJ Mourits; The Netherlands Foundation for the detection of hereditary tumours, Leiden, NL: HF Vasen. The HEBON study is supported by the Dutch Cancer Society grants NKI1998-1854, NKI2004-3088, NKI2007-3756 and the ZonMW grant 91109024.
Helsinki Breast Cancer Study (HEBCS)
HEBCS acknowledge Drs. Kirsimari Aaltonen, Carl Blomqvist and Päivi Heikkilä and research nurse Irja Erkkilä for their help with the patient data and samples. The HEBCS study has been financially supported by the Helsinki University Central Hospital Research Fund, Academy of Finland (132473), the Finnish Cancer Society, and the Sigrid Juselius Foundation.
Interdisciplinary Health Research International Team Breast Cancer Susceptibility (INHERIT BRCA)
We would like to thank Martine Tranchant for skilful technical assistance. J.S. is Chairholder of the Canada Research Chair in Oncogenetics. This work was supported by the Canadian Institutes of Health Research for the "CIHR Team in Familial Risks of Breast Cancer" program and by the Canadian Breast Cancer Research Alliance-grant #019511.

ILUH

The ILUH group was supported by the Icelandic Association "Walking for Breast Cancer Research" and by the Landspítali University Hospital Research Fund.

Istituto Oncologico Veneto Hereditary Breast and Ovarian Cancer Study (IOVHBOCS)

This study was supported by "Ministero della Salute" ("Progetto Tumori Femminili and grant numbers RFP5 2006-5-341353, ACC2/R6.9")

Kathleen Cuninghame Consortium for Research into Familial Breast Cancer (kConFab)

We wish to thank Heather Thorne, Eveline Niedermayr, all the kConFab research nurses and staff, the heads and staff of the Family Cancer Clinics, and the Clinical Follow Up Study (funded 2001-2009 by NHMRC and currently by the National Breast Cancer Foundation and Cancer Australia #628333) for their contributions to this resource, and the many families who contribute to kConFab. kConFab is supported by grants from the National Breast Cancer Foundation, the National Health and Medical Research Council (NHMRC) and by the Queensland Cancer Fund, the Cancer Councils of New South Wales, Victoria, Tasmania and South Australia, and the Cancer Foundation of Western Australia. ABS and GCT are NHMRC Fellows. Mayo Clinic Study

The MAYO study was supported by NIH grants CA116167, CA128978, a Specialized Program of Research Excellence (SPORE) in Breast Cancer (CA116201), and awards from the Komen Foundation for the Cure and the Breast Cancer Research Foundation.

Memorial Sloan-Kettering Cancer Center (MSKCC)

The study is supported by grant from the Breast Cancer Research Foundation, Starr Cancer Consortium, Norman and Carol Stone Genetic Research Fund, The Robert and Kate Niehaus Clinical Genetics Initiative at MSKCC

National Cancer Institute study (NCI)

The research of Drs. PL Mai and MH Greene was supported by the Intramural Research Program of the US National Cancer Institute, and by support services contracts NO2-CP-11019-50 and N02-CP-65504 with Westat, Inc., Rockville, MD.

N.N. Petrov Institute of Oncology

This work has been supported by the Russian Federation for Basic Research (grants 10-04-92601, 10-04-92110, 11-04-00227), the Federal Agency for Science and Innovations (contract 02.740.11.0780), the Commission of the European Communities (grant PITN-GA-2009-238132) and through a Royal Society International Joint grant (JP090615).

Ontario Cancer Genetics Network (OCGN)

We wish to thank Mona Gill, Lucine Collins, Nalan Gokgoz, Teresa Selander, Nayana Weerasooriya and members of the Ontario Cancer Genetics Network for their contributions to the study.

The Ohio State University Comprehensive Cancer Center (OSU-CCG)

This work was supported by the Ohio State University Comprehensive Cancer Center. We thank Kevin Sweet and Leigha Senter for accrual of study participants and data management. The Human Genetics Sample bank prepared DNA samples and the OSU Nucleic Acids Shared Resource assisted with genotyping.

Swedish BRCA1 and BRCA2 study (SWE-BRCA)

SWE-BRCA collaborators: Margareta Nordling, Annika Bergman and Zakaria Einbeigi, Gothenburg, Sahlgrenska University Hospital; Sigrun Liedgren, Linköping University Hospital; Åke Borg, Niklas Loman, Håkan Olsson, Ulf Kristoffersson, Helena Jernström, Katja Harbst, Lund University Hospital; Annika Lindblom, Annelie Liljegren, Gisela Barbany-Bustinza and Johanna Rantala, Stockholm, Karolinska University Hospital; Beatrice Melin, Henrik Grönberg, Eva-Lena Stattin and Monica Emanuelsson, Umeå University Hospital; Hans Ehrencrona, Richard Rosenquist and Niklas Dahl, Uppsala University Hospital.

University California San Francisco (UCSF)

The study received funding from the NIH, NCI Bay Area Breast Cancer SPORE (P50 CA 058207) and the Avon Foundation. We acknowledge support from the UCSF Helen Diller Family Comprehensive Cancer Center

UK and Gilda Radner Familial Ovarian Cancer Registries (UKGRFOCR)

UKFOCR was supported by a project grant from CRUK to Paul Pharoah. We thank Simon Gayther, Carole Pye, Patricia Harrington and Eva Wozniak for their contributions towards the UKFOCR. We'd like to acknowledge the Roswell Park Alliance Foundation for their continued support of the Gilda Radner Ovarian Family Cancer Registry. GRFOCR would like to acknowledge

Lara Sucheston (Department of Cancer Prevention and Control) and Kunle Odunsi (Departments Gynecologic Oncology and Immunology). University of Pennsylvania (UPENN)

Work is supported by grants from the Breast Cancer Research Foundation (to KLN), MacDonald Family Foundation (SMD) and Komen Foundation (SMD)

Author details

¹Department of Laboratory Medicine and Pathobiology, University of Toronto, Toronto, ON M5S 1A8, Canada; Department of Laboratory Medicine, and the Keenan Research Centre of the Li Ka Shing Knowledge Institute, St Michael's Hospital, 30 Bond Street, Toronto, ON M5B 1W8, Canada.

²Department of Laboratory Medicine and Pathology, Mayo Clinic, 200 First Street SW, Rochester, MN 55905, USA. ³Centre for Cancer Genetic Epidemiology, Department of Public Health and Primary Care, University of Cambridge, 2 Worts Causeway, Cambridge, CB1 8RN, UK. ⁴Department of Medicine, Abramson Cancer Center, Perelman School of Medicine at the University of Pennsylvania, 3400 Civic Center Boulevard, Philadelphia, PA 19104, USA. ⁵Faculty of Medicine, University of Southampton, University Hospital Southampton NHS Foundation Trust, Tremona Road, Southampton, SO16 6YD, UK. ⁶Department of Obstetrics and Gynecology, University of Helsinki and Helsinki University Central Hospital, Haartmaninkatu 8, 00290 Helsinki, Finland. ⁷Department of Preventive Medicine, Keck School of Medicine, University of Southern California, 2001 N Soto Street, Los Angeles, CA 90089-9237, USA. ⁸Department of Medicine, Memorial Sloan-Kettering Cancer Center and Weill Cornell Medical College, 1275 York Ave, New York, NY 10065, USA. ⁹National Cancer Institute, Division of Cancer Epidemiology and Genetics, Hormonal and Reproductive Epidemiology Branch, 6120 Executive Blvd., Rockville, MD 20852, USA. ¹⁰Queensland Institute of Medical Research, 300 Herston Rd, Herston, Brisbane, QLD 4006, Australia. ¹¹Centre of Familial Breast and Ovarian Cancer, Department of Gynaecology and Obstetrics and Centre for Integrated Oncology (CIO), University Hospital of Cologne, Kerpener Str. 62, Cologne, 50931, Germany. ¹²Unité Mixte de Génétique Constitutionnelle des Cancers Fréquents, Centre Hospitalier Universitaire de Lyon/Centre Léon Bérard, 28 rue Laënnec, Lyon 69373, France. ¹³INSERM U1052, CNRS UMR5286, Université Lyon 1, Cancer Research Center of Lyon, 28 rue Laënnec, Lyon 69373, France. ¹⁴Department of Molecular and Regenerative Medicine, Hematology, Oncology and Transfusion Medicine Center, Vilnius University Hospital Santariskiu Clinics, Santariskiu st 2, LT-08661 Vilnius and State Research Institute Innovative Medicine Center, Zygimantu st. 9, LT-01102 Vilnius, Lithuania. ¹⁵Center for Genomic Medicine, Rigshospitalet, Copenhagen University Hospital, Blegdamsvej 9, DK-2100 Copenhagen, Denmark. ¹⁶Department of Oncology, Rigshospitalet Bldg. 4262, Copenhagen University Hospital, Blegdamsvej 9, DK-2100 Copenhagen, Denmark. ¹⁷Human Genetics Group, Human Cancer Genetics Programme, Spanish National Cancer Research Centre, C/Melchor Fernández Almagro 3, Madrid, 28029, Spain and the Spanish Network on Rare Diseases (CIBERER). ¹⁸Institute of Biology and Molecular Genetics, Universidad de Valladolid (IBGM-UVA), C/Sanz y Forés, N° 3, Valladolid, 47003, Spain. ¹⁹Instituto de investigación sanitaria de Aragón (IIS), Hospital clinico Universitario "Lozano Blesa", San Juan Bosco 15, Zaragoza, 50009, Spain.

²⁰Molecular Diagnostics Laboratory, IRRP, National Center for Scientific Research Demokritos, Patriarchou Gregoriou E' & Neapoleos Str, Ag. Paraskevi 15310, Athens, Greece. ²¹Human Genetics Group and Genotyping Unit, Human Cancer Genetics Programme, Spanish National Cancer Research Centre, C/Melchor Fernández Almagro 3, Madrid, 28029, Spain and the Spanish Network on Rare Diseases (CIBERER). ²²Unit of Molecular Bases of Genetic Risk and Genetic Testing, Department of Preventive and Predictive Medicine, Fondazione IRCCS Istituto Nazionale Tumori (INT), via Giacomo Venezian 1, 20133 Milan, Italy; IFOM, Fondazione Istituto FIRC di Oncologia Molecolare, via Adamello 16, 20139 Milan, Italy. ²³Unit of Medical Genetics, Department of Preventive and Predictive Medicine, Fondazione IRCCS Istituto Nazionale Tumori (INT), via Giacomo Venezian 1, Milan, 20133, Italy. ²⁴Division of Cancer Prevention and Genetics, Istituto Europeo di Oncologia, via Ripamonti 435, Milan, 20141, Italy. ²⁵Unit of Experimental Oncology 1, Centro di Riferimento Oncologico, IRCCS, Aviano (PN), Italy. ²⁶Department of Genetics, Biology and Biochemistry, University of Turin, Turin, Italy. ²⁷Medical Genetics Unit, Department of Clinical Physiopathology, University of Florence, Firenze, Italy. ²⁸Department of Molecular Medicine, "Sapienza" University of Rome, Rome, Italy. ²⁹Division of Medical Oncology, Regina Elena Cancer Institute, Rome, Italy. ³⁰Department of Experimental Oncology,

Istituto Europeo di Oncologia, Milan, Italy; Consortium for Genomics Technology (Cogentech), Milan, Italy. ³¹Molecular Genetics of Breast Cancer, Deutsches Krebsforschungszentrum (DKFZ), Heidelberg, Germany. ³²Department of Epidemiology, Netherlands Cancer Institute, Amsterdam, The Netherlands. ³³Department of Clinical Genetics, VU Medical Center, Amsterdam, The Netherlands. ³⁴Department of Human Genetics and Department of Clinical Genetics, Leiden University Medical Center, Leiden, The Netherlands. ³⁵Department of Clinical Genetics and GROW, School for Oncology and Developmental Biology, MUMC, Maastricht, The Netherlands. ³⁶Department of Human Genetics 849, Radboud University Nijmegen Medical Centre, P.O. BOX 9101, 6500 HB Nijmegen, The Netherlands. ³⁷Department of Human Genetics 836, Radboud University Nijmegen Medical Centre, P.O. BOX 9101, 6500 HB Nijmegen, The Netherlands. ³⁸Department of Medical Oncology, Family Cancer Clinic, Erasmus University Medical Center, Rotterdam, The Netherlands. ³⁹Department of Surgical Oncology, Family Cancer Clinic, Erasmus University Medical Center, Rotterdam, The Netherlands. ⁴⁰Department of Medical Genetics, University Medical Center Utrecht, The Netherlands. ⁴¹Department of Clinical Genetics, Academic Medical Center, Amsterdam, The Netherlands. ⁴²Barts Cancer Institute, Queen Mary University of London, Centre for Tumour Biology, Charterhouse Square, London, UK. ⁴³Genetic Medicine, Manchester Academic Health Sciences Centre, Central Manchester University Hospitals NHS Foundation Trust, Manchester, UK. ⁴⁴Oncogenetics Team, The Institute of Cancer Research and Royal Marsden NHS Foundation Trust, UK. ⁴⁵Clinical Genetics, Guy's and St. Thomas' NHS Foundation Trust, London, UK. ⁴⁶Yorkshire Regional Genetics Service, Leeds, UK. ⁴⁷Ferguson-Smith Centre for Clinical Genetics, Yorkhill Hospitals, Glasgow, UK. ⁴⁸Sheffield Clinical Genetics Service, Sheffield Children's Hospital, Sheffield, UK. ⁴⁹Clinical Genetics Department, St Michael's Hospital, Bristol, UK. ⁵⁰North West Thames Regional Genetics Service, Kennedy-Galton Centre, Harrow, UK. ⁵¹North of Scotland Regional Genetics Service, NHS Grampian and University of Aberdeen, Foresterhill, Aberdeen, UK. ⁵²Nottingham Clinical Genetics Service, Nottingham University Hospitals NHS Trust, UK. ⁵³Cheshire and Merseyside Clinical Genetics Service, Liverpool Women's NHS Foundation Trust, Liverpool, UK. ⁵⁴Leicestershire Clinical Genetics Service, University Hospitals of Leicester NHS Trust, UK. ⁵⁵North East Thames Regional Genetics Service, Great Ormond Street Hospital for Children NHS Trust, London, UK. ⁵⁶All Wales Medical Genetics Service, Glan Clwyd Hospital, Rhyl, UK. ⁵⁷All Wales Medical Genetics Services, Singleton Hospital, Swansea, UK. ⁵⁸Department of Pathology and Laboratory Medicine, University of Kansas Medical Center, Kansas City, KS, USA. ⁵⁹Institute for Medical Informatics, Statistics and Epidemiology, University of Leipzig, Leipzig, Germany. ⁶⁰Department of Gynaecology and Obstetrics, Division of Tumour Genetics, Klinikum rechts der Isar, Technical University, Munich, Germany. ⁶¹Department of Gynaecology and Obstetrics, Ludwig-Maximilians University, Munich, Germany. ⁶²Department of Gynaecology and Obstetrics, University Hospital of Schleswig-Holstein (UKSH), Campus Kiel, Christian-Albrechts University, Kiel, Germany. ⁶³Department of Gynaecology and Obstetrics, University Hospital Düsseldorf, Heinrich-Heine University, Düsseldorf, Germany. ⁶⁴Institute of Human Genetics, Department of Human Genetics, Heidelberg University Hospital, Heidelberg, Germany. ⁶⁵Department of Gynaecology and Obstetrics, University Hospital, Ulm, Germany. ⁶⁶Institute of Cell and Molecular Pathology, Hannover Medical School, Hannover, Germany. ⁶⁷Department of Gynaecology and Obstetrics, University Hospital Carl Gustav Carus, Technical University, Dresden, Germany. ⁶⁸Institute of Human Genetics, University of Münster, Münster, Germany. ⁶⁹Institute of Human Genetics, Campus Virchow Klinikum, Charité Berlin, Germany. ⁷⁰Centre of Familial Breast and Ovarian Cancer, Department of Medical Genetics, Institute of Human Genetics, University Würzburg, Germany. ⁷¹Institute of Human Genetics, University Regensburg, Germany. ⁷²Institute of Human Genetics, University Leipzig, Germany. ⁷³Institute of Human Genetics, University Hospital, Frankfurt a.M., Germany. ⁷⁴Breakthrough Breast Cancer Research Centre, Institute of Cancer Research, UK. ⁷⁵Service de Génétique Oncologique, Institut Curie, Paris, France. ⁷⁶Genetic Unit, Avicenne Hospital, Assistance Publique-Hôpitaux de Paris, Paris, France; Sud-Francilien Hospital, Evry-Corbeil, France; University Hospital, Clermont-Ferrand, France. ⁷⁷Centre François Baclesse, Caen, France. ⁷⁸Unité d'Oncogénétique, CLCC Paul Strauss, Strasbourg, France. ⁷⁹Service de Génétique, Centre Hospitalier Universitaire Bretonneau, Tours, France. ⁸⁰Unité d'Oncogénétique, CHU Arnaud de Villeneuve, Montpellier, France. ⁸¹Service de Génétique Clinique Chromosomique et Moléculaire, Centre Hospitalier Universitaire de St Etienne, St Etienne, France. ⁸²Cancer Genetics Unit, INSERM U916, Institut Bergonié, Université de Bordeaux, Bordeaux, France. ⁸³Unité INSERM U830, Institut Curie, Paris, France; Université Paris Descartes, Faculté de Médecine, Paris, France. ⁸⁴Lombardi Comprehensive Cancer Center, Georgetown University, Washington DC, USA. ⁸⁵Molecular Oncology Laboratory, Hospital Clinico San Carlos, Madrid, Spain. ⁸⁶Department of Clinical Genetics, Helsinki University Central Hospital, Meilahdentie 2, 00290 Helsinki, Finland. ⁸⁷Hereditary Cancer Program, Institut Català d'Oncologia, Hospital Duran i Reynals - Bellvitge Biomedical Research Institute (IDIBELL), L'Hospitalet de Llobregat, Barcelona, Spain. ⁸⁸Department of Pathology, Landspítali University Hospital, Reykjavik, Iceland, Faculty of Medicine, University of Iceland, Reykjavik, Iceland. ⁸⁹Cancer Genomics Laboratory, Centre Hospitalier Universitaire de Québec, 2705 Laurier Boulevard, T3-57, Quebec City, QC, Canada. ⁹⁰Cancer Genomics Laboratory, Centre Hospitalier Universitaire de Québec, 2705 Laurier Boulevard, T3-57, Quebec City, QC, Canada; Canada Research Chair in Oncogenetics, Department of Molecular Medicine, Faculty of Medicine, Laval University, QC, Canada. ⁹¹Immunology and Molecular Oncology Unit, Istituto Oncologico Veneto IOV - IRCCS, Via Gattamelata 64, 35128 Padua, Italy. ⁹²Department of Oncology and Surgical Sciences, University of Padua and Istituto Oncologico Veneto IOV - IRCCS, Via Gattamelata 64, 35128 Padua, Italy. ⁹³Peter MacCallum Cancer Centre, Melbourne, VIC 3052, Australia. ⁹⁴Department of Anatomical Pathology, Prince of Wales Hospital, Randwick, NSW 2031, Australia. ⁹⁵Abramson Cancer Center and University of Pennsylvania Perelman School of Medicine, Philadelphia, PA, USA. ⁹⁶University of Chicago, Chicago, IL, USA. ⁹⁷City of Hope Comprehensive Cancer Center and Department of Population Sciences, Beckman Research Institute, City of Hope, Duarte, CA, USA. ⁹⁸Departments of Medicine, and Preventive Medicine and Public Health, Creighton University, Omaha, NE, USA. ⁹⁹Jonsson Comprehensive Cancer Center at the University of California, Los Angeles, CA, USA. ¹⁰⁰Department of Internal Medicine and Harold C. Simmons Comprehensive Cancer Center, University of Texas, Southwestern Medical Center, Dallas, TX; USA; Department of Pediatrics, University of Texas Health Science Center at San Antonio, San Antonio, TX, USA. ¹⁰¹Department of Health Sciences Research, Mayo Clinic, Rochester, MN, USA. ¹⁰²Department of Medical Genetics, Mayo Clinic, Rochester, MN, USA. ¹⁰³University of Delaware, Newark, DE, USA. ¹⁰⁴Epidemiology Research Program, American Cancer Society, Atlanta, GA, USA. ¹⁰⁵Department of Obstetrics/Gynaecology and Comprehensive Cancer Center, Medical University of Vienna, Vienna, Austria. ¹⁰⁶Clinical Genetics Branch, Division of Cancer Epidemiology and Genetics, US National Cancer Institute, Rockville, MD, USA. ¹⁰⁷Laboratory of Molecular Oncology, N.N. Petrov Institute of Oncology, St.-Petersburg, Russia. ¹⁰⁸Department of Laboratory Medicine and Pathobiology, University of Toronto, Toronto, ON, Canada; Department of Laboratory Medicine, and the Keenan Research Centre of the Li Ka Shing Knowledge Institute, St Michael's Hospital, Toronto, ON, Canada. ¹⁰⁹Samuel Lunenfeld Research Institute, Mount Sinai Hospital, Toronto; Department of Laboratory Medicine and Pathobiology, University of Toronto, ON, Canada. ¹¹⁰Cancer Care Ontario, ON, Canada. ¹¹¹Departments of Molecular Virology, Immunology and Medical Genetics and Internal Medicine, The Ohio State University Comprehensive Cancer Center, Columbus, OH, USA. ¹¹²Clinical Genetics, Rigshospitalet, Copenhagen University, Copenhagen, Denmark. ¹¹³Clinical Genetics, Odense University Hospital, Odense, Denmark. ¹¹⁴Department of Clinical Genetics, Aarhus University Hospital, Aarhus, Denmark. ¹¹⁵Clinical Genetics, Vejle Hospital, Denmark. ¹¹⁶Section of Genetic Oncology, Dept. of Laboratory Medicine, University and University Hospital of Pisa, Pisa, Italy. ¹¹⁷Department of Clinical Genetics, Lund University Hospital, Lund, Sweden. ¹¹⁸Oncological Centre, Lund University Hospital, Lund, Sweden. ¹¹⁹Department of Oncology, Karolinska University Hospital, Stockholm, Sweden. ¹²⁰Division of Clinical Genetics, Department of Clinical and Experimental Medicine, Linköping University, Linköping, Sweden. ¹²¹Department of Oncology, Sahlgrenska University Hospital, Gothenburg, Sweden. ¹²²Department of Population Sciences, the Beckman Research Institute of the City of Hope, Duarte, CA, USA. ¹²³UCSF Cancer Risk Program and Departments of Medicine and Epidemiology and Biostatistics, University of California San Francisco, San Francisco, CA, USA. ¹²⁴Department of Oncology, University of Cambridge, Cambridge, UK. ¹²⁵Department of Cancer Prevention and Control, Roswell Park Cancer Institute, Buffalo, NY, USA. ¹²⁶Women's Cancer Program at the Samuel Oschin Comprehensive Cancer Institute, Cedars-Sinai Medical Center, Los Angeles, CA, USA. ¹²⁷Department of Epidemiology, Cancer Prevention Institute of California, Fremont, CA, USA. ¹²⁸Fox Chase Cancer Center,

Philadelphia, PA, USA. ¹²⁹Department of Oncological Sciences, Huntsman Cancer Institute, University of Utah, Salt Lake City, UT, USA. ¹³⁰Genetic Epidemiology Laboratory, Department of Pathology, University of Melbourne, Melbourne, VIC, Australia. ¹³¹Centre for Molecular, Environmental, Genetic and Analytic Epidemiology, University of Melbourne, Melbourne, VIC, Australia. ¹³²Department of Epidemiology, Columbia University, New York, NY, USA. ¹³³Department of Cancer Biology, Dana-Farber Cancer Institute, Boston, MA, USA. ¹³⁴Department of Dermatology, University of Utah School of Medicine, Salt Lake City, UT, USA. ¹³⁵Samuel Lunenfeld Research Institute, Mount Sinai Hospital, 600 University Avenue, Toronto, ON, M5G 1X5, Canada; Cancer Care Ontario, Departments of Molecular Genetics and Laboratory Medicine and Pathobiology, University of Toronto, ON, Canada. ¹³⁶Breast Cancer Family Registry. ¹³⁷Cancer Genetics Network "Groupe Génétique et Cancer", Fédération Nationale des Centres de Lutte Contre le Cancer. ¹³⁸Netherlands Cancer Institute, Amsterdam, The Netherlands. ¹³⁹Ontario Cancer Genetics Network, Cancer Care Ontario, 620 University Avenue, Toronto, ON M5G 2L7, Canada. ¹⁴⁰Karolinska Institute, Stockholm, Sweden.

Authors' contributions

AMM, ILA, ACA drafted the initial manuscript. ACA developed the analytical methods and performed the statistical analysis. AMM, ACA, ILA, FJC, DB, SMD, DE, HN, SJR, MR, MS, ABS and BW are members of the CIMBA pathology working group and participated in the design of the study. LM and DB are the CIMBA database managers. AL wrote computer programs for the analysis. SH and OMS reviewed, recoded and classified the BRCA1 and BRCA2 mutations in CIMBA. GCT initiated and coordinated CIMBA. DFE participated in the study design and advised on the statistical analysis. RJ, TvOH, FCNi, BE, AO, IMR, MD, JG, MP, JB, PP, SM, BP, DZ, EC, BB, AV, BP, LP, LO, AS, LB, PR, UH, MV, HEJMH, JW, EBGG, MRN, MK, CS, MMATL, RBvdL, TvO, MR, DF, JLI, DGE, FL, RE, LI, JA, RD, JC, AD, HD, HG, JE, CH, JB, LES, E McC, AM, SP, AG, RKS, KR, CE, AM, IR, NA, DN, CS, HD, DG, KK, SPA, RVM, IS, BF, WH, DS, HG, VCM, FCFC, SM, ML, NBK, AH, PB, DM, JPF, IM, PP, IC, ML, CK, ML, NS, DSL, CI, TC, MH, TH, KA,IB, CL, RBB, PS, MD, JS, MM, ST, EDA, SF, MY, TR, OO, JNW, HTL, PAG, GET, XW, ZF, VSP, NML, CS, KO, RS, MG, JB, NK, CFS, MKT, DGK, AFR, PLM, MHG, EI, FPO, HO, GG, AET, AMG, MT, TAK, UBJ, ABS, MAC, MS, KH, AvW, BA, MSA, PK, YCD, SLN, MB, PDPP, KBM, KLN, BYK, JG, EMJ, MJD, SMB, MCS, JLH, MBT, WC, AFM and DG acquired phenotypic data and DNA samples or performed SNP genotyping. All authors read and approved the final manuscript.

Competing interests

The authors declare that they have no competing interests.

Received: 27 April 2011 Accepted: 2 November 2011

Published: 2 November 2011

References

1. Ford D, Easton DF, Bishop DT, Narod SA, Goldgar DE: **Risks of cancer in BRCA1-mutation carriers. Breast Cancer Linkage Consortium.** *Lancet* 1994, **343**:692-695.
2. The Breast Cancer Linkage Consortium: **Cancer risks in BRCA2 mutation carriers.** *J Natl Cancer Inst* 1999, **91**:1310-1316.
3. Thompson D, Easton DF: **Cancer incidence in BRCA1 mutation carriers.** *J Natl Cancer Inst* 2002, **94**:1358-1365.
4. Mavaddat N, Pharoah PD, Blows F, Driver KE, Provenzano E, Thompson D, Macinnis RJ, Shah M, Easton DF, Antoniou AC: **Familial relative risks for breast cancer by pathological subtype: a population-based cohort study.** *Breast Cancer Res* 2010, **12**:R10.
5. Thompson D, Easton D: **The genetic epidemiology of breast cancer genes.** *J Mammary Gland Biol Neoplasia* 2004, **9**:221-236.
6. Antoniou A, Pharoah PD, Narod S, Risch HA, Eyfjord JE, Hopper JL, Loman N, Olsson H, Johannsson O, Borg A, Pasini B, Radice P, Manoukian S, Eccles DM, Tang N, Olah E, Anton-Culver H, Warner E, Lubinski J, Gronwald J, Gorski B, Tulinius H, Thorlacius S, Eerola H, Nevanlinna H, Syrjäkoski K, Kallioniemi OP, Thompson D, Evans C, Peto J, et al: **Average risks of breast and ovarian cancer associated with BRCA1 or BRCA2 mutations detected in case Series unselected for family history: a combined analysis of 22 studies.** *Am J Hum Genet* 2003, **72**:1117-1130.
7. Begg CB, Haile RW, Borg A, Malone KE, Concannon P, Thomas DC, Langholz B, Bernstein L, Olsen JH, Lynch CF, Anton-Culver H, Capanu M, Liang X, Hummer AJ, Sima C, Bernstein JL: **Variation of breast cancer risk among BRCA1/2 carriers.** *JAMA* 2008, **299**:194-201.
8. Ford D, Easton DF, Stratton M, Narod S, Goldgar D, Devilee P, Bishop DT, Weber B, Lenoir G, Chang-Claude J, Sobol H, Teare MD, Struwing J, Arason A, Scherneck S, Peto J, Rebbeck TR, Tonin P, Neuhausen S, Barkardottir R, Eyfjord J, Lynch H, Ponder BA, Gayther SA, Zelada-Hedman M, et al: **Genetic heterogeneity and penetrance analysis of the BRCA1 and BRCA2 genes in breast cancer families. The Breast Cancer Linkage Consortium.** *Am J Hum Genet* 1998, **62**:676-689.
9. Hopper JL, Southey MC, Dite GS, Jolley DJ, Giles GG, McCredie MR, Easton DF, Venter DJ: **Population-based estimate of the average age-specific cumulative risk of breast cancer for a defined set of protein-truncating mutations in BRCA1 and BRCA2. Australian Breast Cancer Family Study.** *Cancer Epidemiol Biomarkers Prev* 1999, **8**:741-747.
10. Milne RL, Osorio A, Cajal TR, Vega A, Llorc G, de la Hoya M, Diez O, Alonso MC, Lazaro C, Blanco I, Sánchez-de-Abajo A, Caldés T, Blanco A, Graña B, Durán M, Velasco E, Chirivella I, Cardeñosa EE, Tejada MI, Beristain E, Miramar MD, Calvo MT, Martínez E, Guillén C, Salazar R, San Román C, Antoniou AC, Orioste M, Benítez J: **The average cumulative risks of breast and ovarian cancer for carriers of mutations in BRCA1 and BRCA2 attending genetic counseling units in Spain.** *Clin Cancer Res* 2008, **14**:2861-2869.
11. Simchoni S, Friedman E, Kaufman B, Gershoni-Baruch R, Orr-Urtreger A, Kedari-Barnes I, Shiri-Sverdllov R, Dagan E, Tsabari S, Shohat M, Catane R, King MC, Lahad A, Levy-Lahad E: **Familial clustering of site-specific cancer risks associated with BRCA1 and BRCA2 mutations in the Ashkenazi Jewish population.** *Proc Natl Acad Sci USA* 2006, **103**:3770-3774.
12. Struwing JP, Hartge P, Wacholder S, Baker SM, Berlin M, McAdams M, Timmerman MM, Brody LC, Tucker MA: **The risk of cancer associated with specific mutations of BRCA1 and BRCA2 among Ashkenazi Jews.** *N Engl J Med* 1997, **336**:1401-1408.
13. Sorlie T, Tibshirani R, Parker J, Hastie T, Marron JS, Nobel A, Deng S, Johnsen H, Pesich R, Geisler S, Demeter J, Perou CM, Lønning PE, Brown PO, Borresen-Dale AL, Botstein D: **Repeated observation of breast tumour subtypes in independent gene expression data sets.** *Proc Natl Acad Sci USA* 2003, **100**:8418-8423.
14. van 't Veer LJ, Dai H, van de Vijver MJ, He YD, Hart AA, Mao M, Peterse HL, van der Kooy K, Marton MJ, Witteveen AT, Schreiber GJ, Kerkhoven RM, Roberts C, Linsley PS, Bernards R, Friend SH: **Gene expression profiling predicts clinical outcome of breast cancer.** *Nature* 2002, **415**:530-536.
15. Lakhani SR, Jacquemier J, Sloane JP, Gusterson BA, Anderson TJ, van de Vijver MJ, Farid LM, Venter D, Antoniou A, Storfer-Isser A, Smyth E, Steel CM, Haites N, Scott RJ, Goldgar D, Neuhausen S, Daly PA, Ormiston W, McManus R, Scherneck S, Ponder BA, Ford D, Peto J, Stoppa-Lyonnet D, Bignon YJ, Struwing JP, Spurr NK, Bishop DT, Klijn JG, Devilee P, et al: **Multifactorial analysis of differences between sporadic breast cancers and cancers involving BRCA1 and BRCA2 mutations.** *J Natl Cancer Inst* 1998, **90**:1138-1145.
16. Lakhani SR, Reis-Filho JS, Fulford L, Penault-Llorca F, van der Vijver M, Parry S, Bishop T, Benitez J, Rivas C, Bignon YJ, Chang-Claude J, Hamann U, Cornelisse CJ, Devilee P, Beckmann MW, Nestle-Krämling C, Daly PA, Haites N, Varley J, Lalloo F, Evans G, Maugard C, Meijers-Heijboer H, Klijn JG, Olah E, Gusterson BA, Piloni S, Radice P, Scherneck S, Sobol H, et al: **Prediction of BRCA1 status in patients with breast cancer using estrogen receptor and basal phenotype.** *Clin Cancer Res* 2005, **11**:5175-5180.
17. Bane AL, Beck JC, Bleiweiss I, Buys SS, Catalano E, Daly MB, Giles G, Godwin AK, Hibshoosh H, Hopper JL, John EM, Layfield L, Longacre T, Miron A, Senie R, Southey MC, West DW, Whittmore AS, Wu H, Andrulis IL, O'Malley FP: **BRCA2 mutation-associated breast cancers exhibit a distinguishing phenotype based on morphology and molecular profiles from tissue microarrays.** *Am J Surg Pathol* 2007, **31**:121-128.
18. Ahmed S, Thomas G, Ghoussaini M, Healey CS, Humphreys MK, Platte R, Morrison J, Maranian M, Pooley KA, Luben R, Eccles D, Evans DG, Fletcher O, Johnson N, dos Santos Silva I, Peto J, Stratton MR, Rahman N, Jacobs K, Prentice R, Anderson GL, Rajkovic A, Curb JD, Ziegler RG, Berg CD, Buys SS, McCarty CA, Feigelson HS, Calle EE, Thun MJ, et al: **Newly discovered breast cancer susceptibility loci on 3p24 and 17q23.2.** *Nat Genet* 2009, **41**:585-590.
19. Easton DF, Pooley KA, Dunning AM, Pharoah PD, Thompson D, Ballinger DG, Struwing JP, Morrison J, Field H, Luben R, Wareham N, Ahmed S, Healey CS, Bowman R, SEARCH collaborators, Meyer KB,

- Haiman CA, Kolonel LK, Henderson BE, Le Marchand L, Brennan P, Sangrajrang S, Gaborieau V, Odefrey F, Shen CY, Wu PE, Wang HC, Eccles D, Evans DG, Peto J, *et al*: **Genome-wide association study identifies novel breast cancer susceptibility loci.** *Nature* 2007, **447**:1087-1093.
20. Hunter DJ, Kraft P, Jacobs KB, Cox DG, Yeager M, Hankinson SE, Wacholder S, Wang Z, Welch R, Hutchinson A, Wang J, Yu K, Chatterjee N, Orr N, Willett WC, Colditz GA, Ziegler RG, Berg CD, Buys SS, McCarty CA, Feigelson HS, Calle EE, Thun MJ, Hayes RB, Tucker M, Gerhard DS, Fraumeni JF Jr, Hoover RN, Thomas G, Chanock SJ: **A genome-wide association study identifies alleles in FGFR2 associated with risk of sporadic postmenopausal breast cancer.** *Nat Genet* 2007, **39**:870-874.
21. Stacey SN, Manolescu A, Sulem P, Rafnar T, Gudmundsson J, Gudjonsson SA, Masson G, Jakobsdottir M, Thorlacius S, Helgason A, Aben KK, Strobbe LJ, Albers-Akkers MT, Swinkels DW, Henderson BE, Kolonel LN, Le Marchand L, Millastre E, Andres R, Godino J, Garcia-Prats MD, Polo E, Tres A, Mouty M, Saemundsdottir J, Backman VM, Gudmundsson L, Kristjansson K, Bergthorsson JT, Kostic J, *et al*: **Common variants on chromosomes 2q35 and 16q12 confer susceptibility to estrogen receptor-positive breast cancer.** *Nat Genet* 2007, **39**:865-869.
22. Stacey SN, Manolescu A, Sulem P, Thorlacius S, Gudjonsson SA, Jonsson GF, Jakobsdottir M, Bergthorsson JT, Gudmundsson J, Aben KK, Strobbe LJ, Swinkels DW, van Engelenburg KC, Henderson BE, Kolonel LN, Le Marchand L, Millastre E, Andres R, Saez B, Lambea J, Godino J, Polo E, Tres A, Picelli S, Rantala J, Margolin S, Jonsson T, Sigurdsson H, Jonsdottir T, Hrafnkelsson J, *et al*: **Common variants on chromosome 5p12 confer susceptibility to estrogen receptor-positive breast cancer.** *Nat Genet* 2008, **40**:703-706.
23. Thomas G, Jacobs KB, Kraft P, Yeager M, Wacholder S, Cox DG, Hankinson SE, Hutchinson A, Wang Z, Yu K, Chatterjee N, Garcia-Closas M, Gonzalez-Bosquet J, Prokunina-Olsson L, Orr N, Willett WC, Colditz GA, Ziegler RG, Berg CD, Buys SS, McCarty CA, Feigelson HS, Calle EE, Thun MJ, Diver R, Prentice R, Jackson R, Kooperberg C, Chlebowski R, Lissowska J, *et al*: **A multistage genome-wide association study in breast cancer identifies two new risk alleles at 1p11.2 and 14q24.1 (RAD51L1).** *Nat Genet* 2009, **41**:579-584.
24. Turnbull C, Ahmed S, Morrison J, Pernet D, Renwick A, Maranian M, Seal S, Ghoussaini M, Hines S, Healey CS, Hughes D, Warren-Perry M, Tapper W, Eccles D, Evans DG, Breast Cancer Susceptibility Collaboration (UK), Hooping M, Schutte M, van den Ouweland A, Houlston R, Ross G, Langford C, Pharoah PD, Stratton MR, Dunning AM, Rahman N, Easton DF: **Genome-wide association study identifies five new breast cancer susceptibility loci.** *Nat Genet* 2010, **42**:504-507.
25. Zeger SL, Liang KY, Albert PS: **Models for longitudinal data: a generalized estimating equation approach.** *Biometrics* 1988, **44**:1049-1060.
26. Broeks A, Schmidt MK, Sherman ME, Couch FJ, Hopper JL, Dite GS, Apicella C, Smith LD, Hammet F, Southey MC, Van 't Veer LJ, de Groot R, Smit VT, Fasching PA, Beckmann MW, Jud S, Ekici AB, Hartmann A, Hein A, Schulz-Wendtland R, Birkwinkel B, Marime F, Schneeweiss A, Sinn HP, Sohn C, Tchatchou S, Bojesen SE, Nordestgaard BG, Flyger H, Ørsted DD, *et al*: **Low penetrance breast cancer susceptibility loci are associated with specific breast tumour subtypes: findings from the Breast Cancer Association Consortium.** *Hum Mol Genet* 2011, **20**:3289-3303.
27. Garcia-Closas M, Hall P, Nevanlinna H, Pooley K, Morrison J, Richesson DA, Bojesen SE, Nordestgaard BG, Axelsson CK, Arias JI, Milne RL, Ribas G, González-Neira A, Benítez J, Zamora P, Brauch H, Justenhoven C, Hamann U, Ko YD, Bruening T, Haas S, Dörk T, Schürmann P, Hillemanns P, Bogdanova N, Bremer M, Karstens JH, Fagerholm R, Aaltonen K, Aittomäki K, *et al*: **Heterogeneity of breast cancer associations with five susceptibility loci by clinical and pathological characteristics.** *PLoS Genet* 2008, **4**: e1000054.
28. Antoniou AC, Kartsonaki C, Sinilnikova O, Soucy P, McGuffog L, Healey S, Lee A, Peterlongo P, Manoukian S, Peissel B, Zaffaroni D, Cattaneo E, Barile M, Pensotti V, Pasini B, Dolcetti R, Giannini G, Putignano AL, Varesco L, Radice P, Mai PL, Greene MH, Andrulis IL, Glendon G, Ozcelik H, Thomassen M, Gerdes AM, Kruse TA, Birk Jensen U, Crüger DG, *et al*: **Common alleles at 6q25.1 and 1p11.2 are associated with breast cancer risk for BRCA1 and BRCA2 mutation carriers.** *Hum Mol Genet* 2011, **20**:3304-3321.
29. Antoniou AC, Spurdle AB, Sinilnikova OM, Healey S, Pooley KA, Schmutzler RK, Versmold B, Engel C, Meindl A, Arnold N, Hofmann W, Sutter C, Niederacher D, Deissler H, Caldes T, Kämpjärvi K, Nevanlinna H, Simard J, Beesley J, Chen X, Kathleen Cuninghams Consortium for Research into Familial Breast Cancer, Neuhausen SL, Rebbeck TR, Wagner T, Lynch HT, Isaacs C, Weitzel J, Ganz PA, Daly MB, Tomlinson G, *et al*: **Common breast cancer-predisposition alleles are associated with breast cancer risk in BRCA1 and BRCA2 mutation carriers.** *Am J Hum Genet* 2008, **82**:937-948.
30. Antoniou AC, Sinilnikova OM, McGuffog L, Healey S, Nevanlinna H, Heikkinen T, Simard J, Spurdle AB, Beesley J, Chen X, Kathleen Cuninghams Foundation Consortium for Research into Familial Breast Cancer, Neuhausen SL, Ding YC, Couch FJ, Wang X, Fredericksen Z, Peterlongo P, Peissel B, Bonanni B, Viel A, Bernard L, Radice P, Szabo CI, Foretova L, Zikan M, Claes K, Greene MH, Mai PL, Rennett G, Lejbkowitz F, *et al*: **Common variants in LSP1, 2q35 and 8q24 and breast cancer risk for BRCA1 and BRCA2 mutation carriers.** *Hum Mol Genet* 2009, **18**:4442-4456.
31. Antoniou AC, Beesley J, McGuffog L, Sinilnikova OM, Healey S, Neuhausen SL, Ding YC, Rebbeck TR, Weitzel JN, Lynch HT, Isaacs C, Ganz PA, Tomlinson G, Olopade OI, Couch FJ, Wang X, Lindor NM, Pankratz VS, Radice P, Manoukian S, Peissel B, Zaffaroni D, Barile M, Viel A, Allavena A, Dall'Olio V, Peterlongo P, Szabo CI, Zikan M, Claes K, *et al*: **Common breast cancer susceptibility alleles and the risk of breast cancer for BRCA1 and BRCA2 mutation carriers: implications for risk prediction.** *Cancer Res* 2010, **70**:9742-9754.
32. Reeves GK, Travis RC, Green J, Bull D, Tipper S, Baker K, Beral V, Peto R, Bell J, Zelenika D, Lathrop M, Million Women Study Collaborators: **Incidence of breast cancer and its subtypes in relation to individual and multiple low-penetrance genetic susceptibility loci.** *JAMA* 2010, **304**:426-434.
33. Chenevix-Trench G, Milne RL, Antoniou AC, Couch FJ, Easton DF, Goldgar DE: **An international initiative to identify genetic modifiers of cancer risk in BRCA1 and BRCA2 mutation carriers: the Consortium of Investigators of Modifiers of BRCA1 and BRCA2 (CIMBA).** *Breast Cancer Res* 2007, **9**:104.
34. 2011 [http://www.srl.cam.ac.uk/consortia/cimba/eligibility/eligibility.html].
35. Antoniou AC, Goldgar DE, Andrieu N, Chang-Claude J, Brohet R, Rookus MA, Easton DF: **A weighted cohort approach for analysing factors modifying disease risks in carriers of high-risk susceptibility genes.** *Genet Epidemiol* 2005, **29**:1-11.
36. Antoniou AC, Sinilnikova OM, Simard J, Leone M, Dumont M, Neuhausen SL, Struewing JP, Stoppa-Lyonnet D, Barjhoux L, Hughes DJ, Coupiet I, Belotti M, Lasset C, Bonadona V, Bignon YJ, Genetic Modifiers of Cancer Risk in BRCA1/2 Mutation Carriers Study (GEMO), Rebbeck TR, Wagner T, Lynch HT, Domchek SM, Nathanson KL, Garber JE, Weitzel J, Narod SA, Tomlinson G, Olopade OI, Godwin A, Isaacs C, Jakubowska A, Lubinski J, *et al*: **RAD51 135G→C modifies breast cancer risk among BRCA2 mutation carriers: results from a combined analysis of 19 studies.** *Am J Hum Genet* 2007, **81**:1186-1200.
37. Antoniou AC, Pharoah PD, McMullan G, Day NE, Ponder BA, Easton D: **Evidence for further breast cancer susceptibility genes in addition to BRCA1 and BRCA2 in a population-based study.** *Genet Epidemiol* 2001, **21**:1-18.
38. Mavaddat N, Rebbeck TR, Lakhani SR, Easton DF, Antoniou AC: **Incorporating tumour pathology information into breast cancer risk prediction algorithms.** *Breast Cancer Res* 2010, **12**:R28.
39. Lange K, Weeks D, Boehnke M: **Programs for Pedigree Analysis: MENDEL, FISHER, and dGENE.** *Genet Epidemiol* 1988, **5**:471-472.
40. Antoniou AC, Cunningham AP, Peto J, Evans DG, Lalloo F, Narod SA, Risch HA, Eyfjord JE, Hopper JL, Southey MC, Olsson H, Johannsson O, Borg A, Pasini B, Radice P, Manoukian S, Eccles DM, Tang N, Olah E, Anton-Culver H, Warner E, Lubinski J, Gronwald J, Gorski B, Tryggvadottir L, Syrjäkoski K, Kallioniemi OP, Eerola H, Nevanlinna H, Pharoah PD, *et al*: **The BOADICEA model of genetic susceptibility to breast and ovarian cancers: updates and extensions.** *Br J Cancer* 2008, **98**:1457-1466.
41. Boos DD: **On generalised score tests.** *American Statistician* 1992, **46**:327-333.
42. Stacey SN, Sulem P, Zanon C, Gudjonsson SA, Thorleifsson G, Helgason A, Jonasdottir A, Besenbacher S, Kostic JP, Fackenthal JD, Huo D, Adebamowo C, Ogundiran T, Olson JE, Fredericksen ZS, Wang X, Look MP, Sieuwerts AM, Martens JW, Pajares J, Garcia-Prats MD, Ramon-Cajal JM, de Juan A, Panadero A, Ortega E, Aben KK, Vermeulen SH, Asadzadeh F, van Engelenburg KC, Margolin S, *et al*: **Ancestry-shift refinement mapping of the C6orf97-ESR1 breast cancer susceptibility locus.** *PLoS Genet* 2010, **6**: e1001029.

43. Zheng W, Long J, Gao YT, Li C, Zheng Y, Xiang YB, Wen W, Levy S, Deming SL, Haines JL, Gu K, Fair AM, Cai Q, Lu W, Shu XO: **Genome-wide association study identifies a new breast cancer susceptibility locus at 6q25.1.** *Nat Genet* 2009, **41**:324-328.
44. Antoniou AC, Wang X, Fredericksen ZS, McGuffog L, Tarrell R, Sinilnikova OM, Healey S, Morrison J, Kartsonaki C, Lesnick T, Ghoussaini M, Barrowdale D, EMBRACE, Peock S, Cook M, Oliver C, Frost D, Eccles D, Evans DG, Eeles R, Izatt L, Chu C, Douglas F, Paterson J, Stoppa-Lyonnet D, Houdayer C, Mazoyer S, Giraud S, Lasset C, Remenieras A, *et al*: **A locus on 19p13 modifies risk of breast cancer in BRCA1 mutation carriers and is associated with hormone receptor-negative breast cancer in the general population.** *Nat Genet* 2010, **42**:885-892.
45. Milne RL, Benitez J, Nevanlinna H, Heikkinen T, Aittomaki K, Blomqvist C, Arias JI, Zamora MP, Burwinkel B, Bartram CR, Meindl A, Schmutzler RK, Cox A, Brock I, Elliott G, Reed MW, Southey MC, Smith L, Spurdle AB, Hopper JL, Couch FJ, Olson JE, Wang X, Fredericksen Z, Schürmann P, Bremer M, Hillemanns P, Dörk T, Devilee P, van Asperen CJ, *et al*: **Risk of estrogen receptor-positive and -negative breast cancer and single-nucleotide polymorphism 2q35-rs13387042.** *J Natl Cancer Inst* 2009, **101**:1012-1018.
46. Mavaddat N, Antoniou AC, Easton DF, Garcia-Closas M: **Genetic susceptibility to breast cancer.** *Mol Oncol* 2010, **4**:174-191.
47. Wacholder S, Han SS, Weinberg CR: **Inference from a multiplicative model of joint genetic effects or ovarian cancer risk.** *J Natl Cancer Inst* 2011, **103**:82-83.
48. Hammond ME, Hayes DF, Dowsett M, Allred DC, Hagerty KL, Badve S, Fitzgibbons PL, Francis G, Goldstein NS, Hayes M, Hicks DG, Lester S, Love R, Mangu PB, McShane L, Miller K, Osborne CK, Paik S, Perlmutter J, Rhodes A, Sasano H, Schwartz JN, Sweep FC, Taube S, Torlakovic EE, Valenstein P, Viale G, Visscher D, Wheeler T, Williams RB, *et al*: **American Society of Clinical Oncology/College of American Pathologists guideline recommendations for immunohistochemical testing of estrogen and progesterone receptors in breast cancer (unabridged version).** *Arch Pathol Lab Med* 2010, **134**:e48-e72.
49. Collins LC, Botero ML, Schnitt SJ: **Bimodal frequency distribution of estrogen receptor immunohistochemical staining results in breast cancer: an analysis of 825 cases.** *Am J Clin Pathol* 2005, **123**:16-20.
50. Milne RL, Antoniou AC: **Genetic modifiers of cancer risk for BRCA1 and BRCA2 mutation carriers.** *Ann Oncol* 2011, **22**(Suppl 1):i11-i17.
51. Fletcher O, Johnson N, Orr N, Hosking FJ, Gibson LJ, Walker K, Zelenika D, Gut I, Heath S, Palles C, Coupland B, Broderick P, Schoemaker M, Jones M, Williamson J, Chilcott-Burns S, Tomczyk K, Simpson G, Jacobs KB, Chanock SJ, Hunter DJ, Tomlinson IP, Swerdlow A, Ashworth A, Ross G, dos Santos Silva I, Lathrop M, Houlston RS, Peto J: **Novel breast cancer susceptibility locus at 9q31.2: results of a genome-wide association study.** *J Natl Cancer Inst* 2011, **103**:425-435.
52. Gaudet MM, Kirchoff T, Green T, Vijai J, Korn JM, Guiducci C, Segre AV, McGee K, McGuffog L, Kartsonaki C, Morrison J, Healey S, Sinilnikova OM, Stoppa-Lyonnet D, Mazoyer S, Gauthier-Villars M, Sobol H, Longy M, Frenay M, GEMO Study Collaborators, Hogervorst FB, Rookus MA, Collée JM, Hoogerbrugge N, van Roozendaal KE, HEBON Study Collaborators, Piedmonte M, Rubinstein W, Nerenstone S, Van Le L, *et al*: **Common genetic variants and modification of penetrance of BRCA2-associated breast cancer.** *PLoS Genet* 2010, **6**:e1001183.

doi:10.1186/bcr3052

Cite this article as: Mulligan *et al*: Common breast cancer susceptibility alleles are associated with tumour subtypes in BRCA1 and BRCA2 mutation carriers: results from the Consortium of Investigators of Modifiers of BRCA1/2. *Breast Cancer Research* 2011 **13**:R110.

Submit your next manuscript to BioMed Central and take full advantage of:

- Convenient online submission
- Thorough peer review
- No space constraints or color figure charges
- Immediate publication on acceptance
- Inclusion in PubMed, CAS, Scopus and Google Scholar
- Research which is freely available for redistribution

Submit your manuscript at
www.biomedcentral.com/submit

