



## Using additional information on working hours to predict coronary heart disease: a cohort study.

Mika Kivimäki, G David Batty, Mark Hamer, Jane E. Ferrie, Jussi Vahtera, Marianna Virtanen, Michael G. Marmot, Archana Singh-Manoux, Martin J. Shipley

### ► To cite this version:

Mika Kivimäki, G David Batty, Mark Hamer, Jane E. Ferrie, Jussi Vahtera, et al.. Using additional information on working hours to predict coronary heart disease: a cohort study.: Working hours and CHD risk prediction. *Annals of Internal Medicine*, 2011, 154 (7), pp.457-63. 10.1059/0003-4819-154-7-201104050-00003 . inserm-00672633

HAL Id: inserm-00672633

<https://inserm.hal.science/inserm-00672633>

Submitted on 3 Oct 2012


**HAL** is a multi-disciplinary open access archive for the deposit and dissemination of scientific research documents, whether they are published or not. The documents may come from teaching and research institutions in France or abroad, or from public or private research centers.

L'archive ouverte pluridisciplinaire **HAL**, est destinée au dépôt et à la diffusion de documents scientifiques de niveau recherche, publiés ou non, émanant des établissements d'enseignement et de recherche français ou étrangers, des laboratoires publics ou privés.

## ONLINE APPENDIX:

### **Utility of adding information on working hours to the prediction of coronary heart disease: the Whitehall study**

Mika Kivimäki, G. David Batty, Mark Hamer, Jane E. Ferrie, Jussi Vahtera, Marianna Virtanen, Michael G. Marmot, Archana Singh-Manoux and Martin J. Shipley



eFigure1. Flow-diagram of the sample selection

**eTable1. Comparison between included and excluded participants free of prevalent coronary heart disease at study baseline.**

<b>Characteristic*</b>	<b>Included</b>		<b>Excluded</b>		<b>P-value</b>
	<b>N</b>	<b>M<math>\pm</math>SD/%</b>	<b>N</b>	<b>M<math>\pm</math> SD/%</b>	
Age, y	7095	48.8 $\pm$ 5.7	1437	53.6 $\pm$ 6.3	<0.001
Sex, %					<0.001
Male	4986	70.3	864	60.1	
Female	2109	29.7	573	39.9	
Ethnic group, %					<0.001
White	6465	91.1	1253	87.2	
Non-white	630	8.9	184	12.8	
Incident coronary heart disease at follow-up					0.177
None	6903	97.3	1384*	96.5	
Non-fatal myocardial infarction	163	2.3	40	2.8	
Coronary death	29	0.4	10	0.7	

\* 3 Participants excluded because of loss to follow-up

**eTable2. Crude coronary heart disease event rates by categories of 10-year predicted risk calculated using the Framingham risk score alone and in combination with information on long working hours.**

<b>10-year predicted risk category, %</b>	<b>Person-years</b>	<b>N of participants</b>	<b>N of events</b>	<b>Rate*</b>	<b>Rate ratio</b>
<b>Framingham alone</b>					
<5.0	74,695.9	6554	150	2.0	1.00
5.0 – 9.9	4,696.5	436	34	7.2	3.61 (2.48 to 5.23)
≥10.0	1,018.6	105	8	7.9	3.91 (1.92 to 7.96)
<b>Framingham and working hours</b>					
<5.0	74,125.7	6498	143	1.9	1.00
5.0 – 9.9	5,228.0	489	38	7.3	3.77 (2.63 to 5.39)
≥10.0	1,057.1	108	11	10.4	5.39 (2.92 to 9.96)

\*Rate per 1,000 person years

**eTable3. Reclassification of the predicted 10-year risk of incident coronary heart disease for a risk score based on both the Framingham risk score and long working hours compared to the Framingham risk score alone among subgroups of the Whitehall participants.**

		Reclassified			
Subgroup	N	Increased risk	Decreased risk	Net correctly reclassified (%)	
Excluding non - white	CHD-case	159	11	3	5.0%
	Non-case	6306	125	90	-0.6%
	NRI (95%CI)				4.5% (-0.2 to 9.1)
	P-value				p=0.058
Excluding those on medication*	CHD-case	163	12	4	4.9%
	Non-case	6495	118	84	-0.5%
	NRI (95%CI)				4.4% (-0.4 to 9.2)
	P-value				p=0.075
Non-diabetics	CHD-case	180	13	4	5.0%
	Non-case	6716	126	95	-0.5%
	NRI (95%CI)				4.5% (0.0 to 9.0)
	P-value				p=0.049

\*Antihypertensive medication (n=394), lipid-lowering drugs (n=43) and anti-platelet agents (n=17).