

## **IFN-gamma mediates the rejection of haematopoietic stem cells in IFN-gammaR1-deficient hosts.**

Martin Rottman, Claire Soudais, Guillaume Vogt, Laurent Renia, Jean-François Emile, Hélène Decaluwe, Jean-Louis Gaillard, Jean-Laurent Casanova

► **To cite this version:**

Martin Rottman, Claire Soudais, Guillaume Vogt, Laurent Renia, Jean-François Emile, et al.. IFN-gamma mediates the rejection of haematopoietic stem cells in IFN-gammaR1-deficient hosts.. PLoS Medicine, Public Library of Science, 2008, 5 (1), pp.e26. <10.1371/journal.pmed.0050026>. <inserm-00700812>

**HAL Id: inserm-00700812**

**<https://www.hal.inserm.fr/inserm-00700812>**

Submitted on 24 May 2012

**HAL** is a multi-disciplinary open access archive for the deposit and dissemination of scientific research documents, whether they are published or not. The documents may come from teaching and research institutions in France or abroad, or from public or private research centers.

L'archive ouverte pluridisciplinaire **HAL**, est destinée au dépôt et à la diffusion de documents scientifiques de niveau recherche, publiés ou non, émanant des établissements d'enseignement et de recherche français ou étrangers, des laboratoires publics ou privés.

# IFN- $\gamma$ Mediates the Rejection of Haematopoietic Stem Cells in IFN- $\gamma$ R1-Deficient Hosts

Martin Rottman<sup>1,2,3</sup>✉, Claire Soudais<sup>1,2</sup>✉\*, Guillaume Vogt<sup>1,2</sup>, Laurent Renia<sup>4,5,6</sup>, Jean-François Emile<sup>7</sup>, Hélène Decaluwe<sup>1,2</sup>, Jean-Louis Gaillard<sup>3</sup>, Jean-Laurent Casanova<sup>1,2,8</sup>

**1** Laboratoire de Génétique Humaine des Maladies Infectieuses, INSERM, U550, Paris, France, **2** Université Paris René Descartes, Faculté de Médecine Necker-Enfants Malades, Paris, France, **3** Hôpital Raymond Poincaré, Faculté de Médecine Paris-Ile de France-Ouest, UPRES Sud, EA3647, Laboratoire de Microbiologie, Garches, France, **4** Institut Cochin, INSERM, U567, Paris, France, **5** CNRS, UMR8104, Paris, France, **6** Université René Descartes, Hôpital Cochin, Paris, France, **7** Hôpital Ambroise Paré, Laboratoire d'Anatomo-Pathologie, Boulogne, France, **8** Unité d'Immunologie et Hématologie Pédiatriques, Hôpital Necker-Enfants Malades, Paris, France

**Funding:** This work received financial support from the association Vaincre La Mucoviscidose. The Laboratory of Human Genetics of Infectious Diseases is supported in part by grants from the Schlumberger and BNP Paribas Foundations, the March of Dimes, and by EU grant QLK2-CT-2002-00846. Jean-Laurent Casanova is an International Scholar of the Howard Hughes Medical Institute. C. Soudais is supported by the Association Française contre les Myopathies. The funders had no role in study design, data collection and analysis, decision to publish, or preparation of the manuscript.

**Competing Interests:** The authors have declared that no competing interests exist.

**Academic Editor:** Adrian J Thrasher, Institute of Child Health, United Kingdom

**Citation:** Rottman M, Soudais C, Vogt G, Renia L, Emile JF, et al. (2008) IFN- $\gamma$  mediates the rejection of haematopoietic stem cells in IFN- $\gamma$ R1-deficient hosts. *PLoS Med* 5(1): e26. doi:10.1371/journal.pmed.0050026

**Received:** May 9, 2007

**Accepted:** December 10, 2007

**Published:** January 29, 2008

**Copyright:** © 2008 Rottman et al. This is an open-access article distributed under the terms of the Creative Commons Attribution License, which permits unrestricted use, distribution, and reproduction in any medium, provided the original author and source are credited.

**Abbreviations:** BCG, *Mycobacterium bovis* bacillus Calmette-Guérin; DKO, double knock-out; HSCT, haematopoietic stem cell transplantation; IFN- $\gamma$ , Interferon- $\gamma$ ; IFN- $\gamma$ R1, Interferon- $\gamma$  receptor 1; rAAV-IFN- $\gamma$ , recombinant adeno-associated viral vector encoding IFN- $\gamma$

\* To whom correspondence should be addressed. E-mail: soudais@necker.fr

✉ These authors contributed equally to this work.

## ABSTRACT

### Background

Interferon- $\gamma$  receptor 1 (IFN- $\gamma$ R1) deficiency is a life-threatening inherited disorder, conferring predisposition to mycobacterial diseases. Haematopoietic stem cell transplantation (HSCT) is the only curative treatment available, but is hampered by a very high rate of graft rejection, even with intra-familial HLA-identical transplants. This high rejection rate is not seen in any other congenital disorders and remains unexplained. We studied the underlying mechanism in a mouse model of HSCT for IFN- $\gamma$ R1 deficiency.

### Methods and Findings

We demonstrated that HSCT with cells from a syngenic C57BL/6 *Ifngr1*<sup>+/+</sup> donor engrafted well and restored anti-mycobacterial immunity in naive, non-infected C57BL/6 *Ifngr1*<sup>-/-</sup> recipients. However, *Ifngr1*<sup>-/-</sup> mice previously infected with *Mycobacterium bovis* bacillus Calmette-Guérin (BCG) rejected HSCT. Like infected IFN- $\gamma$ R1-deficient humans, infected *Ifngr1*<sup>-/-</sup> mice displayed very high serum IFN- $\gamma$  levels before HSCT. The administration of a recombinant IFN- $\gamma$ -expressing AAV vector to *Ifngr1*<sup>-/-</sup> naive recipients also resulted in HSCT graft rejection. Transplantation was successful in *Ifngr1*<sup>-/-</sup>  $\times$  *Ifng*<sup>-/-</sup> double-mutant mice, even after BCG infection. Finally, efficient antibody-mediated IFN- $\gamma$  depletion in infected *Ifngr1*<sup>-/-</sup> mice in vivo allowed subsequent engraftment.

### Conclusions

High serum IFN- $\gamma$  concentration is both necessary and sufficient for graft rejection in IFN- $\gamma$ R1-deficient mice, inhibiting the development of heterologous, IFN- $\gamma$ R1-expressing, haematopoietic cell lineages. These results confirm that IFN- $\gamma$  is an anti-haematopoietic cytokine in vivo. They also pave the way for HSCT management in IFN- $\gamma$ R1-deficient patients through IFN- $\gamma$  depletion from the blood. They further raise the possibility that depleting IFN- $\gamma$  may improve engraftment in other settings, such as HSCT from a haplo-identical or unrelated donor.

*The Editors' Summary of this article follows the references.*

## Introduction

Complete deficiency of the ligand-binding chain of the interferon- $\gamma$  receptor (IFN- $\gamma$ R1) is an autosomal recessive disorder described in 1996 as the first genetic aetiology of the syndrome of Mendelian Susceptibility to Mycobacterial Diseases (“MSMD”: MIM 209950) [1–3]. Causal mutations either abrogate cell surface IFN- $\gamma$ R1 expression [4,5] or prevent IFN- $\gamma$  recognition due to the production of surface-expressed, non-functional receptors [6]. Both types of IFN- $\gamma$ R1 deficiency result in a complete loss of cellular responses to IFN- $\gamma$ . This disorder confers a profound and selective susceptibility to weakly virulent mycobacteria, such as *Mycobacterium bovis* bacillus Calmette Guérin (BCG) vaccines and environmental mycobacteria [7], as reviewed in [3]. Humans are also susceptible to the more virulent *M. tuberculosis* [7,8]. Other infectious diseases are rare, with the exception of salmonellosis, which has been diagnosed in several patients [9]. Listeriosis and a few viral diseases were each diagnosed in single patients [7,10]. Humans do not produce mature granulomas in response to mycobacteria, and instead display poorly delimited, poorly differentiated, multibacillary tissue lesions [11]. They present with early-onset, disseminated, recurrent, and multiple mycobacterial infections. Most patients die in early childhood, with only one-third surviving to the age of 15 years [7].

The treatment of patients with complete IFN- $\gamma$ R1 deficiency is particularly difficult [3,7]. Antibiotics alone do not achieve permanent clinical remission. Unlike patients with other MSMD aetiologies, those with complete IFN- $\gamma$ R1 (or IFN- $\gamma$ R2) deficiency do not benefit from exogenous IFN- $\gamma$  administration, owing to the lack of a specific receptor. The remission of mycobacterial infection following the first case of haematopoietic stem cell transplantation (HSCT) in such an individual provided proof-of-principle that IFN- $\gamma$ R1 deficiency is primarily a haematopoietic disorder, despite the ubiquitous expression of IFN- $\gamma$ R1 [12]. However, subsequent attempts revealed a very high rate of primary and secondary rejection in HLA-identical HSCT [13–15]. Nine patients received a total of 12 transplants. Four of these patients died within eight months of transplantation—from mycobacterial disease in two cases—and three individuals rejected the graft. Two of the five patients who survived presented only very low levels of chimerism and no chimerism was observed in a third. One individual had a low-grade infection at the time of the treatment [15]. Only three individuals have remained free from infectious complications [13,15] and are currently healthy, nine, seven, and seven years after HSCT. This rate of HLA-identical graft rejection is by far the highest reported for HSCT to treat primary immunodeficiencies or congenital disorders [16]. These observations indicate that HSCT is potentially curative in IFN- $\gamma$ R1-deficient individuals, but associated with a particularly high and unexplained rate of graft rejection, resulting in unacceptable morbidity and mortality rates.

We used mice selectively deficient in IFN- $\gamma$  R1 (*Ifngr*<sup>−/−</sup>), IFN- $\gamma$  (*Ifng*<sup>−/−</sup>), or both (*Ifngr*<sup>−/−</sup> × *Ifng*<sup>−/−</sup>) [17,18], as a means of investigating the mechanism of graft rejection in IFN- $\gamma$ R1-deficient patients. These mice are susceptible to several intracellular pathogens [17,19–23]. In particular, *Ifngr*<sup>−/−</sup> mice are highly susceptible to mycobacteria, such as *M. tuberculosis* [24,25], *M. avium* [26] and BCG [27]. *Ifngr*<sup>−/−</sup> mice

die within nine weeks of the intravenous injection of  $2 \times 10^7$  cells of BCG, whereas control mice survive such injections [27]. BCG is also the most common pathogen in IFN- $\gamma$ R1-deficient individuals [7]. On histological examination, these patients have tissue lesions containing large numbers of acid-fast BCG, with small numbers of structurally impaired granulomas and high tissue loads of mycobacteria [4,11]. The same infectious phenotype has been observed in *Ifng*<sup>−/−</sup> mice infected with BCG [18]. Thus, the extreme susceptibility of *Ifngr*<sup>−/−</sup> mice to mycobacterial infections, including BCG in particular, mimics the human condition, making these mice an ideal model for studies of the mechanisms underlying the HSCT failure observed in IFN- $\gamma$ R1-deficient humans.

## Materials and Methods

### Animals

Specific pathogen-free, *Ifngr*<sup>+/+</sup> C57BL/6 mice were purchased from Charles River (L'Arbresle, France) and used at six to nine weeks of age. Interferon gamma receptor chain 1 (*Ifngr*<sup>−/−</sup>, stock no. 003288, N10) [17] and interferon gamma (*Ifng*<sup>−/−</sup>, stock no. 002287, N10) [18] mice were purchased from JaxLab and reared at the CDTA (Centre de Distribution, Typage et Archivage animal, Orléans, France). Both *Ifngr*<sup>−/−</sup> and *Ifng*<sup>−/−</sup> mice were backcrossed onto the C57BL/6 background for more than ten generations. Double knock-out (DKO) mice (*Ifngr*<sup>−/−</sup> × *Ifng*<sup>−/−</sup>) were generated in a pathogen-free animal facility, by mating homozygous *Ifng*<sup>−/−</sup> mice with *Ifngr*<sup>−/−</sup> mice. The resulting F1 animals were then intercrossed to generate double homozygous mice, identified by PCR on tail DNA (primers and conditions available from the JaxLab Web site [<http://jaxmice.jax.org/strain/002287.html> and <http://jaxmice.jax.org/strain/003288.html>]). All experiments and procedures were performed in accordance with French Ministry of Agriculture regulations for animal experimentation (1987) and the guidelines of our institution's animal welfare committee.

### Haematopoietic Stem Cell Transfer

Recipient mice were subjected to intensive or milder doses (550 rads to 1,200 rads) of body irradiation, using a Cs137 radioactive source (Pasteur Institute). Immune reconstitution was then initiated by the intravenous administration, via a lateral tail vein, of two million total bone marrow cells freshly isolated from donor mice. Briefly, bone marrow cells were flushed into PBS from the femurs and tibias of donor mice. Red blood cells were lysed and the remaining cells were counted before bone marrow cell transfer. After engraftment, mice were kept in a specific-pathogen-free animal facility, in filter-topped cages in an isolation room. All caging procedures and manipulations were carried out in a laminar flow hood. Mice were used for experimental infection eight weeks after HSCT.

### Analysis of Chimerism and Reconstitution

The extent of haematopoietic reconstitution by the donor phenotype was assessed every two weeks, during the eight weeks before experimental infection, using specific antibodies. Immunofluorescence analyses were carried out on whole blood. Briefly, donor bone marrow cells expressed the Ly5.1 marker, whereas recipient leukocytes expressed the Ly5.2 marker. Chimerism was therefore assessed as the

percentage of cells expressing the Ly5.1 isotype (engraftment). Antibodies against the following surface antigens (all from BD Bioscience Pharmingen) were used, as FITC or PE conjugates, to evaluate peripheral reconstitution: TCR $\alpha\beta$ , B220, Mac-1 and GRI. Blood (25  $\mu$ l) was first blocked with 24G2 serum, then stained with specific antibodies. Red blood cells were then lysed with PharM Lyse buffer and flow cytometry was carried out on the remaining cells (FACScan, Becton Dickinson).

### Experimental Infection with BCG and Determination of the Number of Colony-Forming Units

BCG strain Pasteur 1173P<sub>2</sub> was used [28]. BCG was prepared as described elsewhere [29], frozen in aliquots and stored at  $-80^{\circ}\text{C}$ . For each infection experiment, groups of non-irradiated animals were included as positive and negative controls of infection. Bacteria were counted in the spleen and liver, as described elsewhere [29]. In brief, blood samples were taken from six to eight mice for each point. The mice were then killed, and their organs were dissected out, placed in 2 ml screw-cap tubes filled with sterile water and homogenized with a 5 mm stainless steel ball (SKF) using a mini-8 bead-beater (Biospec). Spots (50  $\mu$ l) of serial five-fold dilutions of this suspension were plated on Middlebrook 7H11 medium supplemented with OADC (Difco). The plates were incubated for 20 d at  $37^{\circ}\text{C}$ , under a humidified atmosphere containing 5% CO<sub>2</sub>, and colonies were counted using a stereoscopic binocular microscope with a detection threshold of 170 CFU per organ.

### ELISA

Serum was recovered from whole blood samples following coagulation. Serum aliquots were frozen at  $-20^{\circ}\text{C}$  for quantitative sandwich immunoassays for IFN- $\gamma$ . Mouse ELISA kits from R&D Systems were used, according to the manufacturer's recommendations, in all experiments. The detection limit of the kit was less than 2 pg/ml.

### RNA Purification and Northern Blot Analysis

Tissue samples were lysed in 1 ml of Trizol (Invitrogen) and the resulting suspensions were immediately frozen and stored at  $-80^{\circ}\text{C}$ . Total RNA was extracted with a kit based on the acid phenol-guanidine method, according to the manufacturer's instructions. Total RNA concentration was estimated by spectrophotometry. Total RNA (15  $\mu$ g in 50% formamide) was blotted onto a nylon membrane. The membrane was incubated for three hours at  $80^{\circ}\text{C}$  and was then washed in 2 $\times$ SSC before hybridisation with murine IFN- $\gamma$  and GAPDH probes generated by PCR (primers available on request) and labelled with <sup>32</sup>P by random priming. Each autoradiograph was densitometrically scanned and levels of the specific IFN- $\gamma$  transcripts were normalised with respect to GAPDH transcript levels.

### Preparation and Use of Recombinant Adeno-Associated Vectors and Injection

The mouse IFN- $\gamma$  cDNA was amplified by PCR from the L929 cell line and inserted into convenient restriction sites in the pGG1 vector. The resulting plasmid was sequenced to check the integrity of the mIFN- $\gamma$  cDNA. We prepared rAAV-IFN- $\gamma$  particles, as previously described [30], by a helper-virus-free method involving the triple transfection of HEK

293 cells. Physical particles were estimated by dot plots, and  $1.7 \times 10^{12}$  such particles were obtained per ml of preparation [31]. We assessed rAAV-IFN- $\gamma$  particle function in vivo, by injecting  $5 \times 10^{10}$  rAAV particles, in a final volume of 100  $\mu$ l, into the left gastrocnemius muscle of recipient *Ifngr*<sup>-/-</sup> mice. Serum samples were recovered at various time points after injection and mIFN- $\gamma$  was quantified by ELISA. A rAAV vector including the *lacZ* gene was used as negative control.

### Antibody-Mediated Cytokine Neutralisation

Animals were infected with  $10^6$  CFU of BCG. Fourteen days later, they received an intraperitoneal injection of a mixture of 0.8 mg of purified rat anti-mouse IFN- $\gamma$  antibody (clone XMG1 [32]) and 0.8 mg of purified rat anti-mouse IL-12p40 antibody (clone C15-1 [33]). Antibodies were purified from the culture supernatant by ammonium sulphate precipitation and shown to function effectively in a mouse model of *Plasmodium* infection [34]. The control group received an injection of 1.6 mg of irrelevant purified rat IgG antibody (Sigma). In all groups, the injection was repeated on days 15, 17, 20, and 24. Bone marrow from *Ifngr*<sup>+/+</sup> donors was transferred to the recipient mice 23 d after sublethal body irradiation (550 rad). Serum IFN- $\gamma$  concentration and chimerism were monitored throughout the experiment.


### Histology

Organs were fixed by immersion for 24 h in 3.7% formaldehyde and were then transferred to 70% ethanol for 24 h to 48 h before embedding in paraffin. Sections (5  $\mu$ m) were cut on a rotary microtome, stretched in a water bath, mounted on glass slides and stained with haematoxylin-eosin-saffron (HES) or by the Zielh-Neelsen method.

## Results

### *Ifngr*<sup>-/-</sup> Mice Are Susceptible to BCG Infection

C57BL/6 *Ifngr*<sup>-/-</sup> mice were unable to control BCG infection following intravenous injection with  $\geq 10^2$  CFU (Figure 1A). Deficient animals survived  $29 \pm 2$  weeks (mean  $\pm$  SD) following challenge with  $10^2$  CFU, whereas no mortality was observed in wild-type C57BL/6 control mice (*Ifngr*<sup>+/+</sup>) during 12 mo of follow-up, even after infection with the highest counts of CFU. The survival of deficient animals was inversely proportional to the size of the inoculum (Figure 1A), with mortality ranging from 29 wk with  $10^2$  CFU to 12 wk with  $10^6$  CFU. *Ifngr*<sup>-/-</sup> mice challenged i.v. with  $10^6$  CFU of BCG were unable to control the infection: the bacterial load in the spleen increased to  $7.4 \pm 0.3$  (Log<sub>10</sub> CFU) by 45 d after infection and exceeded 8 Log<sub>10</sub> CFU in dying animals (Figure 1B). In *Ifngr*<sup>+/+</sup> mice, the BCG burden was controlled, with a decrease in CFU observed by day 45, and CFU numbers falling to 5 Log<sub>10</sub> by day 90. Granuloma formation was altered in BCG-infected *Ifngr*<sup>-/-</sup> mice. Two weeks after infection, *Ifngr*<sup>-/-</sup> mice had fewer and smaller granulomas in both the spleen and the liver than infected *Ifngr*<sup>+/+</sup> control mice. Granulomas were mostly lymphoid, with no recruitment of epithelioid cells, whereas the granulomas observed in *Ifngr*<sup>+/+</sup> BCG-infected mice contained mostly epithelioid cells and a few lymphocytes (unpublished data). By day 90 post-infection, *Ifngr*<sup>-/-</sup> mice presented massive mycobacterial dissemination associated with large necrotising granulomas. In contrast, a very small number of small, well-delimited granulomas were



**Figure 1. *Ifngr1*<sup>-/-</sup> Mice Are Susceptible to BCG Infection**  
 (A) *Ifngr1*<sup>-/-</sup> mice were infected with between 10<sup>2</sup> and 10<sup>6</sup> CFU of BCG, and animal survival (five animals per group) was monitored thereafter. (B) Splenic mycobacterial loads were determined on days 4, 15, 45, and 90 in infected *Ifngr1*<sup>-/-</sup> and *Ifngr1*<sup>+/+</sup> mice; means of five animals per point are shown.  
 doi:10.1371/journal.pmed.0050026.g001

observed in infected *Ifngr1*<sup>+/+</sup> control mice (unpublished data). These data are consistent with the absence of mature granulomas observed in patients entirely lacking IFN- $\gamma$ R1 [11]. Our results confirm that *Ifngr1*<sup>-/-</sup> mice are highly susceptible to *M. bovis* BCG infection, consistent with previous reports [27], and validate this model for the study of IFN- $\gamma$ R1 deficiency in humans.

### HSCT Restores Anti-Mycobacterial Immunity in *Ifngr1*<sup>-/-</sup> Mice

We subjected C57BL/6 *Ifngr1*<sup>-/-</sup> mice to HSCT with sex- and age-matched syngenic C57BL/6 *Ifngr1*<sup>+/+</sup> donors with an intensive total body irradiation conditioning regimen (1,200 rad). Several doses of bone marrow were tested and we found that 2 million cells was the dose most comparable to HSCT in humans. Leukocyte chimerism was complete, with low levels of residual autologous cells nine wk after treatment. Peripheral reconstitution was achieved, with the expected counts of lymphoid (T and B) and myeloid (macrophages and


granulocytes) cells (Figure 2A, unpublished data). We then evaluated the ability of the recipient mice to control BCG infection. Ten weeks after HSCT, animals were challenged i.v. with 10<sup>6</sup> CFU of BCG. They were killed 45 d later and BCG load in the spleen was determined. *Ifngr1*<sup>-/-</sup> animals transplanted with *Ifngr1*<sup>-/-</sup> bone marrow were unable to control BCG infection, like *Ifngr1*<sup>+/+</sup> animals receiving *Ifngr1*<sup>-/-</sup> bone marrow (Figure 2B). *Ifngr1*<sup>-/-</sup> animals receiving *Ifngr1*<sup>+/+</sup> bone marrow controlled the infection as efficiently as *Ifngr1*<sup>+/+</sup> mice receiving *Ifngr1*<sup>+/+</sup> bone marrow or mice with no graft (Figure 2B). HSCT recipients conditioned with a milder regimen (550 rads) displayed mixed chimerism, with only about 50% donor Ly5.1 lymphocytes (Figure 2C). The donor haematopoietic compartment had nonetheless restored BCG growth control 45 d after infection (Figure 2D). Thus, the restoration of IFN- $\gamma$ R1 expression in the haematopoietic compartment alone, even in only a fraction of the compartment, is sufficient to confer resistance to BCG infection. Mycobacterial disease in mice with IFN- $\gamma$ R1 deficiency therefore results from the specific absence of IFN- $\gamma$ R1 in the haematopoietic compartment, consistent with reported data for HSCT in human patients [12,13].

### Rejection of HSCT in *Ifngr1*<sup>-/-</sup> Mice Previously Infected with BCG

All the IFN- $\gamma$ R1-deficient patients undergoing HSCT had a history of mycobacterial disease, as HSCT has not yet been attempted in an asymptomatic child [12–15]. We thus infected animals with BCG before HSCT. *Ifngr1*<sup>-/-</sup> and *Ifngr1*<sup>+/+</sup> mice were infected intravenously with 10<sup>6</sup> CFU BCG and subjected to HSCT 40 d later, with sex- and age-matched syngenic *Ifngr1*<sup>+/+</sup> donors. Following intense irradiation (1,200 rads), both *Ifngr1*<sup>-/-</sup> and *Ifngr1*<sup>+/+</sup> BCG-infected mice displayed successful immune reconstitution with *Ifngr1*<sup>+/+</sup> bone marrow (unpublished data). However, with a milder conditioning regimen (550 rads) more closely mimicking the situation in human transplant patients in terms of the chimerism post HSCT [12–15], *Ifngr1*<sup>-/-</sup> mice infected with BCG before HSCT rejected the graft, with Ly5.1 lymphocytes from the donor phenotype accounting for less than 2% of the circulating cells, whereas about 50% chimerism was observed in the absence of BCG infection (Figure 3A). Serving as a control, *Ifngr1*<sup>+/+</sup> mice infected with BCG and subjected to HSCT with *Ifngr1*<sup>+/+</sup> donor marrow, under the same milder conditioning regimen, displayed successful engraftment and reconstitution. Bacterial loads, determined 11 wk after infection, reached 8.4 Log<sub>10</sub> CFU per spleen in the *Ifngr1*<sup>-/-</sup> cohort, versus only 4.5 Log<sub>10</sub> CFU in *Ifngr1*<sup>+/+</sup> mice (Figure 3B). Histological observations confirmed dissemination in the *Ifngr1*<sup>-/-</sup> cohort, contrasting with small, well-delimited granulomas in *Ifngr1*<sup>+/+</sup> animals (unpublished data). Thus, *Ifngr1*<sup>-/-</sup> mice infected with BCG reject HSCT, with features mimicking the graft rejection observed in IFN- $\gamma$ R1-deficient patients.

### HSCT Graft Rejection Is Associated with High Serum IFN- $\gamma$ Levels

High serum IFN- $\gamma$  concentration is a hallmark of human complete IFN- $\gamma$ R1 deficiency [35]. We thus monitored serum IFN- $\gamma$  levels in BCG-infected *Ifngr1*<sup>-/-</sup> and *Ifngr1*<sup>+/+</sup> mice. IFN- $\gamma$  was detected as early as 10 d post-infection and its concentration gradually increased towards a plateau at about 6 ng/ml within six wk of infection in *Ifngr1*<sup>-/-</sup> mice, confirming



**Figure 2.** HSCT Restores Anti-Mycobacterial Immunity in *Ifngr1*<sup>-/-</sup> Mice

(A) *Ifngr1*<sup>-/-</sup> and *Ifngr1*<sup>+/+</sup> mice (five animals per group) expressing the Ly5.2 marker were subjected to HSCT with bone marrow from *Ifngr1*<sup>+/+</sup> mice expressing the Ly5.1 marker, after intense irradiation (1,200 rads). Chimerism, assessed by determining the surface expression of Ly5.1 and Ly5.2 on lymphocytes, and peripheral reconstitution, assessed by determining the surface expression of TCR $\alpha\beta$  and B220 markers on lymphocytes, were analysed by flow cytometry nine weeks after HSCT treatment.

(B) *Ifngr1*<sup>-/-</sup> and *Ifngr1*<sup>+/+</sup> mice were subjected to HSCT with bone marrow from *Ifngr1*<sup>-/-</sup> or *Ifngr1*<sup>+/+</sup> mice. HSCT-treated mice were then infected with BCG and bacterial loads were determined 45 d later (five animals per group).


(C) *Ifngr1*<sup>-/-</sup> and *Ifngr1*<sup>+/+</sup> mice expressing the Ly5.2 marker were subjected to HSCT with bone marrow from *Ifngr1*<sup>+/+</sup> mice expressing the Ly5.1 marker, after mild irradiation (550 rads). Chimerism and peripheral reconstitution were analysed by flow cytometry nine weeks after HSCT treatment.

(D) *Ifngr1*<sup>-/-</sup> and *Ifngr1*<sup>+/+</sup> mice were subjected to HSCT with bone marrow from *Ifngr1*<sup>+/+</sup> mice after intense or mild irradiation, infected with BCG and bacterial load was determined 45 d later.

doi:10.1371/journal.pmed.0050026.g002

previous observations [36]. IFN- $\gamma$  remained barely detectable in infected *Ifngr1*<sup>+/+</sup> mice (Figure 4A). Serum IFN- $\gamma$  levels, determined 30 d after BCG infection in *Ifngr1*<sup>-/-</sup> and *Ifngr1*<sup>+/+</sup> mice previously subjected to HSCT, were found to correlate with the control of BCG infection. IFN- $\gamma$  levels were high in *Ifngr1*<sup>-/-</sup> animals engrafted with *Ifngr1*<sup>-/-</sup> bone marrow and only marginally elevated in *Ifngr1*<sup>+/+</sup> animals engrafted with *Ifngr1*<sup>-/-</sup> bone marrow (Figure 4B). In these conditions IFN- $\gamma$  levels were very similar to those in BCG-infected non-transplanted *Ifngr1*<sup>-/-</sup> mice (Figure 4A). IFN- $\gamma$  was almost

undetectable in both *Ifngr1*<sup>-/-</sup> mice engrafted with *Ifngr1*<sup>+/+</sup> bone marrow and *Ifngr1*<sup>+/+</sup> mice engrafted with *Ifngr1*<sup>+/+</sup> bone marrow, as for non-transplanted *Ifngr1*<sup>+/+</sup> animals (Figure 4B). Following HSCT with mild conditioning after *M. bovis* BCG infection, IFN- $\gamma$  concentration seemed to be high only in the cohort of *Ifngr1*<sup>-/-</sup> animals infected at the time of cell transfer, and in such cases, HSCT was unsuccessful (Figures 3A and 4C). These results were confirmed by quantification of IFN- $\gamma$  transcripts in the spleens of infected animals. Infected *Ifngr1*<sup>-/-</sup> mice contained significantly larger amounts of IFN- $\gamma$



**Figure 3.** Rejection of the HSCT Graft in *Ifngr1*<sup>-/-</sup> Mice Previously Infected with BCG

(A) Chimerism was determined by assessing the surface expression of Ly5.1 (donor cells) on lymphocytes, in *Ifngr1*<sup>-/-</sup> and *Ifngr1*<sup>+/+</sup> mice treated by HSCT with bone marrow from *Ifngr1*<sup>+/+</sup> mice, nine weeks post HSCT. *Ifngr1*<sup>-/-</sup> and *Ifngr1*<sup>+/+</sup> mice were infected with BCG either before or after HSCT (five animals per group).

(B) Bacterial loads were determined after the treatment of *Ifngr1*<sup>-/-</sup> and *Ifngr1*<sup>+/+</sup> mice previously infected with BCG (three animals per group), by HSCT with bone marrow from *Ifngr1*<sup>+/+</sup> mice.

doi:10.1371/journal.pmed.0050026.g003

mRNA than infected *Ifngr1*<sup>+/+</sup> animals (Figure 4D). Basal levels of IFN- $\gamma$  were detected in non-infected *Ifngr1*<sup>-/-</sup> and *Ifngr1*<sup>+/+</sup> mice. As expected, no IFN- $\gamma$  mRNA was detected in *Ifngr1*<sup>-/-</sup> mice. The amounts of IFN- $\gamma$  mRNA in infected *Ifngr1*<sup>-/-</sup> and *Ifngr1*<sup>+/+</sup> mice differed by a factor of 2.25 ( $p \leq 0.007$ ). Thus, serum IFN- $\gamma$  concentration increased with BCG infection and was inversely correlated with HSCT engraftment in the murine model of IFN- $\gamma$ R1 deficiency.

#### rAAV-IFN- $\gamma$ Injection Promotes Graft Rejection


We tested the hypothesis that high circulating IFN- $\gamma$  levels are responsible for HSCT graft rejection, using a recombinant adeno-associated viral vector encoding IFN- $\gamma$  (rAAV-IFN- $\gamma$ ). Following the intramuscular injection of  $5 \times 10^{10}$  rAAV-IFN- $\gamma$  physical particles in non-infected *Ifngr1*<sup>-/-</sup> mice, IFN- $\gamma$  was secreted ectopically in the muscle and accumulated in the serum (unpublished data), resulting on day 15 in serum IFN- $\gamma$  levels ranging from 8 ng/ml to 11 ng/ml. Stable and sustained serum IFN- $\gamma$  levels of 6 ng/ml to 7.5 ng/ml were obtained 30 d after rAAV-IFN- $\gamma$  injection (Figure 5A), approaching those previously quantitated after BCG infection of *Ifngr1*<sup>-/-</sup> mice. HSCT with mild conditioning was thus performed in these mice on day 30 following rAAV-IFN- $\gamma$  injection. In these conditions, HSCT graft rejection occurred in *Ifngr1*<sup>-/-</sup> recipients, with donor Ly5.1 lymphocytes accounting for only  $0.46\% \pm 0.1\%$  of cells (Figure 5B). In contrast, control *Ifngr1*<sup>+/+</sup> mice given intramuscular injections of rAAV-IFN- $\gamma$  displayed successful engraftment and immune reconstitution. No detectable accumulation of IFN- $\gamma$  occurred in *Ifngr1*<sup>+/+</sup> mice, probably because of the rapid clearance of IFN- $\gamma$  by IFN- $\gamma$ R1-proficient cells. We detected minor increases in the serum levels of cytokines known to be involved in the IL12/23-IFN- $\gamma$  circuit (IL12p40 in particular) in both naive and infected *Ifngr1*<sup>-/-</sup> and control *Ifngr1*<sup>+/+</sup> mice, at various stages of HSCT, with no repercussions outside this pathway (unpublished data). These data suggest that high circulating IFN- $\gamma$  levels at the time of HSCT promote graft rejection in IFN- $\gamma$ R1-deficient mice, even in the absence of BCG infection.

#### Infected DKO *Ifngr1*<sup>-/-</sup> $\times$ *Ifngr1*<sup>-/-</sup> Mice Show No Graft Rejection after HSCT

When infected with  $10^6$  CFU of BCG, *Ifngr1*<sup>-/-</sup>  $\times$  *Ifngr1*<sup>-/-</sup> double knock-out (DKO) mice were unable to control the infection and died within 12 wk of infection (unpublished data). As expected, IFN- $\gamma$  was undetectable in serum (Figure 5C). Uninfected DKO mice displayed levels of chimerism after HSCT similar to those observed in *Ifngr1*<sup>-/-</sup> mice after total or mild conditioning (unpublished data). DKO mice controlled infection if HSCT with wild-type bone marrow was carried out before BCG infection (unpublished data). We also infected DKO mice with BCG and subjected them to HSCT, as described above. Bone marrow reconstitution was successful, and the level of chimerism obtained was similar to that in *Ifngr1*<sup>+/+</sup> and *Ifngr1*<sup>-/-</sup> HSCT-treated and BCG-infected mice (Figure 5D). Successful HSCT in these recipient mice ruled out a direct role for BCG infection, other than through IFN- $\gamma$  induction, as the BCG burden in DKO mice was similar to that in *Ifngr1*<sup>-/-</sup> mice before cell transfer. We injected  $5 \times 10^{10}$  physical particles of rAAV-IFN- $\gamma$  into DKO mice, resulting in high serum IFN- $\gamma$  levels (4 ng/ml, 30 d after injection; Figure 5A). Following HSCT, engraftment was almost undetectable in these mice, with a mean of  $2.4\% (\pm 0.5\%)$  donor Ly5.1 lymphocytes (Figure 5B). Thus, in the absence of IFN- $\gamma$ , *Ifngr1*<sup>-/-</sup> mice displayed no rejection following HSCT, despite infection with BCG. In contrast, in the presence of high serum levels of IFN- $\gamma$ , rejection occurred in these mice, even in the absence of BCG infection. In conclusion, these data clearly support the hypothesis that high circulating IFN- $\gamma$  levels are necessary and sufficient for HSCT graft rejection in *Ifngr1*<sup>-/-</sup> mice.

#### In Vivo Neutralisation of Circulating IFN- $\gamma$ Allows HSCT Engraftment

High serum IFN- $\gamma$  concentration at the time of HSCT has a deleterious effect on engraftment. We tried to determine whether blood depletion of IFN- $\gamma$  could render HSCT of infected *Ifngr1*<sup>-/-</sup> mice successful. A first series of experiments




**Figure 4.** HSCT Graft Rejection Is Associated with High Serum IFN- $\gamma$  Levels

(A) IFN- $\gamma$  levels were measured over time, in the serum of *Ifngr1*<sup>-/-</sup> and *Ifngr1*<sup>+/+</sup> mice, after BCG infection (five animals per group).  
 (B) IFN- $\gamma$  levels were measured after the HSCT treatment of *Ifngr1*<sup>-/-</sup> and *Ifngr1*<sup>+/+</sup> mice with either *Ifngr1*<sup>-/-</sup> or *Ifngr1*<sup>+/+</sup> bone marrow (five animals per group).  
 (C) IFN- $\gamma$  levels were measured in *Ifngr1*<sup>-/-</sup> and *Ifngr1*<sup>+/+</sup> mice treated by HSCT with *Ifngr1*<sup>+/+</sup> bone marrow, and infected with BCG before or after HSCT (five animals per group).  
 (D) Northern blot analysis of *Ifng* and *GAPDH* mRNA levels in the spleens of *Ifngr1*<sup>-/-</sup> and *Ifngr1*<sup>+/+</sup> mice 30 d after BCG infection (three animals per group). Spleens were removed from animals, directly frozen in 1 ml of Trizol and stored at -80 °C for further preparation. Non-infected *Ifngr1*<sup>-/-</sup>, *Ifngr1*<sup>+/+</sup>, and *Ifng*<sup>-/-</sup> mice were used as controls (two animals per group). Means of arbitrary values obtained after scanning were calculated and normalised with respect to the values obtained for *GAPDH*.  
 doi:10.1371/journal.pmed.0050026.g004

was performed with anti-IFN- $\gamma$  antibody alone. In these conditions IFN- $\gamma$  depletion was not achieved. We thus refined our protocol: animals were infected with BCG and received four intraperitoneal doses of anti-IFN- $\gamma$  plus anti-IL-12 antibodies at two-day intervals, starting from day 14, before transplantation on day 23. Another antibody injection was administered on the day after HSCT. IFN- $\gamma$  depletion was monitored by blood sampling and HSCT was performed with mild conditioning, mimicking that used for human patients (Figure 6A). The injection of specific antibodies into infected *Ifngr1*<sup>-/-</sup> mice kept serum IFN- $\gamma$  levels below 0.5 ng/ml before cell transfer (Figure 6B). The injection of isotype control antibodies resulted in a serum IFN- $\gamma$  concentration of 1.2 ng/ml at the time of HSCT, and ranging from 3 ng/ml to 5 ng/ml


thereafter (Figure 6C). In three of the six animals treated with specific antibodies, serum IFN- $\gamma$  levels did not exceed 3 ng/ml in the first week after HSCT and even decreased to 0.1 ng/ml by nine weeks post-HSCT (Figure 6B). Three other animals behaved like the isotype control-treated group, with serum IFN- $\gamma$  levels remaining high throughout the experiment. HSCT outcome was found to be strictly correlated with serum IFN- $\gamma$  concentration. Chimerism with bone marrow was observed in the three animals in which serum IFN- $\gamma$  concentrations remained low (Figure 6D, left). In contrast, the graft was rejected in the three animals in which serum IFN- $\gamma$  levels were not controlled; no donor cells were found in such animals (Figure 6D, right). Moreover, bacterial disease was cured in the three animals in which HSCT was successful





**Figure 5.** rAAV-IFN- $\gamma$  Injection Promotes Graft Rejection

(A) A rAAV-IFN- $\gamma$  vector was designed, particles were produced and  $5 \times 10^{10}$  physical particles injected into the gastrocnemius of *Ifngr*<sup>-/-</sup> mice. Serum IFN- $\gamma$  levels were determined after rAAV-IFN- $\gamma$  injection in *Ifngr1*<sup>-/-</sup>, *Ifngr1*<sup>+/+</sup> and DKO (*Ifngr1*<sup>-/-</sup>  $\times$  *Ifngr*<sup>-/-</sup>) mice (eight animals per group). (B) *Ifngr1*<sup>-/-</sup>, *Ifngr1*<sup>+/+</sup>, and DKO mice were injected with rAAV-IFN- $\gamma$ , and HSCT was performed with bone marrow from *Ifngr1*<sup>+/+</sup> mice. The percentage chimerism over time was monitored by determining the surface expression of Ly5.1 (donor cells) on lymphocytes (eight animals per group). (C) Infected DKO mice showed no HSCT graft rejection. IFN- $\gamma$  levels were measured following BCG infection, 30 d after infection (eight animals per group). (D) DKO mice were infected with BCG and the percentage chimerism over time was monitored by determining the surface expression of Ly5.1 (donor cells) on lymphocytes (eight animals per group). doi:10.1371/journal.pmed.0050026.g005



**Figure 6.** The Depletion of Circulating IFN- $\gamma$  Is Sufficient to Allow HSCT Engraftment

Groups of six *Ifng*<sup>-/-</sup> mice were infected with BCG and injected with a mixture of specific antibodies against IFN- $\gamma$  and IL-12 or with a control isotype antibody. HSCT was carried out with bone marrow from *Ifng*<sup>+/+</sup> mice and serum IFN- $\gamma$  levels were monitored over time (A). Data are plotted individually for each animal treated with specific antibodies (B) and control isotype antibodies (C). The percentage chimerism was evaluated and representative FACS analyses are shown for engraftment (D), corresponding to three animals with low serum levels of IFN- $\gamma$  (left) and rejection, corresponding to six animals with high serum levels of IFN- $\gamma$  (right).  
doi:10.1371/journal.pmed.0050026.g006

(unpublished data). Our data therefore demonstrate that efficient IFN- $\gamma$  depletion by antibody administration improves engraftment in *Ifngr1*<sup>-/-</sup> mice.

## Discussion

We have shown that the surface expression of functional IFN- $\gamma$ R1 in the haematopoietic compartment alone—actually in only about half of that compartment—is sufficient to protect mice against BCG infection. Similarly, Yap and Sher previously showed that IFN- $\gamma$ R1 expression in the haematopoietic compartment was sufficient to restore resistance to the intracellular macrophage-tropic bacterium *Listeria monocytogenes* in IFN- $\gamma$ R1-deficient mice [37]. However, they also showed that IFN- $\gamma$ R1 expression on both haematopoietic and non-haematopoietic cells was required to confer resistance to the macrophage-tropic intracellular protozoon *Toxoplasma gondii* [37]. Consistent with these findings, Dal Canto and Virgin showed that IFN- $\gamma$  acted directly on both haematopoietic and non-haematopoietic cells during infection with the medial smooth muscle murine tropic  $\gamma$ -herpesvirus-68 [38]. Thus, despite the almost ubiquitous expression of IFN- $\gamma$ R1 on both haematopoietic and non-haematopoietic cells [39–41], BCG infection can be controlled—as attested by bacterial killing, granuloma structure, and animal survival—by the restricted action of IFN- $\gamma$  on the haematopoietic compartment. Results obtained in our murine model are consistent with data from patients with complete IFN- $\gamma$ R1 deficiency undergoing HSCT [12–15] and demonstrate that susceptibility to BCG, and by extension to other mycobacteria, is a haematopoietic disease in persons with IFN- $\gamma$ R1 deficiency. In mice and humans, the extra-haematopoietic expression of IFN- $\gamma$ R1 is redundant for anti-mycobacterial protective immunity.

Likewise, in both mice and humans, the rate of HSCT graft rejection is also very high in individuals lacking IFN- $\gamma$ R1 and infected with mycobacteria. Moreover, high levels of circulating IFN- $\gamma$  account for this high rate of rejection. *Ifngr1*<sup>-/-</sup> mice reject even syngeneic grafts from IFN- $\gamma$ R1-expressing mice, in all conditions resulting in high serum IFN- $\gamma$  levels, including mycobacterial disease and intramuscular injections of rAAV-IFN- $\gamma$ . We have shown that IFN- $\gamma$  is necessary and sufficient for HSCT graft rejection. IFN- $\gamma$  has already been shown to have a direct inhibitory effect on myeloid, erythroid, megakaryocyte, and multipotent colony formation in human cultures in vitro [42,43]. Moreover, in long-term bone marrow cultures—the in vitro assay best mimicking the complex interactions occurring in intact bone marrow—Selleri et al. showed that local IFN- $\gamma$  overproduction by stromal cells inhibits haematopoiesis [44]. Two mechanisms seem to be responsible for the anti-haematopoietic effect of IFN- $\gamma$ : the direct killing of stem cells and the inhibition of cell cycling [44,45]. The exposure of primitive human haematopoietic stem cells to IFN- $\gamma$  in vitro increases Fas antigen expression, thereby rendering the cells more susceptible to apoptosis [46]. In vivo studies in mouse models have confirmed these results. Transgenic mice expressing multiple copies of the IFN- $\gamma$  gene, leading to high levels of IFN- $\gamma$  production in the bone marrow and thymus, display hypocellularity, and multiple alterations of the immune system [47]. There is also a strong correlation between the degree of haematopoietic suppression and the level of IFN- $\gamma$  in vitro [48,49]. Altogether, these observations most likely

account for the detrimental effects of high levels of circulating IFN- $\gamma$  in IFN- $\gamma$ R1-deficient mice and humans undergoing HSCT.

The negative impact of high IFN- $\gamma$  levels, leading to HSCT graft rejection, has important clinical implications for the treatment of people with complete IFN- $\gamma$ R1 deficiency. Attempts could be made to decrease levels of circulating IFN- $\gamma$  before HSCT. As recently shown, antibiotics may not eradicate *M. fortuitum* infection, but they do markedly decrease serum IFN- $\gamma$  concentration before HSCT [15]. Prolonged treatment with multiple anti-mycobacterial antibiotics is therefore required, but is unlikely to decrease IFN- $\gamma$  levels sufficiently and is unlikely to cure most of the patients [7]. In this study, we explored the use of specific antibodies to deplete the blood of IFN- $\gamma$ . As anti-IFN- $\gamma$  antibodies alone were not sufficient for IFN- $\gamma$  depletion, we also used anti-IL-12p40 antibodies [50] to achieve efficient depletion of IFN- $\gamma$  in the serum. Infected IFN- $\gamma$ R1-deficient mice treated in this way displayed engraftment and control of the mycobacterial infection after HSCT. Anti-IL-12 and anti-IFN- $\gamma$  antibodies have already been used to treat human immune diseases, such as psoriasis and Crohn's disease [51–53]. The clinical use of these antibodies before HSCT is therefore feasible in IFN- $\gamma$ R1-deficient patients. Our study paves the way for improvements in transplantation conditions in people with IFN- $\gamma$ R1 deficiency, or with other diseases associated with elevated levels of circulating IFN- $\gamma$ , such as IFN- $\gamma$ R2 deficiency [54]. We can also speculate that the use of such antibodies may also reduce the risks of HSCT rejection, improving engraftment, in other settings, such as HSCT from a haplo-identical or matched unrelated donor [16,55]. Although we did not observe any detrimental effect of rAAV-IFN- $\gamma$  in an HLA-identical setting in our mouse model, the depletion of IFN- $\gamma$  in patients with conditions associated with high rates of graft rejection, such as HLA-II deficiency [56], or in other patients undergoing haplo-identical transplantations, might be beneficial. Further experimental studies in mice should be done before the corresponding clinical trials can be undertaken.

## Acknowledgments

We would like to thank Anne-Marie Douar from Genethon, Evry for rAAV production. We also thank Joachim Roesler for personal communications concerning patients and Alain Fischer for initially suggesting the potential role of IFN- $\gamma$  in graft rejection. We would like to thank all members of the Laboratory of Human Genetics of Infectious Diseases for helpful discussions.

**Author contributions.** MR, CS, JLG, and JLC designed the study and contributed to the writing. MR and CS performed the experiments. MR, CS, GV, LR, JFE, HD, JLG, and JLC analyzed the data. LR provided the antibody and contributed to the design of the cytokine neutralisation experiment. JFE did the histology analysis.

## References

1. Hamosh A, Scott AF, Amberger JS, Bocchini CA, McKusick VA (2005) Online Mendelian Inheritance in Man (OMIM), a knowledgebase of human genes and genetic disorders. *Nucleic Acids Res* 33: D514–D517.
2. Casanova JL, Abel L (2002) Genetic dissection of immunity to mycobacteria: the human model. *Annu Rev Immunol* 20: 581–620.
3. Filipe-Santos O, Bustamante J, Chappier A, Vogt G, de Beaucoudrey L, et al. (2006) Inborn errors of IL-12/23- and IFN-gamma-mediated immunity: molecular, cellular, and clinical features. *Semin Immunol* 18: 347–361.
4. Jouanguy E, Altare F, Lamhamedi S, Revy P, Emile JF, et al. (1996) Interferon-gamma-receptor deficiency in an infant with fatal bacille Calmette-Guerin infection. *N Engl J Med* 335: 1956–1961.
5. Newport MJ, Huxley CM, Huston S, Hawrylowicz CM, Oostra BA, et al. (1996) A mutation in the interferon-gamma-receptor gene and susceptibility to mycobacterial infection. *N Engl J Med* 335: 1941–1949.
6. Jouanguy E, Dupuis S, Pallier A, Doffinger R, Fondaneche MC, et al. (2000)

- In a novel form of IFN-gamma receptor 1 deficiency, cell surface receptors fail to bind IFN-gamma. *J Clin Invest* 105: 1429–1436.
7. Dorman SE, Picard C, Lammas D, Heyne K, van Dissel JT, et al. (2004) Clinical features of dominant and recessive interferon gamma receptor 1 deficiencies. *Lancet* 364: 2113–2121.
  8. Alcais A, Fieschi C, Abel L, Casanova JL (2005) Tuberculosis in children and adults: two distinct genetic diseases. *J Exp Med* 202: 1617–1621.
  9. MacLennan C, Fieschi C, Lammas D, Picard C, Dorman SE, et al. (2004) Interleukin (IL)-12 and IL-23 are key cytokines for immunity against *Salmonella* in humans. *J Infect Dis* 190: 1755–1757.
  10. Roesler J, Kofink B, Wendisch J, Heyden S, Paul D, et al. (1999) *Listeria monocytogenes* and recurrent mycobacterial infections in a child with complete interferon-gamma-receptor (IFN $\gamma$ R1) deficiency: mutational analysis and evaluation of therapeutic options. *Exp Hematol* 27: 1368–1374.
  11. Emile JF, Patey N, Altare F, Lamhamedi S, Jouanguy E, et al. (1997) Correlation of granuloma structure with clinical outcome defines two types of idiopathic disseminated BCG infection. *J Pathol* 181: 25–30.
  12. Reuter U, Roesler J, Thiede C, Schulz A, Classen CF, et al. (2002) Correction of complete interferon-gamma receptor 1 deficiency by bone marrow transplantation. *Blood* 100: 4234–4235.
  13. Roesler J, Horwitz ME, Picard C, Bordigoni P, Davies G, et al. (2004) Hematopoietic stem cell transplantation for complete IFN-gamma receptor 1 deficiency: a multi-institutional survey. *J Pediatr* 145: 806–812.
  14. Horwitz ME, Uzel G, Linton GF, Miller JA, Brown MR, et al. (2003) Persistent *Mycobacterium avium* infection following nonmyeloablative allogeneic peripheral blood stem cell transplantation for interferon-gamma receptor-1 deficiency. *Blood* 102: 2692–2694.
  15. Chantrain CF, Bruwier A, Richard B, Largent V, Chapgier A, et al. (2006) Successful hematopoietic stem cell transplantation in a child with active disseminated *Mycobacterium fortuitum* infection and interferon-gamma receptor 1 deficiency. *Bone Marrow Transplant* 38: 75–76.
  16. Antoine C, Muller S, Cant A, Cavazzana-Calvo M, Veys P, et al. (2003) Long-term survival and transplantation of haemopoietic stem cells for immunodeficiencies: report of the European experience 1968–99. *Lancet* 361: 553–560.
  17. Huang S, Hendriks W, Althage A, Hemmi S, Bluethmann H, et al. (1993) Immune response in mice that lack the interferon-gamma receptor. *Science* 259: 1742–1745.
  18. Dalton DK, Pitts-Meek S, Keshav S, Figari IS, Bradley A, et al. (1993) Multiple defects of immune cell function in mice with disrupted interferon-gamma genes. *Science* 259: 1739–1742.
  19. Swihart K, Fruth U, Messmer N, Hug K, Behin R, et al. (1995) Mice from a genetically resistant background lacking the interferon gamma receptor are susceptible to infection with *Leishmania major* but mount a polarized T helper cell 1-type CD4+ T cell response. *J Exp Med* 181: 961–971.
  20. Zhao YX, Tarkowski A (1995) Impact of interferon-gamma receptor deficiency on experimental *Staphylococcus aureus* septicemia and arthritis. *J Immunol* 155: 5736–5742.
  21. Mastroeni P, Clare S, Khan S, Harrison JA, Hormaeche CE, et al. (1999) Interleukin 18 contributes to host resistance and gamma interferon production in mice infected with virulent *Salmonella typhimurium*. *Infect Immun* 67: 478–483.
  22. Rijneveld AW, Lauw FN, Schultz MJ, Florquin S, Te Velde AA, et al. (2002) The role of interferon-gamma in murine pneumococcal pneumonia. *J Infect Dis* 185: 91–97.
  23. Cantin E, Tanamachi B, Openshaw H (1999) Role for gamma interferon in control of herpes simplex virus type 1 reactivation. *J Virol* 73: 3418–3423.
  24. Flynn JL, Chan J, Triebold KJ, Dalton DK, Stewart TA, et al. (1993) An essential role for interferon gamma in resistance to *Mycobacterium tuberculosis* infection. *J Exp Med* 178: 2249–2254.
  25. Garcia I, Miyazaki Y, Marchal G, Lesslauer W, Vassalli P (1997) High sensitivity of transgenic mice expressing soluble TNFR1 fusion protein to mycobacterial infections: synergistic action of TNF and IFN-gamma in the differentiation of protective granulomas. *Eur J Immunol* 27: 3182–3190.
  26. Doherty TM, Sher A (1997) Defects in cell-mediated immunity affect chronic, but not innate, resistance of mice to *Mycobacterium avium* infection. *J Immunol* 158: 4822–4831.
  27. Kamijo R, Le J, Shapiro D, Havell EA, Huang S, et al. (1993) Mice that lack the interferon-gamma receptor have profoundly altered responses to infection with *Bacillus Calmette-Guérin* and subsequent challenge with lipopolysaccharide. *J Exp Med* 178: 1435–1440.
  28. Gheorghui M, Augier J, Lagrange P (1983) Maintenance and control of the French BCG strain 1173-P2 (primary and secondary seeding lots). *Bull Inst Pasteur* 81: 281–288.
  29. Ehlers S, Richter E (2001) Differential requirement for interferon-gamma to restrict the growth of or eliminate some recently identified species of nontuberculous mycobacteria in vivo. *Clin Exp Immunol* 124: 229–238.
  30. Douar AM, Poulard K, Stockholm D, Danos O (2001) Intracellular trafficking of adeno-associated virus vectors: routing to the late endosomal compartment and proteasome degradation. *J Virol* 75: 1824–1833.
  31. Snyder RO, Spratt SK, Lagarde C, Bohl D, Kaspar B, et al. (1997) Efficient and stable adeno-associated virus-mediated transduction in the skeletal muscle of adult immunocompetent mice. *Hum Gene Ther* 8: 1891–1900.
  32. Cherwinski HM, Schumacher JH, Brown KD, Mosmann TR (1987) Two types of mouse helper T cell clone. III. Further differences in lymphokine synthesis between Th1 and Th2 clones revealed by RNA hybridization, functionally monospecific bioassays, and monoclonal antibodies. *J Exp Med* 166: 1229–1244.
  33. Wysocka M, Kubin M, Vieira LQ, Ozmen L, Garotta G, et al. (1995) Interleukin-12 is required for interferon-gamma production and lethality in lipopolysaccharide-induced shock in mice. *Eur J Immunol* 25: 672–676.
  34. Belnoue E, Costa FT, Frankenberg T, Vigarito AM, Voza T, et al. (2004) Protective T cell immunity against malaria liver stage after vaccination with live sporozoites under chloroquine treatment. *J Immunol* 172: 2487–2495.
  35. Fieschi C, Dupuis S, Picard C, Smith CI, Holland SM, et al. (2001) High levels of interferon gamma in the plasma of children with complete interferon gamma receptor deficiency. *Pediatrics* 107: E48.
  36. Kamijo R, Gercitano J, Shapiro D, Green SJ, Aguet M, et al. (1995) Generation of nitric oxide and clearance of interferon-gamma after BCG infection are impaired in mice that lack the interferon-gamma receptor. *J Inflamm* 46: 23–31.
  37. Yap GS, Sher A (1999) Effector cells of both nonhemopoietic and hemopoietic origin are required for interferon (IFN)-gamma- and tumor necrosis factor (TNF)-alpha-dependent host resistance to the intracellular pathogen, *Toxoplasma gondii*. *J Exp Med* 189: 1083–1092.
  38. Dal Canto AJ, Swanson PE, O'Guin AK, Speck SH, Virgin HW (2001) IFN-gamma action in the media of the great elastic arteries, a novel immunoprivileged site. *J Clin Invest* 107: R15–22.
  39. Celada A (1988) The interferon gamma receptor. *Lymphokine Res* 7: 61–73.
  40. Valente G, Ozmen L, Novelli F, Geuna M, Palestro G, et al. (1992) Distribution of interferon-gamma receptor in human tissues. *Eur J Immunol* 22: 2403–2412.
  41. Farrar MA, Schreiber RD (1993) The molecular cell biology of interferon-gamma and its receptor. *Annu Rev Immunol* 11: 571–611.
  42. Broxmeyer HE, Williams DE, Lu L, Cooper S, Anderson SL, et al. (1986) The suppressive influences of human tumor necrosis factors on bone marrow hematopoietic progenitor cells from normal donors and patients with leukemia: synergism of tumor necrosis factor and interferon-gamma. *J Immunol* 136: 4487–4495.
  43. Raefsky EL, Platanius LC, Zoumbos NC, Young NS (1985) Studies of interferon as a regulator of hematopoietic cell proliferation. *J Immunol* 135: 2507–2512.
  44. Selleri C, Maciejewski JP, Sato T, Young NS (1996) Interferon-gamma constitutively expressed in the stromal microenvironment of human marrow cultures mediates potent hematopoietic inhibition. *Blood* 87: 4149–4157.
  45. Selleri C, Sato T, Anderson S, Young NS, Maciejewski JP (1995) Interferon-gamma and tumor necrosis factor-alpha suppress both early and late stages of hematopoiesis and induces programmed cell death. *J Cell Physiol* 165: 538–546.
  46. Maciejewski JP, Selleri C, Anderson S, Young NS (1995) Fas antigen expression on CD34+ human marrow cells is induced by interferon-gamma and tumor necrosis factor-alpha and potentiates cytokine-mediated hematopoietic suppression in vitro. *Blood* 85: 3183–3190.
  47. Young HA, Klinman DM, Reynolds DA, Grzegorzewski KJ, Nii A, et al. (1997) Bone marrow and thymus expression of interferon-gamma results in severe B-cell lineage reduction, T-cell lineage alterations, and hematopoietic progenitor deficiencies. *Blood* 89: 583–595.
  48. Zoumbos NC, Djeu JY, Young NS (1984) Interferon is the suppressor of hematopoiesis generated by stimulated lymphocytes in vitro. *J Immunol* 133: 769–774.
  49. Angulo I, Rodriguez R, Garcia B, Medina M, Navarro J, et al. (1995) Involvement of nitric oxide in bone marrow-derived natural suppressor activity; its dependence on IFN $\gamma$ . *J Immunol* 155: 15–26.
  50. Toichi E, Torres G, McCormick TS, Chang T, Mascelli MA, et al. (2006) An anti-IL-12p40 antibody down-regulates type 1 cytokines, chemokines, and IL-12/23 in psoriasis. *J Immunol* 177: 4917–4926.
  51. Kaufmann CL, Aria N, Toichi E, McCormick TS, Cooper KD, et al. (2004) A phase I study evaluating the safety, pharmacokinetics, and clinical response of a human IL-12p40 antibody in subjects with plaque psoriasis. *J Invest Dermatol* 123: 1037–1044.
  52. Mannon PJ, Fuss IJ, Mayer L, Elson CO, Sandborn WJ, et al. (2004) Anti-interleukin-12 antibody for active Crohn's disease. *N Engl J Med* 351: 2069–2079.
  53. Skurkovich B, Skurkovich S (2003) Anti-interferon-gamma antibodies in the treatment of autoimmune diseases. *Curr Opin Mol Ther* 5: 52–57.
  54. Vogt G, Chapgier A, Yang K, Chuzhanova N, Feinberg J, et al. (2005) Gains of glycosylation comprise an unexpectedly large group of pathogenic mutations. *Nat Genet* 37: 692–700.
  55. Ruggeri L, Aversa F, Martelli MF, Velardi A (2006) Allogeneic hematopoietic transplantation and natural killer cell recognition of missing self. *Immunological Reviews* 214: 202–218.
  56. Renella R, Picard C, Neven B, Ouachée-Chardin M, Casanova JL, et al. (2006) Human leucocyte antigen-identical haematopoietic stem cell transplantation in major histocompatibility complex class II immunodeficiency: reduced survival correlates with an increased incidence of acute graft-versus-host disease and pre-existing viral infections. *Br J Haematol* 136: 510–516.

## Editors' Summary

**Background.** Normally, the body's immune system efficiently recognizes and kills bacteria and viruses, but in some rare inherited disorders ("primary immunodeficiencies") part of the immune system works poorly or is missing. This leaves affected individuals susceptible to infections. People with one of these disorders—interferon-gamma receptor 1 (IFN- $\gamma$ R1) deficiency—are very susceptible to infections with mycobacteria. Except for *Mycobacterium tuberculosis* and *M. leprae* (which cause tuberculosis and leprosy, respectively), mycobacteria rarely cause human disease. However, most people with IFN- $\gamma$ R1 deficiency die during childhood from multiple, widespread mycobacterial infections, because IFN- $\gamma$ R1 deficiency disables a specific part of their immune system. When most bacteria enter the body, immune system cells called macrophages engulf and kill them, but mycobacteria actually multiply inside macrophages. This infection stimulates lymphocytes and other immune system cells to release IFN- $\gamma$ , which binds to IFN- $\gamma$ R1 on uninfected macrophages, activates them, and recruits them to the infection site. Here, they form a "granuloma," a mass of macrophages and activated lymphocytes that "walls off" the infection. Granuloma formation does not occur in patients with IFN- $\gamma$ R1 deficiency, so mycobacteria (including the usually benign tuberculosis vaccination strain *M. bovis* BCG) spread throughout the body with disastrous consequences.

**Why Was This Study Done?** The only effective treatment for patients with IFN- $\gamma$ R1 deficiency is hematopoietic stem cell transplantation (HSCT). HSCs are the source of all the immune system cells, so transplantation of HSCs from a donor with a normal *IFNGR1* gene can provide a patient who has IFN- $\gamma$ R1 deficiency with a new immune system that can combat mycobacterial infections. Unfortunately, in this particular immune deficiency, the new HSCs cannot engraft, even when the patient's own immune system is disabled before HSCT by intensive chemotherapy, and when the donor cells come from a close relative and are a good immunological match. In this study, the researchers have investigated why rejection is so common in IFN- $\gamma$ R1 deficiency using a mouse strain called C57BL/6 *Ifngr1*<sup>-/-</sup>—C57BL/6 denotes the genetic background of these mice and *Ifngr1*<sup>-/-</sup> indicates that, like human patients, these mice make no IFN- $\gamma$ R1.

**What Did the Researchers Do and Find?** *Ifngr1*<sup>-/-</sup> mice, the researchers report, cannot control *M. bovis* BCG infections and do not form mature granulomas just like human patients with IFN- $\gamma$ R1 deficiency. Wild-type C57BL/6 (*Ifngr1*<sup>+/+</sup>) mice, however, rapidly control *M. bovis* BCG infections and form mature granulomas. *Ifngr1*<sup>+/+</sup> HSC transplanted into mycobacteria-free *Ifngr1*<sup>-/-</sup> mice survived well and protected the mice against

later mycobacterial challenge but *Ifngr1*<sup>-/-</sup> mice infected with *M. bovis* BCG before HSCT rejected the transplanted HSCs. Mycobacteria-infected *Ifngr1*<sup>-/-</sup> mice and human patients with IFN- $\gamma$ R1 deficiency have blood high levels of IFN- $\gamma$ . Could this be responsible for HSCT rejection? To find out, the researchers expressed IFN- $\gamma$  in uninfected *Ifngr1*<sup>-/-</sup> mice before HSCT. As in infected mice, these grafts failed. Conversely, transplanted HSCs survived when transplanted into *Ifngr1*<sup>-/-</sup> mice that had been genetically altered to express no IFN- $\gamma$ , even when these mice were infected with *M. bovis* BCG before transplantation. Finally, when the researchers used antibodies (proteins made by the immune system that recognize specific molecules) to remove circulating IFN- $\gamma$  from infected *Ifngr1*<sup>-/-</sup> mice, HSCT worked well in the animals with the lowest IFN- $\gamma$  levels.

**What Do These Findings Mean?** These findings indicate that in a mouse model of IFN- $\gamma$ R1 deficiency, high circulating IFN- $\gamma$  concentrations drive the rejection of transplanted HSCs and prevent the development of antimycobacterial immunity, probably by directly killing the transplanted cells and/or stopping them multiplying. They also suggest how HSCT could be improved in patients with IFN- $\gamma$ R1 deficiency although, as with all animal studies, the situation in people might turn out to be very different. Importantly, antibodies that reduce circulating IFN- $\gamma$  are already being used to treat other human immune diseases, so the clinical use of these antibodies in patients with IFN- $\gamma$  deficiency before HSCT is feasible. Finally, the researchers speculate that the use of IFN- $\gamma$ -depleting antibodies might be beneficial in other situations where HSCT often fails such as when a close relative is not available as a donor. However, this possibility will need to be thoroughly tested in mice before human clinical trials can be started.

**Additional Information.** Please access these Web sites via the online version of this summary at <http://dx.doi.org/10.1371/journal.pmed.0050026>.

- General information about primary immunodeficiencies is available from the US National Institute of Child Health and Human Development
- Online Mendelian Inheritance in Man (OMIM) provides information about familial predisposition to mycobacterial disease
- Wikipedia has pages on hematopoietic stem cell transplantation and on interferon- $\gamma$  (note that Wikipedia is a free online encyclopedia that anyone can edit; available in several languages)
- The Human Genetics of Infectious Diseases Lab focuses on the genetic basis of predisposition or resistance to infectious diseases in humans

