

## Longitudinal study assessing the return of chloroquine susceptibility of *Plasmodium falciparum* isolates from travelers returning from West Africa

Myriam Gharbi, Jennifer Flegg, Véronique Hubert, Eric Kendjo, Jessica Metcalf, Philippe Guerin, Jacques Le Bras

### ► To cite this version:

Myriam Gharbi, Jennifer Flegg, Véronique Hubert, Eric Kendjo, Jessica Metcalf, et al.. Longitudinal study assessing the return of chloroquine susceptibility of *Plasmodium falciparum* isolates from travelers returning from West Africa. Challenges in malaria research, Oct 2012, Basel, Switzerland. BMC, 11 (Suppl 1), pp.P38, 2012, <10.1186/1475-2875-11-S1-P38>. <inserm-00742701>

**HAL Id: inserm-00742701**

**<http://www.hal.inserm.fr/inserm-00742701>**

Submitted on 17 Oct 2012

**HAL** is a multi-disciplinary open access archive for the deposit and dissemination of scientific research documents, whether they are published or not. The documents may come from teaching and research institutions in France or abroad, or from public or private research centers.

L'archive ouverte pluridisciplinaire **HAL**, est destinée au dépôt et à la diffusion de documents scientifiques de niveau recherche, publiés ou non, émanant des établissements d'enseignement et de recherche français ou étrangers, des laboratoires publics ou privés.

POSTER PRESENTATION

Open Access

# Longitudinal study assessing the return of chloroquine susceptibility of *Plasmodium falciparum* isolates from travelers returning from West Africa

Myriam Gharbi<sup>1,2,3</sup>, Jennifer Flegg<sup>4,5</sup>, Véronique Hubert<sup>6,7</sup>, Eric Kendjo<sup>6,8</sup>, Jessica Metcalf<sup>9</sup>, Philippe J Guerin<sup>3,4,5,10\*</sup>, Jacques Le Bras<sup>1,2,6,7</sup>

From Challenges in malaria research  
Basel, Switzerland. 10-12 October 2012

## Background

From the 1940s up to the 1990s, chloroquine (CQ) was the main malaria therapy worldwide. Following the CQ resistance burden in Africa, most African countries have discontinued CQ during the past 2 decades, and now promote artemisinin-based combination therapy (ACT), as the first-line treatment for uncomplicated malaria. The policy changed in West Africa during the last decade (2002 in Cameroon; 2003 in Senegal and Cote d'Ivoire; 2004 in Mali). The aim of this study is to describe the evolution of CQ resistance in West Africa, through travellers returning from this region.

## Methods

The study was conducted by the Malaria National Reference Centre, France. The database collated *in vitro* response of reference and clinical isolates for CQ and the *pfcrK76* molecular marker for CQ susceptible *Pf* malaria from travellers returning from Cameroon, Senegal, Cote d'Ivoire and Mali. As a proxy of drug pressure, CQ intake for children under five years of age with fever was extracted from the Demographic Health Surveys (DHS) and Multiple Indicator Cluster Surveys (MICS) for the study period [1,2]. Logistic regression models were used to detect trends in the susceptible isolates proportions.

## Results

From 2000 to 2011, around 700 isolates were genotyped for each country. The frequency of the *pfcrK76* wild-type

significantly increased for Cameroon (CM) (from 10% to 41%, Slope=0.09,  $p<10^{-3}$ ), Cote d'Ivoire (CI) (from 37% to 63%, Slope = 0.14,  $p<10^{-3}$ ), and Senegal (SN) (from 22% to 53%, Slope=0.17,  $p<10^{-3}$ ). The geometric mean of the 50% growth inhibition (IC<sub>50</sub>) of CQ decreased from 181 nM (95% confidence interval, 87-374) (25% CQ sensitive) to 51 nM (37-71) (63% CQ sensitive) in CM, from 75nM (43-130) (41% CQ sensitive) to 29nM (22-39) (84% CQ sensitive) in CI and from 86nM (51-145) (41% CQ sensitive) to 39nM (26-60) (75% CQ sensitive) in SN. Analyses performed from 2004 to 2011, when most of West African countries have officially discontinued CQ, confirmed previous results and also show a significant increase of the prevalence of *pfcrK76* wild-type genotype for Mali (ML) (Slope = 0.07,  $p=0.02$ ). Meanwhile, CQ use among children with fever significantly decreased during this period.

## Conclusions

An increase of CQ susceptibility following official withdrawal is observed in travellers returning from Cameroon, Cote d'Ivoire, Mali and Senegal. The length of time between policy changes and their subsequent implementation, as well as the cross resistance between antimalarial drugs, may affect the time for a significant recovery of CQ sensitivity. This information should be compared to country level CQ efficacy data.

## Acknowledgements

Data for DHS was obtained from Macro, Inc, <http://www.measuredhs.org>. Data for MICS was obtained from UNICEF, <http://www.childinfo.org>. This study was supported in part by a grant for doctoral studies to M. Gharbi

<sup>3</sup>Ecole des Hautes Etudes en Santé Publique (EHESP), Rennes, France  
Full list of author information is available at the end of the article

from the Doctoral Network of the École des Hautes Etudes en Santé Publique, Rennes, France. This abstract is being presented on behalf of the French National Reference Center for Imported Malaria study group.

#### Author details

<sup>1</sup>UMR216: Mère et enfant face aux infections tropicales, Université Paris Descartes-Paris 5 and Institut de Recherche pour le Développement (IRD), Paris, France. <sup>2</sup>WorldWide Antimalarial Resistance Network (WWARN), Paris, France. <sup>3</sup>Ecole des Hautes Etudes en Santé Publique (EHESP), Rennes, France. <sup>4</sup>WorldWide Antimalarial Resistance Network (WWARN), Oxford, UK. <sup>5</sup>Centre for tropical Medicine, Nuffield Department of Clinical Medicine, CCVTM, University of Oxford, Oxford, UK. <sup>6</sup>Centre National de Référence du Paludisme, Paris, France. <sup>7</sup>Service de Parasitologie Mycologie, CHU Bichat-Claude Bernard, APHP, Paris, France. <sup>8</sup>Service de Parasitologie Mycologie, CHU Pitié-Salpêtrière, APHP, Paris, France. <sup>9</sup>Epidemiology and Infectious Diseases, Department of Zoology, University of Oxford, Oxford, UK. <sup>10</sup>UMR S 707: Epidemiology, Information Systems, Modeling, INSERM and Université Pierre et Marie-Curie-Paris, Paris, France.

Published: 15 October 2012

#### References

1. MeasureDHS, ICF Macro: [<http://www.measuredhs.com>].
2. Multiple Indicator Cluster Survey Manual, United Nations Children's Fund: [<http://www.childinfo.org>].

doi:10.1186/1475-2875-11-S1-P38

**Cite this article as:** Gharbi *et al.*: Longitudinal study assessing the return of chloroquine susceptibility of *Plasmodium falciparum* isolates from travelers returning from West Africa. *Malaria Journal* 2012 **11**(Suppl 1): P38.

**Submit your next manuscript to BioMed Central  
and take full advantage of:**

- Convenient online submission
- Thorough peer review
- No space constraints or color figure charges
- Immediate publication on acceptance
- Inclusion in PubMed, CAS, Scopus and Google Scholar
- Research which is freely available for redistribution

Submit your manuscript at  
[www.biomedcentral.com/submit](http://www.biomedcentral.com/submit)

