



HAL
open science

Altered innate functions of myeloid dendritic cells in ANCA-associated vasculitis

Cécile Braudeau, Antoine Néel, Marie Rimbart, Mohamed Hamidou, Régis Josien

► **To cite this version:**

Cécile Braudeau, Antoine Néel, Marie Rimbart, Mohamed Hamidou, Régis Josien. Altered innate functions of myeloid dendritic cells in ANCA-associated vasculitis. 7th European Workshop on Immune-Mediated Inflammatory Diseases, Nov 2012, Noordwijk aan Zee, Netherlands. pp.P20. inserm-00758219

HAL Id: inserm-00758219

<https://www.hal.inserm.fr/inserm-00758219>

Submitted on 28 Nov 2012

HAL is a multi-disciplinary open access archive for the deposit and dissemination of scientific research documents, whether they are published or not. The documents may come from teaching and research institutions in France or abroad, or from public or private research centers.

L'archive ouverte pluridisciplinaire **HAL**, est destinée au dépôt et à la diffusion de documents scientifiques de niveau recherche, publiés ou non, émanant des établissements d'enseignement et de recherche français ou étrangers, des laboratoires publics ou privés.



POSTER PRESENTATION

Open Access

Altered innate functions of myeloid dendritic cells in ANCA-associated vasculitis

Cécile Braudeau^{1,2}, Antoine Néel^{2,3}, Marie Rimbart^{1,2}, Mohamed Hamidou^{2,4}, Régis Josien^{1,2,4*}

From 7th European Workshop on Immune-Mediated Inflammatory Diseases
Noordwijk aan Zee, the Netherlands. 28-30 November 2012

Background

Dendritic cells (DC) are critical effectors of innate and adaptive immunity, acting both as sentinels that detect the presence of pathogens and as key antigen-presenting cells that regulate the adaptive immune response. Therefore, DC play a crucial role in the control of autoimmune responses. We previously showed that blood DC numbers were strongly reduced in ANCA-associated vasculitis (AAV) likely due to their recruitment in tissues. Here, we assessed the ex vivo responsiveness of blood DC from AAV-patients to Toll-like receptors (TLRs) stimulation.

Materials and methods

Blood samples from 10 untreated patients with AAV during flares and before any immunosuppressive treatment (AP) were analyzed, along with 9 AAV patients in remission (RP) and 11 age-matched healthy controls (HC). Intracellular cytokine (IL-12, TNF- α , IFN- α) production by blood DC was assessed by 8-colors flow cytometry after stimulation by Toll-like receptors of whole blood samples.

Results

We found that myeloid DC (mDC) from patients in acute phase exhibited a decreased IL-12 production after TLR3, 4 and 7/8 stimulation compared to patients in remission and healthy controls. These mDC also produced less TNF- α after TLR3 stimulation. Moreover, we observed a reduction in the frequency TNF α -producing plasmacytoid DC (pDC) upon TLR7/8 triggering in AP patients compared to RP patients and HC.

Conclusion

Our data show that circulating mDC from patients with AAV exhibited an altered response to several TLR ligands, with a notable decrease in IL-12 production.

These unexpected results suggest the innate functions of DC especially in response to pathogens are impaired during AAV.

Author details

¹CHU Nantes, Laboratoire d'Immunologie, Nantes, France. ²INSERM Center of Research in Transplantation and Immunology, UMR 1064, Nantes, France. ³CHU Nantes, Service de Médecine Interne, Nantes, France. ⁴Université de Nantes, Faculté de Médecine, Nantes, France.

Published: 28 November 2012

doi:10.1186/1479-5876-10-S3-P20

Cite this article as: Braudeau et al.: Altered innate functions of myeloid dendritic cells in ANCA-associated vasculitis. *Journal of Translational Medicine* 2012 **10**(Suppl 3):P20.

Submit your next manuscript to BioMed Central
and take full advantage of:

- Convenient online submission
- Thorough peer review
- No space constraints or color figure charges
- Immediate publication on acceptance
- Inclusion in PubMed, CAS, Scopus and Google Scholar
- Research which is freely available for redistribution

Submit your manuscript at
www.biomedcentral.com/submit



¹CHU Nantes, Laboratoire d'Immunologie, Nantes, France
Full list of author information is available at the end of the article